

AN ELLIPTIC SURFACE COVERED BY MUMFORD'S FAKE PROJECTIVE PLANE

Dedicated to Professor Masayoshi Nagata on his sixtieth birthday

MASA-NORI ISHIDA

(Received January 28, 1987)

Introduction. In [Mum], Mumford constructed an algebraic surface M of general type with $K_M^2 = 9$ and $p_g = q = 0$. This surface is called Mumford's fake projective plane because it has the same Betti numbers as the complex projective plane (see [BPV, Historical Note]). No other example of fake projective planes in this sense seems to be known up to now.

Since $c_1^2(M) = 3c_2(M) = 9$, the universal covering space of the complex surface M is isomorphic to the unit ball in C^2 by Yau's result. However, Mumford's surface is constructed by means of the theory of the p -adic unit ball by Kurihara [Ku] and Mustafin [Mus]. By the construction of M , there exists an unramified Galois covering $V \rightarrow M$ of order eight. More precisely, a simple group G of order 168 acts on V , and M is the quotient of V by a 2-Sylow subgroup of G .

In this paper, we study the quotient surface $Y = V/G$. Since the action has fixed points, Y has some singular points. We prove that the minimal desingularization \tilde{Y} of Y is an elliptic surface. We also determine the types of the singular fibers of the elliptic fibration.

Mumford's surface M is given as a Z_2 -scheme. Hence it has a modulo 2 reduction M_0 . The normalization \tilde{M}_0 of M_0 is the blowing-up of $P_{F_2}^2$ at the seven F_2 -rational points. In Section 1, we describe explicitly how to recover M_0 from \tilde{M}_0 .

The author expresses his thanks to Professors F. Hirzebruch and I. Nakamura for their interest and suggestion on this work. Some results and techniques in Sections 3 and 4 are due to Nakamura in unpublished notes.

NOTATION. Let X be a scheme over an affine scheme $\text{Spec } A$. When a ring homomorphism $A \rightarrow B$ is given, we denote by X_B the fiber product $X \times_{\text{Spec } A} \text{Spec } B$ and by $X(B)$ the set of B -valued points of X . If X is of finite type and B is an algebraically closed field, then we sometimes treat $X(B)$ as a variety.

1. **The closed fiber of Mumford's surface.** We will recall some notation in Mumford's paper [Mum].

We always restrict ourselves to the case of the base ring \mathbf{Z}_2 . Hence the maximal ideal is generated by 2, and the quotient field is the 2-adic number field \mathbf{Q}_2 . We denote by η and 0 the generic point and the closed point of $\text{Spec } \mathbf{Z}_2$, respectively.

A matrix $\alpha = (a_{i,j})_{i,j=0,1,2} \in GL(3, \mathbf{Q}_2)$ defines a linear automorphism of the vector space $\mathbf{Q}_2 X_0 + \mathbf{Q}_2 X_1 + \mathbf{Q}_2 X_2$ with indeterminates X_0, X_1, X_2 by

$$\alpha(c_0 X_0 + c_1 X_1 + c_2 X_2) = (X_0, X_1, X_2) \alpha^t(c_0, c_1, c_2) = \sum_i (\sum_j a_{i,j} c_j) X_i .$$

Hence the induced automorphism α^\wedge of $P_{\mathbf{Q}_2}^2 = \text{Proj } \mathbf{Q}_2[X_0, X_1, X_2]$ is given in terms of the homogeneous coordinates $(X_0 : X_1 : X_2)$ by

$$\alpha^\wedge(X_0 : X_1 : X_2) = (X_0 : X_1 : X_2) \alpha .$$

Thus the composite $\beta^\wedge \circ \alpha^\wedge$ is equal to $(\alpha\beta)^\wedge$.

The \mathbf{Z}_2 -scheme \mathcal{X} of Kurihara and Mustafin is defined as follows:

Let $P_{\mathbf{F}_2}^2$ be the projective plane with the homogeneous coordinates $(X_0 : X_1 : X_2)$. The closed fiber $P_{\mathbf{F}_2}^2$ has seven \mathbf{F}_2 -rational points and seven \mathbf{F}_2 -rational lines. We first blow up $P_{\mathbf{F}_2}^2$ at these seven \mathbf{F}_2 -rational points, and then blow up the resulting surface further along the proper transform of the union of the seven \mathbf{F}_2 -rational lines. Let U be the union of the generic fiber $P_{\mathbf{Q}_2}^2$ and a sufficiently small open neighborhood of the proper transform of $P_{\mathbf{F}_2}^2$ in the blown-up scheme. For each α in $GL(3, \mathbf{Q}_2)$ we denote by U^α the \mathbf{Z}_2 -scheme such that the generic fiber is equal to $P_{\mathbf{Q}_2}^2$ and that there exists an isomorphism $U \simeq U^\alpha$ which induces α^\wedge on the generic fiber. Then the union $\cup_\alpha U^\alpha$ over all α in $GL(3, \mathbf{Q}_2)$ is patched together to a regular scheme \mathcal{X} with the generic fiber $P_{\mathbf{Q}_2}^2$.

By construction, the action of $GL(3, \mathbf{Q}_2)$ on $P_{\mathbf{Q}_2}^2$ is extended to \mathcal{X} . Mumford found the following discrete subgroup Γ of $GL(3, \mathbf{Q}_2)$. Γ modulo scalar matrices acts on the closed fiber \mathcal{X}_0 freely and induces a quotient \mathcal{X}/Γ as a formal scheme. \mathcal{X}/Γ is algebraized to a projective regular scheme over \mathbf{Z}_2 , and its generic fiber is the fake projective plane.

Γ is contained in the group Γ_1 generated by

$$\sigma = \begin{bmatrix} 1 & 0 & \lambda \\ 0 & 0 & -1 \\ 0 & 1 & -1 \end{bmatrix}, \quad \tau = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 1 + \lambda \\ 0 & 1 & \lambda \end{bmatrix},$$

$$\rho = \begin{bmatrix} 1 & 0 & \lambda \\ 0 & 1 & -\lambda^3/2 \\ 0 & 0 & \lambda^2/2 \end{bmatrix} \quad \text{and} \quad -I_3 = \begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix},$$

where $\lambda = \zeta + \zeta^2 + \zeta^4 = (-1 + \sqrt{-7})/2$ for $\zeta = \exp(2\pi i/7)$. λ is embedded in \mathbf{Z}_2 so that $\lambda = (\text{unit}) \cdot 2$, while its complex conjugate $\bar{\lambda}$ is a unit. There exists a homomorphism $\pi: \Gamma_1 \rightarrow GL(2, \mathbf{F}_7)$ and Γ is given as the inverse image $\pi^{-1}(S)$ of an arbitrary 2-Sylow subgroup S of $GL(2, \mathbf{F}_7)$.

By the matrices in [Mum, p. 243] which describe π , we see that the subgroup of Γ_1 generated by $\{\sigma, \tau, \rho\}$ is mapped onto $SL(2, \mathbf{F}_7)$ by π . Since $-I_3$ is a scalar matrix, the following change of notation does not affect the construction:

MODIFICATION OF THE NOTATION. Γ_1 is replaced by its subgroup of index 2 generated by $\{\sigma, \tau, \rho\}$. The homomorphism π is replaced by one from the new Γ_1 to $PSL(2, \mathbf{F}_7)$. More explicitly, $\pi: \Gamma_1 \rightarrow PSL(2, \mathbf{F}_7)$ is given by

$$\pi(\sigma) = \begin{bmatrix} 2 & 0 \\ 1 & 4 \end{bmatrix}, \quad \pi(\tau) = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \quad \text{and} \quad \pi(\rho) = \begin{bmatrix} 1 & 4 \\ 0 & 1 \end{bmatrix}$$

(see [Mum, p. 243]). The group Γ is also replaced by $\pi^{-1}(S)$ for a 2-Sylow subgroup S of $PSL(2, \mathbf{F}_7)$. In this case, the set of scalar matrices in Γ_1 is $\{(\lambda^2/2)^k I_3 = (\tau\rho)^{3k}; k \in \mathbf{Z}\}$ (cf. [Mum, p. 241]).

From now on, we use this modified notation.

Let $\Gamma_0 = \text{Ker } \pi$. Clearly, Γ_0 is a normal subgroup of Γ_1 . The quotient $G = \Gamma_1/\Gamma_0$ is isomorphic to $PSL(2, \mathbf{F}_7)$ and hence is a simple group of order 168. Since Γ_0 modulo scalar matrices is also a torsionfree cocompact subgroup of $PGL(3, \mathbf{Q}_2)$, the quotient formal scheme \mathcal{X}/Γ_0 can also be algebraized to a projective regular \mathbf{Z}_2 -scheme. We denote the algebraization by V . Then the action of Γ_1 on the scheme \mathcal{X} induces an action of G on V . Since the scalar matrices in Γ_1 are contained in Γ_0 , the induced action is effective. Mumford's fake projective plane is the generic fiber of the quotient $M = V/S$ by the 2-Sylow subgroup S of G .

Since V_η is an unramified cover of degree 8 of Mumford's fake projective plane, the following facts are easily checked.

- (1) V_η is a surface of general type.
- (2) $c_1^2(V_\eta) = 72$.
- (3) $c_2(V_\eta) = 24$.
- (4) $\chi(V_0) = \chi(V_\eta) = 8$.
- (5) $q(V_\eta) = 0$ ([Mum, p. 238]).
- (6) $p_g(V_\eta) = 7$.

In order to describe the closed fiber of M explicitly, we choose the 2-Sylow subgroup S of $G = \Gamma_1/\Gamma_0$ to be the subgroup generated by

$$\begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} 0 & 1 \\ 6 & 0 \end{bmatrix},$$

where we identify G with $PSL(2, F_7)$ by the isomorphism induced by π . S is isomorphic to the dihedral group of order 8. Indeed,

$$\begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix}^4 = I_2, \quad \begin{bmatrix} 0 & 1 \\ 6 & 0 \end{bmatrix}^2 = I_2 \quad \text{and} \quad \begin{bmatrix} 0 & 1 \\ 6 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 6 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix}^{-1}$$

in $PSL(2, F_7)$.

We denote by B the proper transform in \mathcal{X} of the closed fiber $P_{F_2}^2 \subset P_{Z_2}^2 = \text{Proj } Z_2[X_0, X_1, X_2]$. B is an irreducible component of \mathcal{X}_0 and the projection $p: B \rightarrow P_{F_2}^2$ is the blowing-up $P_{F_2}^2$ at the seven F_2 -rational points. We denote by $C(a, b, c)$ the proper transform of the line $aX_0 + bX_1 + cX_2 = 0$ on $P_{F_2}^2$ to B and let $E(a, b, c) := p^{-1}((a, b, c))$ for each triple (a, b, c) of 0 or 1 with not all being zero.

The natural morphism from B to the closed fiber $M_0 = \mathcal{X}_0/\Gamma$ can be regarded as the normalization. Actually, we obtain M_0 by identifying each of suitable seven pairs of $C(a, b, c)$ and $E(a', b', c')$ in B . More precisely, we take $\{\rho\sigma^2\tau, \tau\rho\sigma\tau, \tau^2\rho\tau, \tau^3\rho\sigma\tau^6, \tau^4\rho\sigma^2\tau^5, \tau^5\rho\sigma^2, \tau^6\rho\sigma^2\tau^6\} \subset \Gamma$ as the set of representatives of $S \setminus \{1\}$. Then each element induces an isomorphism of curves on B as follows:

$$\begin{aligned} (\rho\sigma^2\tau)^\wedge &: E(0, 0, 1) \simeq C(1, 1, 0). \\ (\tau\rho\sigma\tau)^\wedge &: E(1, 0, 0) \simeq C(1, 0, 0). \\ (\tau^2\rho\tau)^\wedge &: E(1, 1, 0) \simeq C(0, 1, 0). \\ (\tau^3\rho\sigma\tau^6)^\wedge &: E(1, 1, 1) \simeq C(0, 0, 1). \\ (\tau^4\rho\sigma^2\tau^5)^\wedge &: E(0, 1, 1) \simeq C(1, 0, 1). \\ (\tau^5\rho\sigma^2)^\wedge &: E(1, 0, 1) \simeq C(0, 1, 1). \\ (\tau^6\rho\sigma^2\tau^6)^\wedge &: E(0, 1, 0) \simeq C(1, 1, 1). \end{aligned}$$

In Figure 1, we explicitly describe how these seven pairs are identified. The three points to which the same symbol among A, B, \dots, G is attached are identified to a triple point of M_0 . Here, by $\rho\sigma^2\tau$, the two rational curves $E(0, 0, 1)$ and $C(1, 1, 0)$ are identified in such a way that symbols A, A^*, B come to A^*, A, B , respectively. Consequently, the double curve obtained by this identification has a self-intersection point. Figure 2 indicates the configuration of the double curves on M_0 .

We can check these results by calculating the corresponding action of Γ_1 on the Bruhat-Tits building which is isomorphic to the dual graph of the irreducible components of \mathcal{X}_0 [Mum, p. 235].




FIGURE 1




FIGURE 2

2. Singularities of the quotient surface. Since V is projective and G is finite, the quotient $Y = V/G$ is also a projective \mathbb{Z}_2 -scheme. Although V is regular, Y has some singularities, since the action has fixed points. In this section, we study the singularities of Y .

Let $\bar{\mathbb{Q}}_2$ be the algebraic closure of the 2-adic number field \mathbb{Q}_2 . The discrete valuation v of \mathbb{Q}_2 with $v(2) = 1$ is uniquely extended to a valuation

$$v: \bar{\mathbb{Q}}_2 \rightarrow \mathbb{Q} \cup \{\infty\}.$$

The non-noetherian valuation ring $\bar{\mathbb{Z}}_2 = \{a \in \bar{\mathbb{Q}}_2; v(a) \geq 0\}$ is equal to the integral closure of \mathbb{Z}_2 in $\bar{\mathbb{Q}}_2$. For the maximal ideal $\mathfrak{m} = \{a \in \bar{\mathbb{Z}}_2; v(a) > 0\}$, the residue field $\bar{\mathbb{Z}}_2/\mathfrak{m}$ is equal to the algebraic closure \bar{F}_2 of the prime field F_2 .

In order to describe the geometric points of V_η and Y_η , it is convenient to use the $\bar{\mathbb{Z}}_2$ -valued points of the \mathbb{Z}_2 -scheme \mathcal{X} .

Let $\mathcal{D} := \mathcal{X}(\bar{\mathbb{Z}}_2)$ be the set of $\bar{\mathbb{Z}}_2$ -valued points of \mathcal{X} . Since $\bar{\mathbb{Q}}_2$ is the quotient field of $\bar{\mathbb{Z}}_2$, we have an injection

$$\mathcal{D} \rightarrow \mathcal{X}(\bar{\mathbb{Q}}_2) = P^2(\bar{\mathbb{Q}}_2),$$

where $P^2(\bar{\mathbb{Q}}_2)$ is the projective plane with the coordinates $(X_0: X_1: X_2)$. Hence we use this coordinate system to represent the points of \mathcal{D} through this injection. As we see later, Mumford's fake projective plane is set-theoretically the quotient of \mathcal{D} by $\Gamma \subset GL(3, \bar{\mathbb{Q}}_2)$.

Let $x: \text{Spec}(\bar{\mathbb{Z}}_2) \rightarrow \mathcal{X}$ be a point of \mathcal{D} . Then by composing it with the inclusion $\text{Spec}(\bar{F}_2) \hookrightarrow \text{Spec}(\bar{\mathbb{Z}}_2)$, we get an \bar{F}_2 -valued point of $\mathcal{X}_0 \subset \mathcal{X}$. We denote it by $2\text{-red}(x)$. Let $y \in \mathcal{X}$ be the support point of $2\text{-red}(x)$. Then we get the associated local homomorphism $\mathcal{O}_{y, \mathcal{X}} \rightarrow \bar{\mathbb{Z}}_2$. By this observation, we see that \mathcal{D} is equal to the sum

$$\bigcup_{y \in \mathcal{X}_0} \{x: \mathcal{O}_{y, \mathcal{X}} \rightarrow \bar{\mathbb{Z}}_2; x \text{ is a local } \mathbb{Z}_2\text{-homomorphism}\}.$$

We would like to know which points of $P^2(\bar{\mathbb{Q}}_2)$ are in \mathcal{D} . Since \mathcal{X}_0 is a normal crossing divisor in \mathcal{X} , the points of \mathcal{X}_0 are classified into the following three types: (1) Smooth points of \mathcal{X}_0 . (2) Points lying only on a double curve of \mathcal{X}_0 . (3) Triple points.

Recall that the dual graph which describes the intersections of the components of \mathcal{X}_0 is known as the Bruhat-Tits building. Each irreducible component E of \mathcal{X}_0 corresponds to a free \mathbb{Z}_2 -module $M \subset \mathbb{Q}_2 X_0 + \mathbb{Q}_2 X_1 + \mathbb{Q}_2 X_2$ of rank three modulo the equivalence relation $M \sim 2^*M$. More explicitly, $\text{Proj } S^*M \simeq P^2_{\mathbb{Z}_2}$ for the symmetric algebra S^*M is dominated by \mathcal{X} , and E is the proper transform of the closed fiber. For the detail, see [Mum, p. 235].

(1) Let B be the irreducible component of \mathcal{X}_0 which corresponds to the module $M_0 = \mathbf{Z}_2X_0 + \mathbf{Z}_2X_1 + \mathbf{Z}_2X_2$. The smooth points of \mathcal{X}_0 which are contained in B are exactly those points of $P_{F_2}^2 = \text{Proj } F_2[X_0, X_1, X_2] \subset \text{Proj } \mathbf{Z}_2[X_0, X_1, X_2]$ which are not on the seven F_2 -rational lines on it. These lines are given by $(X_0 = 0)$, $(X_1 = 0)$, $(X_2 = 0)$, $(X_0 + X_1 = 0)$, $(X_0 + X_2 = 0)$, $(X_1 + X_2 = 0)$ and $(X_0 + X_1 + X_2 = 0)$. Hence, a point $x = (x_0 : x_1 : x_2) \in P^2(\bar{\mathbf{Q}}_2)$ is in \mathcal{D} with $2\text{-red}(x)$ in this smooth part if and only if

$$v(x_0) = v(x_1) = v(x_2) = v(x_0 + x_1) = v(x_0 + x_2) = v(x_1 + x_2) = v(x_0 + x_1 + x_2).$$

(2) Let C be the double curve which corresponds to the pair $\mathbf{Z}_2X_0 + \mathbf{Z}_2X_1 + \mathbf{Z}_2X_2/2 \supset \mathbf{Z}_2X_0 + \mathbf{Z}_2X_1 + \mathbf{Z}_2X_2$. It can be shown easily that $2\text{-red}(x)$ of a point $x \in P^2(\bar{\mathbf{Q}}_2)$ is on C and that it is not a triple point if and only if

$$v(x_2) - 1 < v(x_0) = v(x_1) = v(x_0 + x_1) < v(x_2).$$

(3) The triple point P which corresponds to the triple $\mathbf{Z}_2X_0 + \mathbf{Z}_2X_1/2 + \mathbf{Z}_2X_2/2 \supset \mathbf{Z}_2X_0 + \mathbf{Z}_2X_1 + \mathbf{Z}_2X_2/2 \supset \mathbf{Z}_2X_0 + \mathbf{Z}_2X_1 + \mathbf{Z}_2X_2$ is the point $X_1/X_0 = X_2/X_1 = 2X_0/X_2 = 0$ of $\text{Spec } \mathbf{Z}_2[X_1/X_0, X_2/X_1, 2X_0/X_2]$ (see [Mum, p. 234]). Then $2\text{-red}(x)$ is equal to P if and only if

$$v(x_2) - 1 < v(x_0) < v(x_1) < v(x_2).$$

$PGL(3, \mathbf{Q}_2)$ acts transitively on the sets of the irreducible components, the double curves and triple points of \mathcal{X}_0 , respectively. Hence we have the following description of \mathcal{D} .

PROPOSITION 2.1. *Let $x = (x_0 : x_1 : x_2)$ be a point of $P^2(\bar{\mathbf{Q}}_2)$. Then x is in \mathcal{D} if and only if there exists $\alpha \in GL(3, \mathbf{Q}_2)$ such that $(y_0, y_1, y_2) = (x_0, x_1, x_2)\alpha$ satisfies either*

- (i) $v(y_0) = v(y_1) = v(y_2) = v(y_0 + y_1) = v(y_0 + y_2) = v(y_1 + y_2) = v(y_0 + y_1 + y_2)$,
- (ii) $v(y_2) - 1 < v(y_0) = v(y_1) = v(y_0 + y_1) < v(y_2)$ or
- (iii) $v(y_2) - 1 < v(y_0) < v(y_1) < v(y_2)$.

By the above criterion, it is easy to see that any \mathbf{Q}_2 -rational point of $P^2(\bar{\mathbf{Q}}_2)$ is not in \mathcal{D} . In fact, we have the following stronger result.

PROPOSITION 2.2. *Let K be an arbitrary quadratic extension of \mathbf{Q}_2 . If x_0, x_1, x_2 are elements of K , then the point $(x_0 : x_1 : x_2) \in P^2(\bar{\mathbf{Q}}_2)$ is not contained in \mathcal{D} .*

PROOF. Let α be an element of $GL(3, \mathbf{Q}_2)$ and let $(y_0, y_1, y_2) = (x_0, x_1, x_2)\alpha$. Clearly, y_0, y_1, y_2 are also in K . Let \mathcal{O}_K be the integral closure of \mathbf{Z}_2 in K . Since \mathbf{Z}_2 is Henselian, \mathcal{O}_K is also a discrete valuation

ring. Let $u\mathcal{O}_K$ be the maximal ideal of \mathcal{O}_K . Since the ramification index e and the relative degree f satisfy the relation $ef = [K: \mathbb{Q}_2] = 2$, we have two possibilities: Namely,

- (1) $e = 1$ and $f = 2$, i.e., $v(u) = 1$ and $\mathcal{O}_K/u\mathcal{O}_K = \mathbf{F}_4$, or
- (2) $e = 2$ and $f = 1$, i.e., $v(u) = 1/2$ and $\mathcal{O}_K/u\mathcal{O}_K = \mathbf{F}_2$.

We now show that in both cases none of the three conditions in Proposition 2.1 is satisfied. We may assume $y_0, y_1, y_2 \in \mathcal{O}_K$ and one of them is 1 by dividing them by some y_i , if necessary. Let $\bar{y}_0, \bar{y}_1, \bar{y}_2$ be the images of y_0, y_1, y_2 in $\mathcal{O}_K/u\mathcal{O}_K$, respectively.

Case (1). $v(y_0) = v(y_1) = v(y_2) = 0$ implies $\bar{y}_0, \bar{y}_1, \bar{y}_2 \neq 0$. $v(y_0 + y_1) = v(y_0 + y_2) = v(y_1 + y_2) = 0$ implies that $\bar{y}_0, \bar{y}_1, \bar{y}_2$ are distinct elements of \mathbf{F}_4 . Since the sum of the three distinct non-zero elements of \mathbf{F}_4 is zero, we have $v(y_0 + y_1 + y_2) > 0$. Hence (i) of Proposition 2.1 is impossible. Both (ii) and (iii) are obviously impossible, since $v(y_i)$'s are integers.

Case (2). (i) and (ii) are impossible, since $v(y_0) = v(y_1)$ and $\mathcal{O}_K/u\mathcal{O}_K \simeq \mathbf{F}_2$ imply $v(y_0 + y_1) > v(y_0)$. (iii) is also impossible, since $v(y_i)$'s are half integers. q.e.d.

Although $\bar{\mathbf{Z}}_2$ is neither complete nor noetherian, we have the following:

LEMMA 2.3. *Let (A, \mathfrak{m}_A) be a local \mathbf{Z}_2 -algebra essentially of finite type with $2 \in \mathfrak{m}_A$. Then, for the 2-adic completion $i: A \rightarrow A[[2]]$, the induced map*

$$\begin{aligned} i^*: \{f: A[[2]] \rightarrow \bar{\mathbf{Z}}_2; \text{ local } \mathbf{Z}_2\text{-homomorphism}\} \\ \rightarrow \{f: A \rightarrow \bar{\mathbf{Z}}_2; \text{ local } \mathbf{Z}_2\text{-homomorphism}\} \end{aligned}$$

is bijective.

PROOF. Let $f, g: A[[2]] \rightarrow \bar{\mathbf{Z}}_2$ be two local \mathbf{Z}_2 -homomorphisms. Suppose that their restrictions to A are equal. Then they induce the same homomorphism $A/2^n A \rightarrow \bar{\mathbf{Z}}_2/2^n \bar{\mathbf{Z}}_2$ for every $n > 0$. By taking their projective limits, we have a homomorphism $A[[2]] \rightarrow \bar{\mathbf{Z}}_2[[2]]$. Since the natural homomorphism $\bar{\mathbf{Z}}_2 \rightarrow \bar{\mathbf{Z}}_2[[2]]$ is injective, f and g are equal. Hence i^* is injective. We now show the surjectivity. Let $f: A \rightarrow \bar{\mathbf{Z}}_2$ be a local $\bar{\mathbf{Z}}_2$ -homomorphism. Since A is essentially of finite type, the image $f(A)$ is contained in a finite extension of \mathbb{Q}_2 and hence it is a finite \mathbf{Z}_2 -algebra. Hence it is complete in the 2-adic topology. Hence the homomorphism $f: A \rightarrow f(A) \hookrightarrow \bar{\mathbf{Z}}_2$ can be extended to $A[[2]] \rightarrow f(A)$. q.e.d.

Recall that Γ_0 is a normal subgroup of Γ_1 such that $G = \Gamma_1/\Gamma_0$ is isomorphic to $PSL(2, F_7)$. For an element α of Γ_1 , we denote by α^- the

induced automorphism of the \mathbf{Z}_2 -scheme $V = \mathcal{X}/\Gamma_0$.

PROPOSITION 2.4. *There exists a natural map*

$$\varphi: \mathcal{D} \rightarrow V(\bar{\mathbf{Q}}_2)$$

such that the action of Γ_1 on \mathcal{D} and $V(\bar{\mathbf{Q}}_2)$ are compatible with this map, i.e., for an arbitrary element $\alpha \in \Gamma_1$, the diagram

$$\begin{array}{ccc} \mathcal{D} & \xrightarrow{\alpha^\wedge} & \mathcal{D} \\ \varphi \downarrow & & \varphi \downarrow \\ V(\bar{\mathbf{Q}}_2) & \xrightarrow{\alpha^-} & V(\bar{\mathbf{Q}}_2) \end{array}$$

commutes. Furthermore, the induced map $\bar{\varphi}: \mathcal{D}/\Gamma_0 \rightarrow V(\bar{\mathbf{Q}}_2)$ is bijective.

PROOF. Note that $V(\bar{\mathbf{Q}}_2) = V(\bar{\mathbf{Z}}_2)$, since V is proper over \mathbf{Z}_2 . By Lemma 2.3, we have natural bijections

$$\mathcal{D} \simeq \bigcup_{y \in \mathcal{X}_0} \{x: \mathcal{O}_{y, \mathcal{X}}^h \rightarrow \bar{\mathbf{Z}}_2; \text{ local } \mathbf{Z}_2\text{-homomorphism}\}$$

and

$$V(\bar{\mathbf{Q}}_2) \simeq \bigcup_{\bar{y} \in \mathcal{X}_0/\Gamma_0} \{\bar{x}: \mathcal{O}_{\bar{y}, \mathcal{X}/\Gamma_0}^h \rightarrow \bar{\mathbf{Z}}_2; \text{ local } \mathbf{Z}_2\text{-homomorphism}\},$$

where $\mathcal{O}_{y, \mathcal{X}}^h$ (resp. $\mathcal{O}_{\bar{y}, \mathcal{X}/\Gamma_0}^h$) is the local ring at y (resp. \bar{y}) of \mathcal{X} (resp. \mathcal{X}/Γ_0) as a formal scheme, i.e., the 2-adic completion of the usual algebraic local ring. Let $x: \mathcal{O}_{y, \mathcal{X}} \rightarrow \bar{\mathbf{Z}}_2$ be an element of \mathcal{D} . Then, for the image \bar{y} of y in the free quotient \mathcal{X}_0/Γ_0 , we have a natural isomorphism

$$\mathcal{O}_{\bar{y}, \mathcal{X}/\Gamma_0}^h \xrightarrow{\sim} \mathcal{O}_{y, \mathcal{X}}^h.$$

We define $\varphi(x)$ to be the composite

$$\mathcal{O}_{\bar{y}, \mathcal{X}/\Gamma_0} \rightarrow \mathcal{O}_{\bar{y}, \mathcal{X}/\Gamma_0}^h \xrightarrow{\sim} \mathcal{O}_{y, \mathcal{X}}^h \xrightarrow{x'} \bar{\mathbf{Z}}_2,$$

where x' is the homomorphism which satisfies $i^*(x') = x$ for the embedding $i: \mathcal{O}_{y, \mathcal{X}} \rightarrow \mathcal{O}_{y, \mathcal{X}}^h$. Then it is obvious that φ satisfies the assertion of the proposition since \mathcal{X}/Γ_0 is the quotient of the formal scheme \mathcal{X} with respect to a free action. q.e.d.

Now, we study the ramification of the quotient $V_\eta \rightarrow (V/H)_\eta$ with respect to a subgroup $H \subset G$. We need the following elementary ring-theoretic lemmas.

LEMMA 2.5. *Let B be a \mathbf{Z}_2 -algebra of finite type. Assume that a finite group G acts on B as a \mathbf{Z}_2 -algebra, and that a G -invariant maximal ideal \mathfrak{p} contains 2. Then, for the local ring $A = B_\mathfrak{p}$, the ring A^G of G -*

invariant elements of A is essentially of finite type over \mathbf{Z}_2 , and $A^G[[2]]$ is equal to $A[[2]]^G$.

PROOF. Since B is of finite type and G is a finite group, the subring B^G is also of finite type over \mathbf{Z}_2 and B is finite over B^G . Let $\mathfrak{p}^G = B^G \cap \mathfrak{p}$. Then since \mathfrak{p} is G -invariant, $B \setminus \mathfrak{p}$ is a G -invariant multiplicative set with $(B \setminus \mathfrak{p})^G = B^G \setminus \mathfrak{p}^G$. Since G is finite, A is equal to $(B^G \setminus \mathfrak{p}^G)^{-1}B$ and $A^G = (B^G)_{\mathfrak{p}^G}$. Hence A^G is essentially of finite type and A is finite over A^G . There is an exact sequence

$$0 \rightarrow A^G \rightarrow A \xrightarrow{\delta} A^{\oplus |G|} / \Delta(A)$$

of finite A^G -modules, where $\Delta(A)$ is the diagonal and $\delta(a) = (ga)_{g \in G}$. Since $A^G[[2]]$ is flat over A^G , and since $A \otimes_{A^G} A^G[[2]]$ is equal to $A[[2]]$, we get $A^G[[2]] = A[[2]]^G$ by tensoring this exact sequence with $A^G[[2]]$. q.e.d.

LEMMA 2.6. *Let A be a local \mathbf{Z}_2 -algebra essentially of finite type with $2 \in \mathfrak{m}_A$. Let \mathfrak{p} be a prime ideal of A with $2 \notin \mathfrak{p}$ and A/\mathfrak{p} is finite over \mathbf{Z}_2 . Then, for $A' = A[[2]]$, we have $A'_{\mathfrak{p}A'}[[\mathfrak{p}]] = A_{\mathfrak{p}}[[\mathfrak{p}]]$.*

PROOF. Since A/\mathfrak{p} is finite over \mathbf{Z}_2 , the finite A/\mathfrak{p} -module $\mathfrak{p}^n/\mathfrak{p}^{n+1}$ is also a finite \mathbf{Z}_2 -module for every $n \geq 0$. Hence A/\mathfrak{p}^n is a finite \mathbf{Z}_2 -algebra, and is complete in the 2-adic topology. Namely, we have $A/\mathfrak{p}^n = (A/\mathfrak{p}^n)[[2]] = A'/\mathfrak{p}^n A'$. Since $(A/\mathfrak{p}^n)_{\mathfrak{p}/\mathfrak{p}^n} = A_{\mathfrak{p}}/\mathfrak{p}^n A_{\mathfrak{p}}$ and $(A'/\mathfrak{p}^n A')_{\mathfrak{p}A'/\mathfrak{p}^n A'} = A'_{\mathfrak{p}A'}/\mathfrak{p}^n A'_{\mathfrak{p}A'}$, we have $A_{\mathfrak{p}}/\mathfrak{p}^n A_{\mathfrak{p}} = A'_{\mathfrak{p}A'}/\mathfrak{p}^n A'_{\mathfrak{p}A'}$. The lemma is just the projective limit with respect to n of this equality. q.e.d.

Let H be a subgroup of $G = \Gamma_1/\Gamma_0$, and let Γ_H be the pull-back $\pi^{-1}(H) \subset \Gamma_1$. Let x be a point in \mathcal{S} and let $\bar{x} = \varphi(x) \in V(\bar{\mathbf{Q}}_2)$. We denote by $\bar{\Gamma}_1, \bar{\Gamma}_0$ and $\bar{\Gamma}_H$ the images of Γ_1, Γ_0 and Γ_H in $PGL(3, \bar{\mathbf{Q}}_2)$ as in Mumford [Mum, p. 240]. Since $\bar{\Gamma}_H/\bar{\Gamma}_0 \simeq H$ and since $\bar{\Gamma}_0$ acts freely on \mathcal{S} , the isotropy groups

$$T(x, \bar{\Gamma}_H) = \{\alpha^+ \in \bar{\Gamma}_H; \alpha^+(x) = x\} \quad \text{and} \\ T(\bar{x}, H) = \{\alpha^- \in H; \alpha^-(\bar{x}) = \bar{x}\}$$

are isomorphic.

PROPOSITION 2.7. *The singularity of the quotient of $P_{\bar{\mathbf{Q}}_2}^3$ with respect to $T(x, \Gamma_H)$ at the image of x is formally isomorphic to that of the quotient of V with respect to $T(\bar{x}, H)$ at the image of \bar{x} .*

PROOF. Set $T = T(x, \bar{\Gamma}_H)$ and $\bar{T} = T(\bar{x}, H)$. Let y be the support point of 2-red(x) and let $\bar{y} \in V_0$ be the specialization of the support point of \bar{x} . Then by Lemma 2.5, we have

$$(\mathcal{O}_{y,x})^x[[2]] = (\mathcal{O}_{y,x}^h)^x \simeq (\mathcal{O}_{\bar{y},v}^h)^{\bar{x}} = (\mathcal{O}_{\bar{y},v})^{\bar{x}}[[2]] .$$

Let \mathfrak{p} and \mathfrak{p}' be the kernel of the composite homomorphisms

$$(\mathcal{O}_{y,x})^x \rightarrow \mathcal{O}_{y,x} \xrightarrow{x} \bar{\mathcal{Z}}_2 \quad \text{and} \quad (\mathcal{O}_{\bar{y},v})^{\bar{x}} \rightarrow \mathcal{O}_{\bar{y},v} \xrightarrow{\bar{x}} \bar{\mathcal{Z}}_2 ,$$

respectively. Then $((\mathcal{O}_{y,x})^x)_{\mathfrak{p}}$ and $((\mathcal{O}_{\bar{y},v})^{\bar{x}})_{\mathfrak{p}'}$ are the local rings of the support points of x and \bar{x} , respectively. By Lemma 2.6 and the above equality, we have an isomorphism $((\mathcal{O}_{y,x})^x)_{\mathfrak{p}}[[\mathfrak{p}]] \simeq ((\mathcal{O}_{\bar{y},v})^{\bar{x}})_{\mathfrak{p}'}[[\mathfrak{p}']]$. q.e.d.

Now, we study the case $H = G$ and hence $\Gamma_H = \Gamma_1$. We denote by Y the \mathcal{Z}_2 -scheme V/G . Since $T(x, \bar{\Gamma}_1) \simeq T(\bar{x}, G) \subset G$, each element of $T(x, \bar{\Gamma})$ is of finite order. Mumford [Mum, p. 241] has already shown that every element of $\bar{\Gamma}_1$ of finite order is conjugate to one of $\sigma^i \tau^j$ or $(\rho\tau)^i$ for some $0 \leq i \leq 2$ and $0 \leq j \leq 6$. Since $\{\sigma, \tau\}$ generates a non-commutative group of order 21, they are conjugate to one of

$$1, \sigma, \sigma^2, \tau, \tau^2, \dots, \tau^6, (\tau\rho), (\tau\rho)^2 .$$

Since the fixed points of conjugate elements come to the same points in Y , it is sufficient to determine the fixed points of σ, τ and $\tau\rho$ in \mathcal{Z}_0 or \mathcal{D} in order to find out all the ramification points of $f: V \rightarrow Y$.

Before determining the ramification points of $f: V \rightarrow Y$, we have to reformulate some of Mumford's results in a different way.

REMARK 2.8. Mumford has shown the following in his paper.

(i) For the component B of \mathcal{Z}_0 which corresponds to the module $M_0 = \mathcal{Z}_2 X_0 + \mathcal{Z}_2 X_1 + \mathcal{Z}_2 X_2$, the stabilizer $\{\alpha^\wedge \in \bar{\Gamma}_1; \alpha^\wedge(B) = B\}$ is equal to $\bar{\Gamma}_2$ which is the group of order 21 generated by σ and τ (cf. [Mum, p. 241]).

(ii) $\bar{\Gamma}_2$ acts on the F_2 -rational points on B simply transitively (cf. [Mum, p. 242]).

(iii) In particular, if $\alpha^\wedge \in \bar{\Gamma}_1$ fixes B and one F_2 -rational point on it, then $\alpha^\wedge = 1$.

We first determine the fixed points of σ, τ and $\tau\rho$ in the closed fiber \mathcal{Z}_0 . We can do so by looking at the corresponding action on the Bruhat-Tits building as follows:

Let x_0 be a fixed point of σ on \mathcal{Z}_0 . Then there exists an irreducible component B' of \mathcal{Z}_0 which is stable under σ and which contains x_0 . Actually if x_0 is the triple point corresponding to the triple of distinct \mathcal{Z}_2 -submodules $M'_0 \supset M'_1 \supset M'_2$ of $\mathcal{Q}_2 X_0 + \mathcal{Q}_2 X_1 + \mathcal{Q}_2 X_2$ with $M'_i \cong 2M'_0$, then since $\det \sigma = 1$ we have $\sigma(M'_i) = M'_i$ for every i . If x_0 is not triple and is on a double curve of \mathcal{Z}_0 , then σ fixes the two components of \mathcal{Z}_0 which are

adjacent along the double curve since σ is of order three. If x_0 is not on any double curve, then σ stabilizes the unique component which contains x_0 .

Let γ be an element of $\bar{\Gamma}_1$ with $\gamma^\wedge(B) = B'$. Then $(\gamma\sigma\gamma^{-1})^\wedge$ stabilizes B . Since the subgroups of order three of $\bar{\Gamma}_2$ are mutually conjugate, $\gamma\sigma\gamma^{-1}$ is conjugate to σ or σ^2 . Hence the fixed points of σ in B' and B give the same ramification points on Y_0 . It is easy to see that σ has just two fixed points on B . One of them is on $C(1, 0, 0)$ and the other is on $E(1, 0, 0)$, and they are identified by $(\tau\rho\sigma\tau)^\wedge$ in M_0 . The point on $C(1, 0, 0)$ is mapped to the point defined by $X_1^2 + X_1X_2 + X_2^2 = 0$ on the line $X_0 = 0$ in $P^2_{F_2}$ by the natural isomorphism. We denote by w the corresponding ramification point of Y . Clearly, w is of degree two and splits into two points in $Y(\bar{F}_2)$.

Since τ is of order seven, any fixed point of τ in \mathcal{X}_0 is on a stabilized component. Let M'_0 be the module associated to a component of \mathcal{X}_0 stabilized by τ . We may assume $M_0 \supset M'_0$ and $2M_0 \not\supset M'_0$. Since the group generated by τ acts transitively on $(M_0/2M_0) \setminus \{0\}$, we have $M'_0 = M_0$. Hence the fixed points of τ are in B . Later we explicitly determine the fixed points of τ together with those in \mathcal{D} .

Since $\det \tau\rho = \lambda^2/2$, $\tau\rho$ stabilizes no component of \mathcal{X}_0 . Hence it stabilizes no double curve of \mathcal{X}_0 since it is of order three. It is easy to see that $P \in B$ is the unique triple point fixed by $\tau\rho$.

The fixed points of σ , τ and $\tau\rho$ in $P^2(\bar{Q}_2)$ are calculated easily as follows.

$$(1) \quad \sigma = \begin{bmatrix} 1 & 0 & \lambda \\ 0 & 0 & -1 \\ 0 & 1 & -1 \end{bmatrix}, \quad \det(tI_3 - \sigma) = t^3 - 1,$$

$$\begin{array}{llll} \text{eigenvalues} & 1 & \omega & \omega^2 \\ \text{eigenvectors} & (3, \lambda, \lambda) & (0, 1, \omega) & (0, 1, \omega^2), \end{array}$$

where $\omega = (-1 + \sqrt{-3})/2$.

$$(2) \quad \tau = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 1 + \lambda \\ 0 & 1 & \lambda \end{bmatrix},$$

$$\det(tI_3 - \tau) = t^3 - \lambda t^2 - (\lambda + 1)t - 1 = (t - \zeta)(t - \zeta^2)(t - \zeta^4).$$

$$\begin{array}{llll} \text{eigenvalues} & \zeta & \zeta^2 & \zeta^4 \\ \text{eigenvectors} & (1, \zeta, \zeta^2) & (1, \zeta^2, \zeta^4) & (1, \zeta^4, \zeta), \end{array}$$

where $\zeta = \exp(2\pi i/7)$.

$$(3) \quad \tau\rho = \begin{bmatrix} 0 & 0 & \lambda^2/2 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}, \quad \det(tI_3 - \tau\rho) = t^3 - \lambda^2/2,$$

eigenvalues	ε	$\omega\varepsilon$	$\omega^2\varepsilon$
eigenvectors	$(1, \varepsilon, \varepsilon^2)$	$(1, \omega\varepsilon, \omega^2\varepsilon^2)$	$(1, \omega^2\varepsilon, \omega\varepsilon^2)$,

where $\varepsilon = (\lambda^2/2)^{1/3}$.

In case (1), since every component of the eigenvectors are in $\mathbf{Q}_2(\sqrt{-7})$ or $\mathbf{Q}_2(\sqrt{-3})$, the fixed points of σ in $P^2(\bar{\mathbf{Q}}_2)$ are outside \mathcal{S} by Proposition 2.2.

In case (2), set $\tilde{q} = (1, \zeta, \zeta^2)$. Note that $\sigma^\wedge(\tilde{q}) = (1, \zeta^2, \zeta^4)$ and $(\sigma^\wedge)^2(\tilde{q}) = (1, \zeta^4, \zeta)$. Let ζ_0 be the image of ζ in $\mathbf{Z}_2(\zeta)/(2) \simeq \mathbf{F}_8$. Then since $\zeta_0 \in \mathbf{F}_8 \setminus \mathbf{F}_4$, we see that $1 + \zeta_0, 1 + \zeta_0^2, \zeta_0 + \zeta_0^2$ and $1 + \zeta_0 + \zeta_0^2$ are not zero. This implies that $v(1) = v(\zeta) = v(\zeta^2) = v(1 + \zeta) = v(1 + \zeta^2) = v(\zeta + \zeta^2) = v(1 + \zeta + \zeta^2) = 0$. Hence \tilde{q} is a point of \mathcal{S} by Proposition 2.1. We denote by q the image $f \circ \varphi(\tilde{q}) \in Y(\bar{\mathbf{Q}}_2)$.

Since $2\text{-red}(\tilde{q})$ is a smooth point of \mathcal{X}_0 and is on the component B , the isotropy group $T(\tilde{q}, \bar{\Gamma}_1)$ is a subgroup of $\bar{\Gamma}_2$ by Remark 2.8. Since σ does not fix $\tilde{q} \in \mathcal{S}$, we have $T(\tilde{q}, \bar{\Gamma}_1) = \langle \tau \rangle$. As we see later in Remark 2.10, the linear map τ is given locally at \tilde{q} by $(y_1, y_2) \mapsto (\zeta y_1, \zeta^3 y_2)$. Hence the singularity of the quotient at this point is the cyclic quotient singularity of type (7, 3). By Proposition 2.7, the singularity of $Y(\bar{\mathbf{Q}}_2)$ at q is also a cyclic quotient singularity of type (7, 3).

These τ -invariant points of $P^2(\bar{\mathbf{Q}}_2)$ are $\mathbf{Q}_2(\zeta)$ -valued and they are identified to q in $Y(\bar{\mathbf{Q}}_2)$ by σ . Since the action of σ on these three points is compatible with the automorphism of $\mathbf{Q}_2(\zeta)$ defined by $\zeta \mapsto \zeta^2$, we see that q is a \mathbf{Q}_2 -valued point. Since Y is proper over \mathbf{Z}_2 , there exists a \mathbf{Z}_2 -valued point $\bar{q}: \text{Spec } \mathbf{Z}_2 \rightarrow Y$ such that $\bar{q}(\eta) = q$. We see easily that $\bar{q}(0) \in Y_0$ is also a cyclic quotient singularity of type (7, 3). We can see similarly that the fixed points of τ on B are only $(1, \zeta_0, \zeta_0^2), (1, \zeta_0^2, \zeta_0^4)$ and $(1, \zeta_0^4, \zeta_0)$. Since B is the only component of \mathcal{X}_0 stabilized by τ , we see that $\bar{q} \subset Y$ is the unique ramification locus given by τ .

Finally in case (3), set $\tilde{p}_0 = (1, \varepsilon, \varepsilon^2), \tilde{p}_1 = (1, \omega\varepsilon, \omega^2\varepsilon^2)$ and $\tilde{p}_2 = (1, \omega^2\varepsilon, \omega\varepsilon^2)$. Since $v(\lambda^2/2) = 1$, we have $v(\varepsilon) = 1/3$, while $v(\omega) = 0$. Hence these points are in \mathcal{S} by Proposition 2.1. In this case, $2\text{-red}(\tilde{p}_i)$'s are the same triple point $P \in \mathcal{X}_0$. At this point P , the three components of \mathcal{X}_0 which correspond to $\mathbf{Z}_2 X_0 + \mathbf{Z}_2 X_1/2 + \mathbf{Z}_2 X_2/2, \mathbf{Z}_2 X_0 + \mathbf{Z}_2 X_1 + \mathbf{Z}_2 X_2/2$ and $\mathbf{Z}_2 X_0 + \mathbf{Z}_2 X_1 + \mathbf{Z}_2 X_2$ meet together. In particular, the component B contains P . Suppose $\alpha^\wedge \in \bar{\Gamma}_1$ fixes P . Then since $\tau\rho$ cyclically permutes

the three components, $(\tau\rho)^{-i}\alpha^\wedge$ stabilize B for $i = 0, 1$ or 2 . By (iii) of Remark 2.8, we get $\alpha^\wedge = (\tau\rho)^i$. Since the isotropy group of \tilde{p}_i 's are contained in that of P , we have $T(\tilde{p}_i, \bar{\Gamma}_1) = \langle \tau\rho \rangle$.

No $\alpha \in \bar{\Gamma}_1$ maps \tilde{p}_i to another \tilde{p}_j since $\alpha^\wedge(\tilde{p}_i) = \tilde{p}_j$ implies $\alpha \in \langle \tau\rho \rangle$. Hence, the points $\tilde{p}_0, \tilde{p}_1, \tilde{p}_2$ are mapped to distinct points in $Y(\bar{\mathcal{Q}}_2)$. Let them be p_0, p_1 and p_2 , respectively. As in case (2), we see that $Y(\bar{\mathcal{Q}}_2)$ has cyclic quotient singularities of type $(3, 2)$ at these points.

The points $\tilde{p}_0, \tilde{p}_1, \tilde{p}_2$ are solutions of the system of equations $(X_1/X_0) = (X_2/X_1) = (\varepsilon^3 X_0/X_2)$. Since the local ring of \mathcal{X} at P is $\mathbf{Z}_2[X_1/X_0, X_2/X_1, \varepsilon^3 X_0/X_2]_{\mathfrak{m}}$ for the maximal ideal $\mathfrak{m} = (X_1/X_0, X_2/X_1, \varepsilon^3 X_0/X_2)$, the equations give a $\mathbf{Z}_2[\varepsilon]$ -valued point \bar{p} of Y such that $\bar{p}(0) = P$ and that the image of $\bar{p}(\gamma)$ in Y is a $\mathbf{Q}_2(\varepsilon)$ -valued point which splits into the three points p_0, p_1, p_2 in $Y(\bar{\mathcal{Q}}_2)$. Since P is the unique fixed point of $\tau\rho$ in \mathcal{X} , we see that \bar{p} is the unique ramification locus of $f: V \rightarrow Y$ caused by $\tau\rho$.

Thus we conclude:

THEOREM 2.9. *The morphism $f: V \rightarrow Y$ is ramified along \bar{q}, \bar{p} and at the point $w \in Y_0$ of degree two. The restriction to the geometric fibers $f_{\bar{q}_2}: V(\bar{\mathcal{Q}}_2) \rightarrow Y(\bar{\mathcal{Q}}_2)$ is ramified at the point p_0, p_1, p_2 and q . p_0, p_1 and p_2 (resp. q) are cyclic quotient singularities of type $(3, 2)$ (resp. of type $(7, 3)$).*

REMARK 2.10. Let R be the étale finite ring extension $\mathbf{Z}_2[\zeta, \omega]$ of \mathbf{Z}_2 . We can describe the minimal resolution of the singularities along \bar{q} and \bar{p} after the étale base extension $Y_R \rightarrow \text{Spec } R$ of $Y \rightarrow \text{Spec } \mathbf{Z}_2$ as follows:

By the coordinate change

$$(Y_0, Y_1, Y_2) := (X_0, X_1, X_2) \begin{bmatrix} 1 & \zeta & \zeta^2 \\ 1 & \zeta^2 & \zeta^4 \\ 1 & \zeta^4 & \zeta \end{bmatrix}^{-1},$$

of P_R^2 , τ is diagonalized as

$$\begin{bmatrix} \zeta & 0 & 0 \\ 0 & \zeta^2 & 0 \\ 0 & 0 & \zeta^4 \end{bmatrix}$$

and the eigenvectors are $(1, 0, 0)$, $(0, 1, 0)$ and $(0, 0, 1)$. Hence the local ring of Y_R at $\bar{q}(0)$ is formally isomorphic to the localization of the ring of invariants $R[Y_1/Y_0, Y_2/Y_0]^{\tau}$ in the polynomial ring $R[Y_1/Y_0, Y_2/Y_0]$ with respect to the action of τ defined by $Y_1/Y_0 \mapsto \zeta Y_1/Y_0$ and $Y_2/Y_0 \mapsto \zeta^3 Y_2/Y_0$. One can resolve it minimally by the standard method. For any geometric

fiber, the exceptional set is a chain of nonsingular rational curves with the self-intersection numbers $-3, -2, -2$.

The local ring of Y_R at $\bar{p}(0)$ is formally isomorphic to the localization of the ring of invariants $R[X_1/X_0, X_2/X_1, \varepsilon^3 X_0/X_2]^{\tau\rho} \subset R[X_1/X_0, X_2/X_1, \varepsilon^3 X_0/X_2]$ with respect to the automorphism $\tau\rho$ given by $X_1/X_0 \mapsto X_2/X_1, X_2/X_1 \mapsto \varepsilon^3 X_0/X_2, \varepsilon^3 X_0/X_2 \mapsto X_1/X_0$. Note that $\varepsilon^3 = \lambda^2/2$ is a generator of the maximal ideal of the discrete valuation ring R . By the coordinate change

$$(T_0, T_1, T_2) = (X_1/X_0, X_2/X_1, \varepsilon^3 X_0/X_2) \begin{bmatrix} 1 & 1 & 1 \\ 1 & \omega^2 & \omega \\ 1 & \omega & \omega^2 \end{bmatrix},$$

we have

$$\tau\rho = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \omega & 0 \\ 0 & 0 & \omega^2 \end{bmatrix}.$$

Then the ring of invariants is $R[T_0, T_1^3, T_2^3, T_1 T_2]$ with the relation $T_0^3 + T_1^3 + T_2^3 - 3T_0 T_1 T_2 = 27\varepsilon^3$. We see easily that this is a complete intersection of a regular ring. In particular, this is a Gorenstein ring. This singularity is resolved by the blowing up along the prime ideal $(T_1^3, T_2^3, T_1 T_2)$. For the geometric fiber $Y(\bar{Q}_2)$, this is the blowing-up at $\{p_0, p_1, p_2\}$. Since these are cyclic quotient singularities of type $(3, 2)$, this blowing-up gives the minimal resolution of these singular points and each exceptional set is the union of two nonsingular rational curves with the self-intersection numbers -2 intersecting each other at one point.

Thus we minimally resolved the singularities of Y_R along \bar{q} and \bar{p} . Since this resolution is canonical, it descends to a scheme Y' over Z_2 . Clearly, $Y'(\bar{Q}_2)$ is the minimal resolution of $Y(\bar{Q}_2)$.

3. The plurigenera of the quotient surface. In this section, we study pluri-canonical line bundles on V and its quotients.

The component B of \mathcal{Z}_0 is a smooth rational surface, and the fourteen rational curves $C(a, b, c)$'s and $E(a, b, c)$'s form a divisor $A = \cup_{a,b,c} (C(a, b, c) \cup E(a, b, c))$ with only normal crossings in B . For the unramified covering $\mathcal{Z}_0 \rightarrow V_0$, we denote by $B_1, C(a, b, c)_1, E(a, b, c)_1, P_1$ and A_1 the image of $B, C(a, b, c), E(a, b, c), P$ and A in V_0 , respectively. Note that the fixed point $P \in \mathcal{Z}_0$ of $\tau\rho$, is the intersection point of $C(0, 0, 1)$ and $E(1, 0, 0)$. One can check that B_1 has no self-intersection. Hence B_1 is isomorphic to B .

From now on, we mainly treat V and its quotient with respect to a

subgroup of G . Hence, for simplicity, we denote also by σ, τ, ρ their images in G . For an element $\alpha \in G$, we denote by α^- the associated automorphism of V as in Section 2.

Since $M_0 = V_0/S$ consists of only one irreducible component, we have

$$V_0 = \bigcup_{\alpha \in S} B_\alpha \quad \text{where} \quad B_\alpha = \alpha^-(B_1).$$

Here B_α 's cross each other normally and the normalization \tilde{V}_0 is equal to the disjoint union $\coprod_{\alpha \in S} B_\alpha$. Let $\varphi: \tilde{V}_0 \rightarrow V_0$ be the natural morphism.

Since the induced action of G on the set of double curves is transitive, and since the stabilizer of the double curve $D_1 = C(1, 0, 0)_1$ is $\{1, \sigma, \sigma^2\}$, we see that the union D of the double curves is

$$D = \bigcup_{\beta \in G/\langle \sigma \rangle} D_\beta,$$

where $G/\langle \sigma \rangle$ is the set of left cosets $\{\langle \sigma \rangle g; g \in G\}$ and $D_\beta := \beta^-(D_1)$.

Similarly, the stabilizer of P_1 is $\{1, \tau\rho, (\tau\rho)^2\}$ and

$$\{P_\mu = \mu^-(P_1); \mu \in G/\langle \tau\rho \rangle\}$$

is the set of the triple points of V_0 . Note that the set of F_2 -rational points of V_0 is exactly equal to this set.

For the union D of the double curves of V_0 , let $\delta: \tilde{D} \rightarrow V_0$ be the natural morphism from the normalization $\tilde{D} = \coprod_{\beta \in G/\langle \sigma \rangle} D_\beta$ of D to V_0 .

Since the double curves arise from the identification of (-1) -curves and (-2) -curves [Mum, p. 236], there exist morphisms $\varepsilon, \gamma: \tilde{D} \rightarrow \tilde{V}_0$ such that $\varepsilon(D_\beta)^2 = -1$ and $\gamma(D_\beta)^2 = -2$ for every component D_β of \tilde{D} and $\varphi \circ \varepsilon = \varphi \circ \gamma = \delta$. The union $\varepsilon(\tilde{D}) \cup \gamma(\tilde{D})$ is equal to $\coprod_{\alpha \in S} A_\alpha$, where $A_\alpha = \alpha^-(A_1) \subset B_\alpha$.

For any line bundle L on V_0 , the following diagram is exact:

$$H^0(V_0, L) \xrightarrow{\varphi^*} H^0(\tilde{V}_0, \varphi^*L) \xrightarrow[\gamma^*]{\varepsilon^*} H^0(\tilde{D}, \delta^*L).$$

For an equidimensional Gorenstein scheme Z , we denote by ω_Z its canonical invertible sheaf. As is well known for varieties with normal crossing singularities, we have

$$\varphi^* \omega_{V_0} = \omega_{\tilde{V}_0}(\varepsilon(\tilde{D}) \cup \gamma(\tilde{D})) = \bigoplus_{\alpha \in S} \omega_{B_\alpha}(A_\alpha).$$

Hence we get the exact diagram

$$(1) \quad H^0(V_0, \omega_{V_0}^{\otimes m}) \rightarrow \bigoplus_{\alpha \in S} H^0(B_\alpha, \omega_{B_\alpha}^{\otimes m}(mA_\alpha)) \xrightarrow[\gamma^*]{\varepsilon^*} H^0(\tilde{D}, \delta^* \omega_{V_0}^{\otimes m})$$

for every integer m .

On the other hand, $\bigoplus_{\alpha \in S} \varphi_* \mathcal{O}_{B_\alpha}(-A_\alpha)$ is equal to the ideal $I_D \subset \mathcal{O}_{V_0}$

defining D . Hence by the projection formula, we have

$$\omega_{V_0}^{\otimes m} \otimes I_D = \bigoplus_{\alpha \in S} \varphi_* \omega_{B_\alpha}^{\otimes m}((m-1)A_\alpha).$$

Hence we get an exact sequence of \mathcal{O}_{V_0} -modules

$$(2) \quad 0 \rightarrow \bigoplus_{\alpha \in S} \varphi_* \omega_{B_\alpha}^{\otimes m}((m-1)A_\alpha) \rightarrow \omega_{V_0}^{\otimes m} \rightarrow \omega_{V_0}^{\otimes m} \otimes \mathcal{O}_D \rightarrow 0.$$

Now we analyze the sections of $\omega_B^{\otimes m}(mA)$ and $\omega_B^{\otimes m}((m-1)A)$ more precisely.

For the projective plane $P_{F_2}^2$ with the homogeneous coordinate system $(X_0: X_1: X_2)$, we set $y = X_0/X_2$ and $z = X_1/X_2$. Then the rational 2-form $\omega_0 = (dy \wedge dz)/yz$ vanishes nowhere and has a pole of order one along the divisor $(X_0 X_1 X_2 = 0)$. Let $p^* \omega_0$ be the pull-back of ω_0 with respect to the natural morphism $p: B \rightarrow P_{F_2}^2$. Then, the divisor $(p^* \omega_0)$ is equal to

$$E(1, 1, 1) - C(1, 0, 0) - C(0, 1, 0) - C(0, 0, 1) \\ - E(1, 0, 0) - E(0, 1, 0) - E(0, 0, 1).$$

Hence $p^* \omega_0$ is a section of $\omega_B(A)$ with the zero divisor

$$F_0 = C(1, 1, 0) + C(1, 0, 1) + C(0, 1, 1) + C(1, 1, 1) \\ + E(1, 1, 0) + E(1, 0, 1) + E(0, 1, 1) + 2E(1, 1, 1).$$

Let F be a divisor on B which is linearly equivalent to F_0 . Then the images $p(F_0)$ and $p(F)$ in $P_{F_2}^2$ are also linearly equivalent. Since $p(F_0) = (u_0 = 0)$ for $u_0 = (X_0 + X_1)(X_0 + X_2)(X_1 + X_2)(X_0 + X_1 + X_2)$, we see that $p(F)$ is equal to $(f = 0)$ for a homogeneous quartic polynomial $f \in F_2[X_0, X_1, X_2]$.

Since $p^*(u_0 = 0) - F_0 = \sum_{a,b,c} E(a, b, c)$ should be equal to $p^*(f = 0) - F$, the divisor $(f = 0) \subset P_{F_2}^2$ contains all the seven F_2 -rational points of $P_{F_2}^2$. Conversely, if f is a quartic homogeneous polynomial with $f(a, b, c) = 0$ for all triple (a, b, c) of 0 or 1, then $p^*(f = 0) - \sum_{a,b,c} E(a, b, c)$ is effective and linearly equivalent to F_0 . Hence $(f/u_0)p^* \omega_0$ is a section of $\omega_B(A)$.

Thus the space of section of $\omega_B(A)$ is described as

$$(3) \quad H^0(\omega_B(A)) = \left\{ \frac{f}{u_0} \left(\frac{dy}{y} \wedge \frac{dz}{z} \right) \right\},$$

where f runs over the homogeneous polynomials in $F_2[X_0, X_1, X_2]$ of degree 4 such that $f(a, b, c) = 0$ if $a, b, c = 0$ or 1.

Similarly for general $m \in \mathbb{Z}$, we get the following:

$$(4) \quad H^0(\omega_B^{\otimes m}(mA)) = \left\{ \frac{f}{u_0^m} \left(\frac{dy}{y} \wedge \frac{dz}{z} \right)^{\otimes m} \right\},$$

where f runs over the homogeneous polynomials in $F_2[X_0, X_1, X_2]$ of degree $4m$ which has zero of multiplicity at least m at each of the seven F_2 -rational points of $P_{F_2}^2$.

Let $\omega = (f/u_0^m)(dy/y \wedge dz/z)^{\otimes m}$ be an element of $H^0(\omega_B^{\otimes m}(mA))$. Then ω is in $H^0(\omega_B^{\otimes m}((m-1)A))$ if and only if f has the factor $u = X_0X_1X_2(X_0 + X_1)(X_0 + X_2)(X_1 + X_2)(X_0 + X_1 + X_2)$ and f has zero of multiplicity at least $m + 1$ at every F_2 -rational point of $P_{F_2}^2$. Since u has zeros of multiplicity three at these points, we see that

$$(5) \quad H^0(\omega_B^{\otimes m}((m-1)A)) = \left\{ \frac{ug}{u_0^m} \left(\frac{dy}{y} \wedge \frac{dz}{z} \right)^{\otimes m} \right\},$$

where g runs over the homogeneous polynomials in $F_2[X_0, X_1, X_2]$ of degree $4m - 7$ which has zeros of multiplicity at least $m - 2$ at the seven F_2 -rational points of $P_{F_2}^2$.

Let m be an integer greater than one. Since $c_1^2(V_\eta) = 72$ and $\chi(\mathcal{O}_{V_\eta}) = 8$, we have $P_m(V_\eta) = \chi^0(\omega_{V_\eta}^{\otimes m}) = 36m(m-1) + 8$ by the plurigenus formula for surfaces of general type. Hence $H^0(V, \omega_{V_0}^{\otimes m})$ is a free Z_2 -module of rank $36m(m-1) + 8$. By Grothendieck's base change theorem, we have a natural injection

$$i_m: H^0(V, \omega_{V_0}^{\otimes m}) \otimes_{Z_2} F_2 \hookrightarrow H^0(V_0, \omega_{V_0}^{\otimes m}).$$

More generally, let H be a subgroup of G acting freely on V and let $V' = V/H$. Then we have an injection

$$i'_m: H^0(V', \omega_{V'_0}^{\otimes m}) \otimes_{Z_2} F_2 \hookrightarrow H^0(V'_0, \omega_{V'_0}^{\otimes m}).$$

Note that the left hand side is of dimension $(36m(m-1) + 8)/|H|$, since V'_η is also of general type.

PROPOSITION 3.1. *The above homomorphisms i_m and i'_m are isomorphisms for $m = 2$ and 3.*

PROOF. We give the proof only for i_m , since the proof for general i'_m is similar. Suppose $m = 2$. By (5), we have

$$H^0(\omega_B^{\otimes 2}(A)) = \left\{ \frac{(aX_0 + bX_1 + cX_2)u}{u_0^2} \left(\frac{dy}{y} \wedge \frac{dz}{z} \right)^{\otimes 2}; a, b, c \in F_2 \right\}.$$

This is obviously three-dimensional. Hence $\bigoplus_{\alpha \in S} H^0(\omega_{B_\alpha}^{\otimes 2}(A_\alpha))$ is of dimension $8 \times 3 = 24$. On the other hand, V_0 has fifty-six F_2 -rational points $\{P_\mu\}_{\mu \in G/\langle \tau \rho \rangle}$. Hence there exists a natural homomorphism

$$(6) \quad j_2: H^0(V_0, \omega_{V_0}^{\otimes 2}) \rightarrow \bigoplus_{\mu \in G/\langle \tau \rho \rangle} \omega_{V_2}^{\otimes 2}(P_\mu).$$

Here the right hand side is an F_2 -vector space of dimension 56. Hence

it suffices to show that the kernel $\text{Ker } j_2$ is contained in $\bigoplus_{\alpha \in S} H^0(\omega_{B_\alpha}^{\otimes 2}(A_\alpha))$, because then the dimension of $H^0(V_0, \omega_{V_0}^{\otimes 2})$ is at most $24 + 56 = 80$ which is the rank of $H^0(V, \omega^{\otimes 2})$.

Let ω be an element of $\text{Ker } j_2$. We have to show that $\omega|_{D_\beta} = 0$ on each double curve D_β . Set $M_\beta = \delta^* \omega_{V_0}|_{D_\beta}$. Since $\delta^* \omega_{V_0}|_{D_\beta} = \gamma^* \omega_{B_\alpha}(A_\alpha)|_{D_\beta}$ for some $\alpha \in S$, and since $\gamma(D_\beta)$ is a nonsingular rational curve with $\gamma(D_\beta)^2 = -2$, we have

$$\text{deg } M_\beta = \text{deg } \omega_{B_\alpha}|_{\gamma(D_\beta)} + \gamma(D_\beta) \cdot A_\alpha = 0 + 1 = 1.$$

Since $D_\beta \simeq P^1(F_2)$ has three F_2 -rational points and ω is zero there, $\omega|_{D_\beta} \in H^0(M_\beta^{\otimes 2})$ should be zero.

We now consider the case $m = 3$. By (5), $H^0(\omega_B^{\otimes 3}(2A))$ is isomorphic to the module of homogeneous quintic polynomials which have zeros at all the seven F_2 -rational points of $P_{F_2}^2$. It is easy to see that this is of dimension $21 - 7 = 14$. Hence $\bigoplus_{\alpha \in S} H^0(\omega_{B_\alpha}^{\otimes 3}(2A_\alpha))$ is of dimension $8 \times 14 = 112$. Let L be the kernel of the homomorphism

$$(7) \quad j_3: H^0(V_0, \omega_{V_0}^{\otimes 3}) \rightarrow \bigoplus_{\mu \in G/\langle \tau \rho \rangle} \omega_{V_0}^{\otimes 3}(P_\mu) \simeq F_2^{\oplus 56}.$$

Clearly, L is of codimension at most 56 in $H^0(V_0, \omega_{V_0}^{\otimes 3})$. Let D_β be a double curve of V_0 , and let $0, 1, \infty$ be its F_2 -rational points. We consider the restriction map $L \rightarrow H^0(M_\beta^{\otimes 3})$. Since $\text{deg } M_\beta^{\otimes 3} = 3$ and since each element $\omega \in L$ has zeros at $\{0, 1, \infty\}$, the image of this map is in $H^0(M_\beta^{\otimes 3}(-0 - 1 - \infty)) \simeq F_2$. Hence the kernel of the natural homomorphism

$$(8) \quad L \rightarrow \bigoplus_{\beta \in G/\langle \sigma \rangle} H^0(M_\beta^{\otimes 3}(-0 - 1 - \infty)) = F_2^{\oplus 56}$$

is of codimension at most 56. Since the kernel is contained in $\bigoplus_{\alpha \in S} H^0(\omega_{B_\alpha}^{\otimes 3}(2A_\alpha))$, we see that the dimension of $H^0(V_0, \omega_{V_0}^{\otimes 3})$ is at most $112 + 56 + 56 = 224$ which is the rank of $H^0(V, \omega^{\otimes 3})$. Hence i_3 is an isomorphism. q.e.d.

REMARK 3.2. This proof implies that the homomorphisms (6), (7) and (8) are surjective. This is also true for the homomorphism i'_m .

PROPOSITION 3.3. *Let H be a subgroup of G , and let ω be an element of $H^0(V_0, \omega_{V_0}^{\otimes m})$ for $m = 2$ or 3 . If ω is H -invariant, then there exists an element $\tilde{\omega} \in H^0(V, \omega^{\otimes m})$ which is H -invariant and $\tilde{\omega}|_{V_0} = \omega$.*

PROOF. Let S_0 be a 2-Sylow subgroup of H . Then since S_0 is contained in a 2-Sylow subgroup of G , S_0 acts on V freely by a result of Mumford. Let V' be the quotient V/S_0 . Since ω is S_0 -invariant, it descends to an element of $H^0(V'_0, \omega_{V'_0}^{\otimes m})$. By Proposition 3.1, there exists

$\tilde{\omega}' \in H^0(V', \omega_{V'})$ with $\tilde{\omega}'|_{V_0} = \omega$. We regard $\tilde{\omega}'$ as an S_0 -invariant element of $H^0(V, \omega_{V'}^{\otimes 2})$. Let $H = S_0\alpha_1 + \dots + S_0\alpha_n$ be the left coset decomposition of H with respect to S_0 . Let $\tilde{\omega} = \sum_{i=1}^n \alpha_i^*(\tilde{\omega}')$. Then $\tilde{\omega}$ is H -invariant, and $\tilde{\omega}|_{V_0} = n\omega = \omega$, since $n = [H:S_0]$ is an odd number. q.e.d.

THEOREM 3.4. *Let H be a subgroup of G and let m be 2 or 3. Then the homomorphism*

$$H^0(V, \omega_{V'}^{\otimes m})^H \otimes_{\mathbb{Z}_2} \mathbb{F}_2 \rightarrow H^0(V_0, \omega_{V_0}^{\otimes m})^H$$


induced by i_m is an isomorphism.

PROOF. Since the quotient $H^0(V, \omega_{V'}^{\otimes m})/H^0(V, \omega_{V'}^{\otimes m})^H$ is contained in the \mathbb{Q}_2 -module $H^0(V_\eta, \omega_{V_\eta}^{\otimes m})/H^0(V_\eta, \omega_{V_\eta}^{\otimes m})^H$, it is a free \mathbb{Z}_2 -module. Hence $H^0(V, \omega_{V'}^{\otimes m})^H$ is a direct summand of $H^0(V, \omega_{V'}^{\otimes m})$. In particular, the homomorphism is injective. Since $m = 2$ or 3 , it is surjective by Proposition 3.3. q.e.d.

The following shows that the bigenus P_2 of the desingularization of the quotient surface V_η/H is calculated only in terms of the closed fiber V_0 .

PROPOSITION 3.5. *Let H be a subgroup of G , and let \tilde{Z} be the minimal resolution of $Z = V_\eta/H$. Then $P_2(\tilde{Z}) = \dim H^0(V_0, \omega_{V_0}^{\otimes 2})^H$.*

PROOF. By Theorem 3.4, we have $\dim H^0(V_\eta, \omega_{V_\eta}^{\otimes 2})^H = \dim H^0(V_0, \omega_{V_0}^{\otimes 2})^H$. By Theorem 2.9, Z may only have at most cyclic quotient singularities of types (3, 2) or (7, 3), and the morphism $V_\eta \rightarrow Z$ is ramified only at these singular points. Hence an element $s \in H^0(V_\eta, \omega_{V_\eta}^{\otimes 2})^H$ can be regarded as a section of $\omega_{Z'}^{\otimes 2}$, where $Z' = Z \setminus \{\text{singular points}\}$. Note that \tilde{Z} contains Z' as an open subset. It suffices to show that the rational section s of $\omega_{Z'}^{\otimes 2}$ has no pole along the exceptional divisors. This is the case over the cyclic quotient singularities of type (3, 2), since they are rational double points. Let $y \in Z$ be a cyclic quotient singularity of type (7, 3) and let D_1, D_2, D_3 be the exceptional curves for the resolution of y with $D_1^2 = -3, D_2^2 = D_3^2 = -2, D_1 \cdot D_2 = D_2 \cdot D_3 = 1$ and $D_1 \cdot D_3 = 0$.



We can write the divisor (s) on \tilde{Z} as $aD_1 + bD_2 + cD_3 + F$, where the support of F contains none of D_i 's. Let d_i be the intersection number $D_i \cdot F$ for $i = 1, 2, 3$. Since (s) is linearly equivalent to $2K_{\tilde{Z}}$, we have

$$\begin{aligned} 2 &= (s) \cdot D_1 = -3a + b + d_1, \\ 0 &= (s) \cdot D_2 = a - 2b + c + d_2, \\ 0 &= (s) \cdot D_3 = b - 2c + d_3. \end{aligned}$$

By these equalities, we calculate easily that

$$\begin{aligned} 7a &= 3(d_1 - 2) + 2d_2 + d_3, \\ 7b &= 2(d_1 - 2) + 6d_2 + 3d_3, \\ 7c &= (d_1 - 2) + 3d_2 + 5d_3. \end{aligned}$$

Since a, b, c are integers and d_1, d_2, d_3 are nonnegative, we have $a, b, c \geq 0$. Hence s has no pole on \tilde{Z} . q.e.d.

Recall that $\Gamma_2 = \langle \sigma, \tau \rangle$ stabilizes the component B of \mathcal{L}_0 . We denote by G_{21} the injective image of Γ_2 in G . G_{21} is a group of order 21. Since $G_{21} \cap S = \{1\}$, G is equal to the disjoint union $\cup_{\alpha \in S} G_{21}\alpha$. If an element β is in $G_{21}\alpha$, then β induces an isomorphism $(\beta|_{B_1}): B_1 \rightarrow B_\alpha$.

The action of G on V_0 induces an action on the diagram (1). An element $(\omega_\alpha)_{\alpha \in S} \in \bigoplus_{\alpha \in S} H^0(B_\alpha, \omega_{B_\alpha}^{\otimes m}(mA_\alpha))$ is G -invariant if and only if $(\beta|_{B_1})^* \omega_\alpha = \omega_1$ for every $\beta \in G$, where α is the element of S with $\beta \in G_{21}\alpha$. This is also equivalent to the condition that ω_1 is G_{21} -invariant and $\omega_1 = (\alpha|_{B_1})^* \omega_\alpha$ for every $\alpha \in S$.

Suppose (ω_α) is G -invariant. By the diagram (1), (ω_α) is in $H^0(V_0, \omega_{V_0}^{\otimes m})^G$ if and only if $\varepsilon^*((\omega_\alpha)) = \gamma^*((\omega_\alpha))$. Since the action of G on the set of double curves of V_0 is transitive, this equality holds if they coincide on a component of \tilde{D} . Recall that, for $\alpha = \tau\rho\sigma\tau$, $C(1, 0, 0) \subset B$ and $\alpha^\wedge(E(1, 0, 0)) \subset \alpha^\wedge(B)$ form a double curve of \mathcal{L}_0 . The isomorphism κ of the identification $E(1, 0, 0) \rightarrow C(1, 0, 0)$ is given by $(X_1: X_2) \mapsto (X_2: X_1)$.

We set

$$L_m = \{ \omega \in H^0(B, \omega_B^{\otimes m}(mA)) \}^{\Gamma_2}; \kappa^*(\omega|_{C(1,0,0)}) = \omega|_{E(1,0,0)} \}.$$

By the expression (4) for $H^0(B, \omega_B^{\otimes m}(mA))$, we see easily that L_m is naturally isomorphic to $L'_m \subset F_2[X_0, X_1, X_2]$ consisting of Γ_2 -invariant homogeneous polynomials f of degree $4m$ such that $f(1, X_1, X_2)$ has no terms of degree smaller than m and $f(0, X_2, X_1)/X_1^m X_2^m (X_1 + X_2)^m = [f(1, X_1, X_2)]_m$, where $[g]_m$ denotes the homogeneous part of degree m of a polynomial g . Note that f has zero of multiplicity at least m at $(1, 0, 0)$ if and only if $f(1, X_1, X_2)$ has no terms of degree smaller than m . By the above observation, we have the following:

PROPOSITION 3.6. *$H^0(V_0, \omega_{V_0}^{\otimes m})^G$ is isomorphic to L_m by the correspondence $(\omega_\alpha)_{\alpha \in S} \mapsto \omega'_1$ where ω'_1 is the pull-back of ω_1 by the natural isomorphism $B \xrightarrow{\sim} B_1$. Hence it is also isomorphic to L'_m .*

For any $\alpha \in GL(3, F_2)$, we have $\alpha^*(f/u_0^m(dy/y \wedge dz/z)^{\otimes m}) = (\alpha^* f)/u_0^m(dy/y \wedge dz/z)^{\otimes m}$ for $f/u_0^m(dy/y \wedge dz/z)^{\otimes m} \in H^0(B, \omega_B^{\otimes m}(mA))$, where f is a homogeneous polynomial of degree $4m$. Hence, in order to determine the Γ_2 -invariant elements of $H^0(\omega_B^{\otimes m}(mA))$, we have to know those of $F_2[X_0, X_1, X_2]$.

Recall that $\lambda = (-1 + \sqrt{-7})/2$ is embedded in Z_7 so that $\lambda \equiv 0 \pmod{2}$. Hence, for $\zeta = \exp(2\pi i/7)$, $Q_2(\zeta)$ is a cubic extension of Q_2 with the relation $\zeta^3 - \lambda\zeta^2 - (1 + \lambda)\zeta - 1 = 0$. We denote by ζ_0 the modulo 2 reduction of ζ , i.e., ζ_0 is a root of the equation $X^3 + X + 1 = 0$ in $F_2[X]$.

The following method to find Γ_2 -invariant polynomials in $F_2[X_0, X_1, X_2]$ is due to Nakamura.

We set

$$\begin{aligned} Y_0 &= X_0 + \zeta_0^2 X_1 + \zeta_0 X_2, \\ Y_1 &= X_0 + \zeta_0^4 X_1 + \zeta_0^2 X_2, \\ Y_2 &= X_0 + \zeta_0 X_1 + \zeta_0^4 X_2. \end{aligned}$$

Note that this is the modulo 2 reduction of the coordinate change in Remark 2.10, since

$$\begin{bmatrix} 1 & 1 & 1 \\ \zeta_0^2 & \zeta_0^4 & \zeta_0 \\ \zeta_0 & \zeta_0^2 & \zeta_0^4 \end{bmatrix} = \begin{bmatrix} 1 & \zeta_0 & \zeta_0^2 \\ 1 & \zeta_0^2 & \zeta_0^4 \\ 1 & \zeta_0^4 & \zeta_0 \end{bmatrix}^{-1}.$$

Then we have

$$\begin{aligned} \tau(Y_0) &= \zeta_0 Y_0, & \tau(Y_1) &= \zeta_0^2 Y_1, & \tau(Y_2) &= \zeta_0^4 Y_2, \\ \sigma(Y_0) &= Y_2, & \sigma(Y_1) &= Y_0 & \text{and} & \sigma(Y_2) = Y_1. \end{aligned}$$

Thus, if a polynomial f in $\bar{F}_2[Y_0, Y_1, Y_2]$ is τ -invariant, then it is a sum of τ -invariant monomials in Y_0, Y_1 and Y_2 .

A monomial $Y_0^i Y_1^j Y_2^k$ is τ -invariant if and only if $i + 2j + 4k \equiv 0 \pmod{7}$. If it is τ -invariant, then

$$F_{i,j,k} = Y_0^i Y_1^j Y_2^k + Y_0^k Y_1^i Y_2^j + Y_0^j Y_1^k Y_2^i$$

is Γ_2 -invariant. Conversely, every Γ_2 -invariant polynomial in $\bar{F}_2[Y_0, Y_1, Y_2]$ is a linear combination of $F_{i,j,k}$'s.

PROPOSITION 3.7. *For any i, j, k with $i + 2j + 4k \equiv 0 \pmod{7}$, $F_{i,j,k}$ is in $F_2[X_0, X_1, X_2]$. Conversely, every Γ_2 -invariant polynomial in $F_2[X_0, X_1, X_2]$ is a sum of $F_{i,j,k}$'s.*

PROOF. Clearly, $F_{i,j,k} \in F_2(\zeta_0)[X_0, X_1, X_2]$. Let u be the automorphism of $F_2(\zeta_0)[X_0, X_1, X_2]$ defined by $u(X_i) = X_i$ for $i = 0, 1, 2$ and $u(\zeta_0) = \zeta_0^2$. Then, a polynomial f in $F_2(\zeta_0)[X_0, X_1, X_2]$ is in $F_2[X_0, X_1, X_2]$ if and only

if $u(f) = f$. Since $u(Y_0) = Y_1$, $u(Y_1) = Y_2$, $u(Y_2) = Y_0$, we have $u(F_{i,j,k}) = F_{i,j,k}$.

Suppose $F \in F_2[X_0, X_1, X_2]$ is Γ_2 -invariant. Since $F_2[X_0, X_1, X_2] \subset F_2(\zeta_0)[Y_0, Y_1, Y_2]$, F is written uniquely as a linear combination of $F_{i,j,k}$'s with coefficients in $F_2(\zeta_0) \setminus \{0\}$. Since $u(F_{i,j,k}) = F_{i,j,k}$, the coefficients are in $F_2 \setminus \{0\} = \{1\}$. q.e.d.

We denote by Inv_n the F_2 -vector space of Γ_2 -invariant homogeneous polynomials of degree n in $F_2[X_0, X_1, X_2]$. By the above proposition, we can easily find bases for Inv_n for small n 's as follows:

- $\text{Inv}_0 = (1)$.
- $\text{Inv}_1 = \text{Inv}_2 = \{0\}$.
- $\text{Inv}_3 = (\phi_3)$, $\phi_3 = Y_0 Y_1 Y_2$.
- $\text{Inv}_4 = (\phi_4)$, $\phi_4 = Y_0 Y_1^3 + Y_1 Y_2^3 + Y_2 Y_0^3$.
- $\text{Inv}_5 = (\phi_5)$, $\phi_5 = Y_0^3 Y_1^2 + Y_1^3 Y_2^2 + Y_2^3 Y_0^2$.
- $\text{Inv}_6 = (\phi_3^2, \phi_6)$, $\phi_6 = Y_0^5 Y_1 + Y_1^5 Y_2 + Y_2^5 Y_0$.
- $\text{Inv}_7 = (\phi_3 \phi_4, \phi_7)$, $\phi_7 = Y_0^7 + Y_1^7 + Y_2^7$.
- $\text{Inv}_8 = (\phi_4^2, \phi_3 \phi_5)$.

We can also show that Inv_{12} is generated by $\{F_{10,2,0}, F_{3,9,0}, F_{5,6,1}, F_{7,3,2}, F_{4,4,4}\}$. Hence

$$\text{Inv}_{12} = (\phi_3^4, \phi_3^2 \phi_6, \phi_3 \phi_4 \phi_5, \phi_5 \phi_7, \phi_6^2),$$

TABLE 1

f	$f(0, X_2, X_1)$	$f(1, X_1, X_2) \text{ mod}(X_1, X_2)^4$
ϕ_3	$X_1^3 + X_1 X_2^2 + X_2^3$	$1 + X_1^2 + X_1 X_2 + X_2^2 + X_1^3 + X_1^2 X_2 + X_2^3$
ϕ_4	$X_1^4 + X_1^2 X_2^2 + X_2^4$	$1 + X_1^2 + X_1 X_2 + X_2^2 + X_1^2 X_2 + X_1 X_2^2$
ϕ_5	$X_1^5 + X_1 X_2^4 + X_2^5$	$1 + X_1^2 X_2 + X_1 X_2^2$
ϕ_6	$X_1^6 + X_1^4 X_2^2 + X_2^6$	$1 + X_1^2 + X_1 X_2 + X_2^2$
ϕ_7	$X_1^7 + X_1^4 X_2^3 + X_1^2 X_2^5 + X_1 X_2^6 + X_2^7$	$1 + X_1^3 + X_1^2 X_2 + X_2^3$

TABLE 2

f	$f(0, X_2, X_1)$	$f(1, X_1, X_2) \text{ mod}(X_1, X_2)^4$
ϕ_3^2	$X_1^{12} + X_1^8 X_2^4 + X_2^{12}$	1
$\phi_6 \phi_7$	$X_1^{12} + X_1^9 X_2^3 + X_1^8 X_2^4 + X_1^6 X_2^6 + X_1^4 X_2^8 + X_1^3 X_2^9 + X_2^{12}$	$1 + X_1^3 + X_1 X_2^2 + X_2^3$
$\phi_3 \phi_4 \phi_5$	$X_1^{12} + X_1^9 X_2^3 + X_1^8 X_2^4 + X_1^6 X_2^6 + X_1^4 X_2^8 + X_1^3 X_2^9 + X_2^{12}$	$1 + X_1^3 + X_1^2 X_2 + X_2^3$
ϕ_3^4	$X_1^{12} + X_1^4 X_2^8 + X_2^{12}$	1
$\phi_6 \phi_3^2$	$X_1^{12} + X_1^{10} X_2^2 + X_1^8 X_2^4 + X_1^6 X_2^6 + X_1^4 X_2^8 + X_1^2 X_2^{10} + X_2^{12}$	$1 + X_1^2 + X_1 X_2 + X_2^2$

since $F_{10,2,0} = \phi_6^2$, $F_{3,9,0} = \phi_5\phi_7 + \phi_6^2 + \phi_3^2\phi_8$, $F_{5,6,1} = \phi_3\phi_4\phi_5 + \phi_3^4 + \phi_3^2\phi_6$, $F_{7,3,2} = \phi_3^2\phi_6$ and $F_{4,4,4} = \phi_3^4$.

In order to determine L'_m for $m = 2, 3$, we provide the Tables 1 and 2 of $f(0, X_2, X_1)$ and $f(1, X_1, X_2)$ for $f = \phi_i$ and each element of the basis for Inv_{12} . In the tables, we omit the part of degree greater than 3 of $f(1, X_1, X_2)$.

PROPOSITION 3.8. *We have $L'_2 = (\phi_4^2 + \phi_3\phi_5)$ and $L'_3 = (\phi_5\phi_7 + \phi_3^4, \phi_3\phi_4\phi_5 + \phi_6^2)$. In particular, $\dim H^0(V_0, \omega_{V_0}^{\otimes 2})^G = 1$ and $\dim H^0(V_0, \omega_{V_0}^{\otimes 3})^G = 2$.*

PROOF. The second assertion follows from the first by Proposition 3.6. In $\text{Inv}_8 \setminus \{0\}$, only $\phi_4^2 + \phi_3\phi_5$ has zero of multiplicity 2 at $(1, 0, 0)$. For $f = \phi_4^2 + \phi_3\phi_5$, we calculate easily by Table 1 that $[f(1, X_1, X_2)]_2 = f(0, X_2, X_1)/X_1^2X_2^2(X_1 + X_2)^2 = X_1^2 + X_1X_2 + X_2^2$. Hence L'_2 is generated by $\phi_4^2 + \phi_3\phi_5$.

From Table 2, the F_2 -vector space $\{f \in \text{Inv}_{12}; f \text{ has zero of multiplicity } 3 \text{ at } (1, 0, 0)\}$ is of dimension 3 and is generated by $\{\phi_6^2 + \phi_3^4, \phi_5\phi_7 + \phi_3^4, \phi_3\phi_4\phi_5 + \phi_6^2\}$. Hence it is easy to see that $L'_3 = (\phi_5\phi_7 + \phi_3^4, \phi_3\phi_4\phi_5 + \phi_6^2)$. Actually, we have

$$f(0, X_2, X_1)/X_1^3X_2^3(X_1 + X_2)^3 = [f(1, X_1, X_2)]_3 = X_1^3 + X_1X_2^2 + X_2^3$$

for $f = \phi_5\phi_7 + \phi_3^4$, and

$$f(0, X_2, X_1)/X_1^3X_2^3(X_1 + X_2)^3 = [f(1, X_1, X_2)]_3 = X_1^3 + X_1^2X_2 + X_2^3$$

for $f = \phi_3\phi_4\phi_5 + \phi_6^2$.

q.e.d.

We now prove the following:

THEOREM 3.9. *For the minimal resolution Y'_η of $Y_\eta = V_\eta/G$, we have $P_2(Y'_\eta) = P_3(Y'_\eta) = 1$. We can choose as generators of $H^0(Y'_\eta, \omega_{Y'_\eta}^{\otimes 2})$ and $H^0(Y'_\eta, \omega_{Y'_\eta}^{\otimes 3})$, the elements which corresponds to the Γ_2 -invariant polynomials $\phi_4^2 + \phi_3\phi_5$ and $\phi_3^4 + \phi_3\phi_4\phi_5 + \phi_5\phi_7 + \phi_6^2$ by the modulo 2 reduction, respectively.*

PROOF. We have $P_2(Y'_\eta) = 1$ by Propositions 3.5, 3.6 and 3.8. By Proposition 3.8, $H^0(Y'_\eta, \omega_{Y'_\eta}^{\otimes 2})$ is generated by the lifting of the element of $H^0(V_0, \omega_{V_0}^{\otimes 2})^G$ which corresponds to $\phi_4^2 + \phi_3\phi_5$.

By Theorem 2.9 and Remark 2.10, Y has cyclic quotient singularity of type $(7, 3)$ along \bar{q} , and it is minimally resolved simultaneously in Y' .

Let s be a section of $\omega_{Y''}^{\otimes 3}$, where Y'' is the smooth part $Y \setminus \{\bar{p}, \bar{q}, w\}$ of Y . For the resolution of Y_η at q , we define the exceptional divisors D_1, D_2, D_3 in Y'_η and integers a, b, c and $d_1, d_2, d_3 \geq 0$ similarly as in the proof of Proposition 3.5. Then we have

$$7a = 3(d_1 - 3) + 2d_2 + d_3,$$

$$7b = 2(d_1 - 3) + 6d_2 + 3d_3,$$

$$7c = (d_1 - 3) + 3d_2 + 5d_3.$$

Hence $b, c \geq 0$ and $a \geq -1$. In other words, s is regular at the divisors D_2, D_3 and may have a pole of order at most one along D_1 .

Let L_i be the intersection of the closure of D_i with Y'_0 for $i = 1, 2, 3$. Then $L = L_1 \cup L_2 \cup L_3$ is the exceptional curve of $\bar{q}(0) \in Y_0$. Let $U \subset Y'_0$ be a smooth neighborhood of L and let ω_0 be a rational section of $\omega_{Y'_0}^{\otimes 3}$ which is regular outside L . Then, as in the case of the generic fiber, ω_0 may have a pole of order at most one along L_1 . When ω_0 is represented by an element of $H^0(B, \omega_B^{\otimes 3}(3A))^{F_2}$, its regularity at L_1 is examined as follows:

Let $f(Y_0, Y_1, Y_2)$ be the corresponding Γ_2 -invariant homogeneous polynomial of degree 12 in Y_i 's. We take the local coordinate $(y_1, y_2) = (Y_1/Y_0, Y_2/Y_0)$ of the point $(1: \zeta_0: \zeta_0^2) \in \mathbf{P}^2_{F_2} = \text{Proj } \bar{F}_2[X_0, X_1, X_2]$. Then the action of τ is given by $(y_1, y_2) \mapsto (\zeta_0 y_1, \zeta_0^3 y_2)$ (cf. Remark 2.10). In the resolution, L is covered by four affine open sets with coordinates $(y_1^7, y_1^{-3}y_2)$, $(y_1^3y_2^{-1}, y_1^{-2}y_2^3)$, $(y_1^2y_2^{-3}, y_1^{-1}y_2^5)$ and $(y_1y_2^{-5}, y_2^7)$, where the second and the third coordinates are of the neighborhoods of $L_1 \cap L_2$ and $L_2 \cap L_3$, respectively. The divisor L_1 is described as the line $(s = 0)$ with respect to the coordinate $(s, t) = (y_1^7, y_1^{-3}y_2)$. ω_0 is equal to $v \cdot f(1, y_1, y_2)(dy_1 \wedge dy_2)^{\otimes 3}$ for a non-vanishing regular function v on U . In view of the equality $dy_1 \wedge dy_2 = (1/7)s^{-3/7}ds \wedge dt$, we see that ω_0 has a pole at L_1 if and only if $s^{-9/7}g(s, t)$ has a pole along $(s = 0)$, where $g(s, t) = f(1, y_1, y_2)$.

Among τ -invariant monomials of degree 12 in Y_i 's only $s^{-9/7}g(s, t)$ for $Y_0^{10}Y_1^2$ has a pole along $(s = 0)$. Hence ω_0 's which correspond to $\phi_5\phi_7 + \phi_3^4 = F_{10,2,0} + F_{3,9,0} + F_{7,3,2} + F_{4,4,4}$ and $\phi_3\phi_4\phi_5 + \phi_6^2 = F_{10,2,0} + F_{5,6,1} + F_{7,3,2} + F_{4,4,4}$ have a pole along L_1 , while ω_0 for $\phi_3^4 + \phi_3\phi_4\phi_5 + \phi_5\phi_7 + \phi_6^2 = F_{3,9,0} + F_{5,6,1}$ does not.

Let ω_γ be an element of $H^0(Y''_\gamma, \omega_{Y''_\gamma}^{\otimes 3})$ which has nontrivial reduction ω_0 to Y''_0 . Then ω_γ has a pole at D_1 , if so does ω_0 at L_1 . Hence, by Theorem 3.4, there exists $\omega \in H^0(Y''_\gamma, \omega_{Y''_\gamma}^{\otimes 3})$ with a pole along D_1 . Since D_1 is a nonsingular rational curve and $D_1^2 = -3$, we have $\omega_{Y''_\gamma}^{\otimes 3}(D_1)|_{D_1} \simeq \mathcal{O}_{D_1}$. Hence $H^0(Y''_\gamma, \omega_{Y''_\gamma}^{\otimes 3})$ is of codimension one in $H^0(Y''_\gamma, \omega_{Y''_\gamma}^{\otimes 3}(D_1))$, which is isomorphic to $H^0(Y''_\gamma, \omega_{Y''_\gamma}^{\otimes 3})$, since the other singularities p_0, p_1, p_2 are rational double points. Since $H^0(Y''_\gamma, \omega_{Y''_\gamma}^{\otimes 3}) \simeq H^0(V_\gamma, \omega_{V_\gamma}^{\otimes 3})^\sigma$ is of dimension two by Theorem 3.4 and Proposition 3.8, we have $\dim H^0(Y''_\gamma, \omega_{Y''_\gamma}^{\otimes 3}) = 1$. q.e.d.

REMARK 3.10. The Γ_2 -invariant polynomials $f_2 = \phi_4^2 + \phi_3\phi_5$ and $f_3 = \phi_3^4 + \phi_3\phi_4\phi_5 + \phi_5\phi_7 + \phi_6^2$ are equal to $F_{2,6,0} + F_{4,3,1}$ and $F_{3,9,0} + F_{5,6,1}$, respectively. By expressing these polynomials in terms of the coordinates at $L_1 \cap L_2$ and $L_2 \cap L_3$ in the proof of the above theorem, we see that gene-

rators of $H^0(Y'_i, \omega_{Y'_i}^{\otimes m})$ for $m = 2, 3$ and their modulo 2 reductions have no zero along D_i 's and L_i 's, respectively.

4. The minimal resolution of $Y(\bar{Q}_2)$. In this section, we denote by X the normal surface $Y(\bar{Q}_2)$. By Theorem 2.9, X has cyclic quotient singularities p_0, p_1, p_2 and q . Let $\pi: \tilde{X} \rightarrow X$ be the minimal resolution of these singularities. Hence $\tilde{X} = Y'(\bar{Q}_2)$ for Y' in Remark 2.10. We denote by D_1, \dots, D_9 the irreducible divisors of \tilde{X} such that $\pi^{-1}(q) = D_1 + D_2 + D_3$, $\pi^{-1}(p_0) = D_4 + D_5$, $\pi^{-1}(p_1) = D_6 + D_7$ and $\pi^{-1}(p_2) = D_8 + D_9$. We assume $D_1^2 = -3$ and $D_1 \cap D_3 = \emptyset$ as in Section 3. Hence we have $D_i^2 = -2$ for $2 \leq i \leq 9$. Let K_X be a canonical divisor of X . Since X has only cyclic quotient singularities, K_X is a \mathbf{Q} -Cartier divisor. In fact, $21K_X$ is a Cartier divisor.

PROPOSITION 4.1. *The Chern numbers of the nonsingular surface \tilde{X} are $c_1^2(\tilde{X}) = 0$ and $c_2(\tilde{X}) = 12$.*

PROOF. Let $K_{\tilde{X}}$ be the canonical divisor of \tilde{X} which is equal to K_X on $X \setminus \{p_0, p_1, p_2, q\}$. Then $\pi^*K_X - K_{\tilde{X}}$ is a \mathbf{Q} -divisor supported in $D_1 \cup \dots \cup D_9$, i.e., $\pi^*K_X - K_{\tilde{X}} = a_1D_1 + \dots + a_9D_9$ for some $a_1, \dots, a_9 \in \mathbf{Q}$. Since D_i 's are nonsingular rational curves, we have $(\pi^*K_X - K_{\tilde{X}}) \cdot D_i = -K_{\tilde{X}} \cdot D_i = 2 + D_i^2$ for every i .

Then we see easily that

$$\pi^*K_X - K_{\tilde{X}} = (3/7)D_1 + (2/7)D_2 + (1/7)D_3.$$

In particular, we have

$$(1) \quad K_X^2 - K_{\tilde{X}}^2 = (\pi^*K_X - K_{\tilde{X}}) \cdot K_{\tilde{X}} = 3/7.$$

On the other hand, by Theorem 2.9, there exists a finite morphism $f: V(\bar{Q}_2) \rightarrow X$ of degree 168 ramified only at $\{p_0, p_1, p_2, q\}$. Since $c_1^2(V(\bar{Q}_2)) = 72$, we have

$$(2) \quad K_X^2 = 72/168 = 3/7.$$

Hence $c_1^2(\tilde{X}) = K_{\tilde{X}}^2 = 0$ by (1) and (2).

For $c_2(\tilde{X})$, we may let $\bar{Q}_2 = C$ and calculate it as the topological Euler number $e(\tilde{X})$. By Theorem 2.9, $f^{-1}(p_i)$ for $i = 0, 1, 2$ and $f^{-1}(q)$ consist of $168/3 = 56$ and $168/7 = 24$ points, respectively. Since $c_2(V(\bar{Q}_2)) = 24$, we have

$$\begin{aligned} c_2(\tilde{X}) &= (c_2(V(\bar{Q}_2)) - \sum f^{-1}(\{p_0, p_1, p_2, q\})/168 + e(\pi^{-1}(\{p_0, p_1, p_2, q\}))) \\ &= (24 - (3 \cdot 56 + 24))/168 + (3 \cdot 3 + 4) = 12. \end{aligned} \quad \text{q.e.d.}$$

REMARK 4.2. The above proposition implies $\chi(\mathcal{O}_{\tilde{X}}) = 1$ by Noether's

formula. In fact, we have $p_g(\tilde{X}) = q(\tilde{X}) = 0$, since X has a finite covering $M(\mathbb{Q}_2) \rightarrow X$ from Mumford's fake projective plane $M(\mathbb{Q}_2)$ ramified only at finite points.

PROPOSITION 4.3. *\tilde{X} is a minimal elliptic surface, i.e., the Kodaira dimension of \tilde{X} is equal to one.*

PROOF. Suppose \tilde{X} were of general type, and let X' be its minimal model. By the plurigenus formula, we have $P_m(\tilde{X}) = (m(m - 1)/2)K_{X'}^2 + \chi(\mathcal{O}_{\tilde{X}})$ for $m \geq 2$. In particular $P_2(\tilde{X}) \geq 2$. This contradicts Theorem 3.9.

If \tilde{X} were of Kodaira dimension zero, then \tilde{X} is either a $K3$ surface or an Enriques surface, since $q(\tilde{X}) = 0$. These are impossible since $p_g(\tilde{X}) = 0$ and $P_3(\tilde{X}) = 1$ by Theorem 3.9.

Hence \tilde{X} is an elliptic surface and it is minimal by $K_{\tilde{X}}^2 = 0$. q.e.d.

Recall that the \mathbb{Z}_2 -scheme Y' is regular outside the point w in the closed fiber. For each integer m , we denote by $\omega_{Y'}^{\otimes m}$ the maximal torsion-free extension of $\omega_{Y'}^{\otimes m} \setminus \{w\}$ to Y' . We fix sections F_2 and F_3 of $\omega_{Y'}^{\otimes 2}$ and $\omega_{Y'}^{\otimes 3}$ with non-trivial modulo 2 reductions, respectively, which exist by Theorem 3.9. Let E' and E'' be the effective divisors (F_2) and (F_3) of Y' , respectively. Clearly, $3E'$ and $2E''$ are linearly equivalent.

LEMMA 4.4. *E' and E'' are disjoint.*

PROOF. Let $\pi_0: Y'_0 \rightarrow Y_0$ be the natural morphism. We denote by \bar{E}'_0 and \bar{E}''_0 the images by π_0 of the divisors $E'_0 = E' \cap Y'_0$ and $E''_0 = E'' \cap Y'_0$, respectively.

By the definition of E' and E'' and by Theorem 3.9, \bar{E}'_0 and \bar{E}''_0 correspond to the $\bar{\Gamma}_2$ -invariant polynomials $f_2 = \phi_4^2 + \phi_3\phi_5$ and $f_3 = \phi_3^4 + \phi_3\phi_4\phi_6 + \phi_5\phi_7 + \phi_6^2$, respectively. Let \tilde{E}'_0 and \tilde{E}''_0 be the pull-backs of \bar{E}'_0 and \bar{E}''_0 , respectively, by the natural surjective morphism $h: B \rightarrow Y_0$. By Tables 1 and 2, the restrictions of \tilde{E}'_0 and \tilde{E}''_0 to the rational curve $C(1, 0, 0) \subset B$ is defined by $X_1^2 + X_1X_2 + X_2^2$ and $X_1X_2(X_1 + X_2)$, respectively. In particular, they do not intersect each other on the curve. Since G acts transitively on the set of double curves of V_0 , and since B is isomorphic to the component B_1 of V_0 , \tilde{E}'_0 and \tilde{E}''_0 do not intersect each other on the fourteen rational curves in Figure 1 in Section 1. Since the complement of the union of the curves in B is an affine open set, \tilde{E}'_0 and \tilde{E}''_0 have no common components. E'_0 and E''_0 also have no common components, since they do not contain L_i for $i = 1, 2, 3$ by Remark 3.10, and since E'_0 does not have any zero on the other exceptional curves of π_0 .

On the other hand, f_2 and f_3 have zeros of multiplicities 2 and 3, re-

spectively, at the seven F_2 -rational points of $P_{F_2}^2$. Since B is the blowing-up of $P_{F_2}^2$ at the seven F_2 -rational points, the intersection number $\tilde{E}' \cdot \tilde{E}''$ is $\deg f_2 \cdot \deg f_3 - 7 \cdot 2 \cdot 3 = 96 - 42 = 54$. Since $Y_0 \setminus h(C(1, 0, 0))$ is smooth except at the cyclic quotient singularity $\bar{q}(0)$, we can consider the intersection number $\bar{E}' \cdot \bar{E}'' = 54/21 = 18/7$, since h is of degree 21. As in the proof of Proposition 4.1, we have

$$\begin{aligned} \pi_0^* \bar{E}'_0 - E'_0 &= 2((3/7)L_1 + (2/7)L_2 + (1/7)L_3), \\ \pi_0^* \bar{E}''_0 - E''_0 &= 3((3/7)L_1 + (2/7)L_2 + (1/7)L_3) \quad \text{and} \\ \bar{E}'_0 \cdot \bar{E}''_0 - E'_0 \cdot E''_0 &= 2 \cdot 3 \cdot 3/7 = 18/7. \end{aligned}$$

Hence $E'_0 \cdot E''_0 = 0$. We have $E'_0 \cap E''_0 = \emptyset$, since they have no common components. This implies $E' \cap E'' = \emptyset$. q.e.d.

Let $\kappa: Y' \rightarrow P_{z_2}^1$ be the morphism defined by (F_2^3, F_3^2) .

PROPOSITION 4.5. *The induced morphism $\kappa^1_{\bar{q}_2}: \tilde{X} \rightarrow P^1_{\bar{q}_2}$ of the geometric fibers is the elliptic fibration of \tilde{X} . It has just two multiple fibers $3E'_{\bar{q}_2}$ and $2E''_{\bar{q}_2}$, where $E'_{\bar{q}_2}$ and $E''_{\bar{q}_2}$ are the restrictions of E' and E'' to \tilde{X} , respectively.*

PROOF. Let $f: \tilde{X} \rightarrow P^1_{\bar{q}_2}$ be the elliptic fibration, and let $m_1 C_1, \dots, m_n C_n$ be its multiple fibers. By Kodaira's canonical bundle formula [Ko2, Th. 12], we have

$$K_{\tilde{X}} \sim f^{-1}(-x_0) + \sum_{i=1}^n (m_i - 1)C_i,$$

where x_0 is a point of $P^1_{\bar{q}_2}$, since $\deg K_{P^1} + \chi(\mathcal{O}_{\tilde{X}}) = -1$. Since $2K_{\tilde{X}} \sim (n - 2)f^{-1}(x_0) + \sum_{i=1}^n (m_i - 2)C_i$, we have $\dim |2K_{\tilde{X}}| = n - 2$. Hence $n = 2$ by Theorem 3.9. Since $E'_{\bar{q}_2}$ is a unique effective bicanonical divisor, we have $E'_{\bar{q}_2} = (m_1 - 2)C_1 + (m_2 - 2)C_2$. If $m_1, m_2 \geq 3$, $3K_{\tilde{X}} \sim f^{-1}(x_0) + (m_1 - 3)C_1 + (m_2 - 3)C_2$ and hence $\dim |3K_{\tilde{X}}| = 1$. This contradicts Theorem 3.9. Hence we may assume $m_1 = 2$. Since $(m_2 - 2)C_2 = E'_{\bar{q}_2}$, we have $m_2 > 2$. Hence $3K_{\tilde{X}} \sim E''_{\bar{q}_2} = C_1 + (m_2 - 3)C_2$. Since $E'_{\bar{q}_2} \cap E''_{\bar{q}_2} = \emptyset$ by Lemma 4.4, we have $m_2 = 3$.

Thus we have $E'_{\bar{q}_2} = C_2$, $E''_{\bar{q}_2} = C_1$ and $f^{-1}(x_0) \sim 3E'_{\bar{q}_2} \sim 2E''_{\bar{q}_2}$. Hence f is equal to $\kappa_{\bar{q}_2}$ up to automorphism of $P^1_{\bar{q}_2}$. q.e.d.

The connected curves $D_2 \cup D_3$, $D_4 \cup D_5$, $D_6 \cup D_7$ and $D_8 \cup D_9$ are unions of (-2) -curves. Hence they are mapped to points in $P^1_{\bar{q}_2}$ by $\kappa_{\bar{q}_2}$. We denote $y = \kappa_{\bar{q}_2}(D_2 \cup D_3)$ and $z_i = \kappa_{\bar{q}_2}(D_{4+2i} \cup D_{5+2i})$ for $i = 0, 1, 2$.

PROPOSITION 4.6. *$E'_{\bar{q}_2}$, $E''_{\bar{q}_2}$, $D_2 \cup D_3$, $D_4 \cup D_5$, $D_6 \cup D_7$ and $D_8 \cup D_9$ are mapped to distinct points in $P^1_{\bar{q}_2}$ by $\kappa_{\bar{q}_2}$.*

PROOF. By definition, $\kappa_{\bar{Q}_2}(E'_R) = (0:1)$ and $\kappa_{\bar{Q}_2}(E''_R) = (1:0)$. By Remark 3.10, the modulo 2 reduction $L_2 \cup L_3$ of $D_2 \cup D_3$ is contained in neither E'_0 nor E''_0 . Hence the specialization of y in $\mathbf{P}^1_{F_2}$ is neither $(1:0)$ nor $(0:1)$. As we saw immediately before Theorem 2.9, there exists a \mathbf{Z}_2 -morphism $\text{Spec } \mathbf{Z}_2[\varepsilon] \rightarrow \mathcal{X}$ which is fixed by $\tau\rho$, and the induced $\mathbf{Q}_2[\varepsilon]$ -valued point in Y splits to p_0, p_1, p_2 in $Y(\bar{\mathbf{Q}}_2)$ and the image of the closed point is the triple point P of \mathcal{X}_0 . As we saw in the proof of Lemma 4.4, the pull-back of \bar{E}' and \bar{E}'' to $C(1, 0, 0)$ is defined by $X_1^2 + X_1X_2 + X_2^2$ and $X_1X_2(X_1 + X_2)$, respectively. Hence we have $\bar{P} \in \bar{E}''$ and $\bar{P} \notin \bar{E}'$, where \bar{P} is the image of P in Y . Since $D_4 \cup D_5, D_6 \cup D_7$ and $D_8 \cup D_9$ are the exceptional curves of p_0, p_1 and p_2 , respectively, the specialization of z_i 's are all $(1:0)$. We get the following diagram after the base extension in Remark 2.10:

$$\begin{array}{ccccc}
 \text{Spec } K[\varepsilon] \hookrightarrow \text{Spec } R[\varepsilon] & \rightarrow & V'_R & \hookrightarrow & V_R \\
 \downarrow \mu & & \downarrow & & \downarrow \\
 & & Y_R \setminus \bar{E}'_R & \hookrightarrow & Y_R \leftarrow Y'_R \\
 & & \downarrow F_3^2/F_2^3 & & \downarrow \kappa_R \\
 \text{Spec } K[t] \hookrightarrow \text{Spec } R[t] = A^1_R & \hookrightarrow & & \hookrightarrow & \mathbf{P}^1_R.
 \end{array}$$

Here K is the quotient field of $R = \mathbf{Z}_2[\zeta, \omega]$, V'_R a neighborhood of $P_1 \in V_R$, \bar{E}'_R the image of E'_R in Y_R and $A^1_R = \mathbf{P}^1_R \setminus \kappa_R(E'_R)$. It suffices to show that the K -homomorphism $\mu^*: K[t] \rightarrow K[\varepsilon]$ is surjective, since then the image of μ is a separable point of degree 3 while $(1:0)$ is the K -rational point $t = 0$. By the notation in Remark 2.10, we get the following sequence of formal completions of local rings:

$$R[t] \rightarrow R[[T_0, T_1^3, T_2^3, T_1T_2]] \rightarrow R[[T_0, T_1, T_2]] \xrightarrow{l} R[\varepsilon],$$

where T_0, T_1, T_2 have a relation $T_0^3 + T_1^3 + T_2^3 - 3T_0T_1T_2 = 27\varepsilon^3$. l is given by $l(T_0) = 3\varepsilon$ and $l(T_1) = l(T_2) = 0$. The image of t in $R[[T_0, T_1, T_2]]$ is equal to F_3^2/F_2^3 . Since Y is a Gorenstein scheme and since F_2 and F_3 are sections of $\omega_Y^{\otimes m}$ for $m = 2, 3$, respectively, we may regard F_2 and F_3 as elements of $R[[T_0, T_1^3, T_2^3, T_1T_2]]$. By the restriction of the polynomials f_2 and f_3 to $C(1, 0, 0) \subset B$, we see that $F_3 \in (T_0, T_1, T_2) \setminus (T_0, T_1, T_2)^2$ and F_2 is a unit. Hence F_3 has a unit coefficient for T_0 , and hence F_3^2/F_2^3 has a unit coefficient for T_0^2 . This implies that the image of t in $R[\varepsilon]$ is outside R . Hence μ^* is surjective. q.e.d.

Now we can determine the types of the singular fibers:

THEOREM 4.7. *The elliptic fibration $\kappa_{\bar{Q}_2}: \tilde{X} \rightarrow \mathbf{P}^1_{\bar{Q}_2}$ has singular fibers at $\{(1:0), (0:1), y, z_0, z_1, z_2\} \subset \mathbf{P}^1_{\bar{Q}_2}$ and smooth elsewhere. The singular*

fibers over z_0, z_1, z_2 and y are not multiple and are of type I_3 in the notation of [Ko1, Th. 6.2]. The fibers over $(1:0)$ and $(0:1)$ are $2E''_{\bar{q}_2}$ and $3E'_{\bar{q}_2}$, respectively, where $E''_{\bar{q}_2}$ and $E'_{\bar{q}_2}$ are smooth elliptic curves.

PROOF. Each of the fibers over z_0, z_1, z_2 and y contains a union of two (-2) -curves intersecting each other transversally at one point. Hence they are not of type II nor III. Hence the Euler number of the non-elliptic fiber is at least three and is equal to three if and only if it is of type I_3 . Now we apply Kodaira's formula for the second Betti number of an elliptic surface [Ko1, Th. 12.2]. Since $c_2(\tilde{X}) = 12$ by Proposition 4.1, all these fibers are of type I_3 and the other fibers are elliptic curves. The multiple fibers are only $2E''_{\bar{q}_2}$ and $3E'_{\bar{q}_2}$ by Proposition 4.5. q.e.d.

REFERENCES

- [BPV] W. BARTH, C. PETERS AND A. VAN DE VEN, Compact Complex Surfaces, Springer-Verlag, Berlin, Heidelberg, New York, 1984.
- [Ko1] K. KODAIRA, On compact complex analytic surfaces I-III, Ann. of Math. 71 (1960), 111-152; 77 (1963), 563-626; 78 (1963), 1-40.
- [Ko2] K. KODAIRA, On the structure of compact complex analytic surfaces, I, Amer. J. Math. 86 (1964), 751-798.
- [Ku] A. KURIHARA, Construction of p -adic unit balls and the Hirzebruch proportionality, Amer. J. Math. 102 (1980), 565-648.
- [Mum] D. MUMFORD, An algebraic surface with K ample, $(K^2)=9$, $p_g=q=0$, in Contribution to Algebraic Geometry, Johns Hopkins Univ. Press, 233-244, 1979.
- [Mus] G. A. MUSTAFIN, Nonarchimedean uniformization, Math. USSR, Sbornik, 34 (1978), 187-214.

MATHEMATICAL INSTITUTE
TÔHOKU UNIVERSITY
SENDAI, 980
JAPAN