

SUPERPOSITION OPERATORS ON DIRICHLET SPACES

Dedicated to Professor Ronald Getoor on his seventy-fifth birthday

PATRICK J. FITZSIMMONS

(Received November 11, 2002, revised October 3, 2003)

Abstract. In the context of a strongly local Dirichlet space we show that if a function mapping the real line to itself (and fixing the origin) operates by composition on the left to map the Dirichlet space into itself, then the function is necessarily locally Lipschitz continuous. If, in addition, the Dirichlet space contains unbounded elements, then the function must be globally Lipschitz continuous. The proofs rely on a co-area formula for condenser potentials.

1. Introduction. Let $(\mathcal{E}, \mathcal{D})$ be a strongly local regular Dirichlet form as in [13]. Specifically, let E be a locally compact second countable metric space, and m a Radon measure on E with full support. The domain \mathcal{D} of the Dirichlet form \mathcal{E} is a vector space contained and dense in $L^2(m)$. The symmetric bilinear form $\mathcal{E} : \mathcal{D} \times \mathcal{D} \rightarrow \mathbf{R}$ is non-negative definite, and the augmented form $\mathcal{E}_1 : (u, v) \mapsto \mathcal{E}(u, v) + (u, v)_m$ is a Hilbertian inner product on \mathcal{D} . (Here $(u, v)_m := \int_E uv dm$ is the inner product in $L^2(m)$; the associated norm is $\|u\|_2$.) The form \mathcal{E} is *closable* in the sense that if $\{u_n\}$ is an \mathcal{E} -Cauchy sequence with $\lim_n \|u_n\|_2 = 0$, then $\lim_n \mathcal{E}(u_n, u_n) = 0$ as well. Furthermore, $(\mathcal{E}, \mathcal{D})$ enjoys the following contraction property: If $K : \mathbf{R} \rightarrow \mathbf{R}$ satisfies the conditions

$$(1.1) \quad K(0) = 0$$

and

$$(1.2) \quad |K(t) - K(s)| \leq M|t - s| \quad \text{for all } s, t \in \mathbf{R},$$

then

$$(1.3) \quad K \circ u \in \mathcal{D} \quad \text{for all } u \in \mathcal{D},$$

and

$$(1.4) \quad \mathcal{E}(K \circ u, K \circ u) \leq M^2 \cdot \mathcal{E}(u, u) \quad \text{for all } u \in \mathcal{D}.$$

In the sequel we shall describe the property (1.3) by saying that “ K operates on \mathcal{D} ”. Notice that the elements of \mathcal{D} are actually classes of functions equal a.e. with respect to m , but that the superposition operation $u \mapsto K \circ u$ preserves m -classes. The modifier “regular” indicates that $\mathcal{D} \cap C_0(E)$ is \mathcal{E}_1 -dense in \mathcal{D} and uniformly dense in $C_0(E)$. (Here, $C_0(E)$ denotes the class of real-valued continuous functions on E with compact support.) Finally, “strongly local” means that whenever u and v are elements of \mathcal{D} such that the measures $u \cdot m$ and $v \cdot m$ are of

compact support and v is constant on a neighborhood of the support of $u \cdot m$, then necessarily $\mathcal{E}(u, v) = 0$.

Our goal in this paper is to examine the following question: If K operates on \mathcal{D} , must it be Lipschitz continuous, at least locally? This problem has been posed and solved by Marcus and Mizel [17] in the context of Sobolev spaces $W^{1,p}(\Omega)$ ($1 \leq p < \infty$), where the bounded Euclidean domain $\Omega \subset \mathbf{R}^d$ is assumed to satisfy a cone condition. (The critical case $p = d \geq 2$, left open in [17], was treated by Bourdaud [5].) Their work was extended to general Euclidean domains of finite measure by Labutin [15]. (See [2] for a monograph treatment of these and related questions.) The intersection of our context with that of these authors is the case $p = 2$, corresponding to Brownian motion. They found that if K operates on $W^{1,2}(\Omega)$, then K is locally Lipschitz (i.e., K is absolutely continuous and $K' \in L_{\text{loc}}^\infty(\mathbf{R})$); if, in addition, $W^{1,2}(\Omega)$ contains essentially unbounded elements, then $K' \in L^\infty(\mathbf{R})$. We shall see that these two conclusions remain valid for superpositions operating on the Dirichlet space of a general diffusion process. (The second conclusion requires a mild secondary hypothesis in the general case.)

Our methods are quite different from those used in the papers cited above and the constructions involved are less intricate. This simplification is due to a systematic use of condenser potentials, and to a related variant of the co-area formula that is of independent interest. This formula, an explicit instance of the “energy image density” formula of Bouleau and Hirsch [4], was suggested by a calculation found in recent work of Fukushima and Ying [14].

Our arguments are largely probabilistic, and in the balance of this section we shall describe the probabilistic context in which we shall be working, and we shall review some relevant facts. The precise statement of our main result, and its proof, will occupy the next section.

Associated with the Dirichlet form $(\mathcal{E}, \mathcal{D})$ is a symmetric Markov process $X = (\Omega, \mathcal{F}, \mathcal{F}_t, \theta_t, X_t, \mathbf{P}^x)$, with symmetry measure m . The process X is a Hunt process; in particular, the transition semigroup of X , defined by

$$(1.5) \quad P_t f(x) := \mathbf{P}^x[f(X_t)], \quad t \geq 0,$$

maps $b\mathcal{B}(E)$ (the class of bounded real-valued $\mathcal{B}(E)$ -measurable functions on E) into itself. Here $\mathcal{B}(E)$ denotes the class of Borel subsets of E and $\mathbf{P}^x[Z]$ denotes the expectation of the random variable Z with respect to the measure \mathbf{P}^x . As noted already, the semigroup (P_t) is m -symmetric:

$$(1.6) \quad (f, P_t g)_m = (P_t f, g)_m, \quad f, g \in p\mathcal{B}(E).$$

Because $(\mathcal{E}, \mathcal{D})$ is strongly local, X is a *diffusion* in the sense that

- (i) The \mathbf{P}^m -completion $(\mathcal{F}_t)_{t \geq 0}$ of the natural filtration $\sigma\{X_s; 0 \leq s \leq t\}$, $t \geq 0$, is quasi-left-continuous and the lifetime of X , denoted ζ , is an (\mathcal{F}_t) *predictable* stopping time;
- (ii) $t \mapsto X_t$ is continuous on $[0, \zeta[$, \mathbf{P}^m -a.s.

Here $\mathbf{P}^m[\cdot] := \int_E \mathbf{P}^x[\cdot] m(dx)$. Also, ζ , the lifetime of X , is the time at which X retires to the cemetery state Δ , which has been adjoined to E (as the point at infinity if E is not

compact) to allow for the possible mass defect $P_t 1_E(x) < 1$. By convention any function (resp. measure) defined on E (resp. \mathcal{E}) is extended to the cemetery state Δ by declaring its value at Δ (resp. $\{\Delta\}$) to be 0. Consequently, every (\mathcal{F}_t) -stopping time is \mathbf{P}^m -predictable, and every (\mathcal{F}_t) -martingale has continuous paths (\mathbf{P}^m -a.s.). See [21, §47]. As general references on Markov processes, the reader can consult [3] and [21].

Let us now recall more about the specifics of the association between $(\mathcal{E}, \mathcal{D})$ and X . Because of (1.6), (P_t) restricted to $b\mathcal{B}(E) \cap L^2(m)$ extends uniquely to a strongly continuous contraction semigroup of self-adjoint operators in $L^2(m)$. The Dirichlet form \mathcal{E} is given by the formula

$$(1.8) \quad \mathcal{E}(u, v) := \lim_{t \downarrow 0} t^{-1}(u, v - P_t v)_m,$$

and its domain is specified by

$$(1.9) \quad \mathcal{D} := \left\{ u \in L^2(m); \sup_{t > 0} t^{-1}(u, u - P_t u)_m < \infty \right\}.$$

We shall often write $\mathcal{E}(u)$ instead of $\mathcal{E}(u, u)$.

REMARK 1.10. In the discussion at hand, there is nothing to be gained by working in the more general context of quasi-regular Dirichlet forms, as developed in [16]. Indeed, the main result of [7] asserts that every quasi-regular Dirichlet form is quasi-homeomorphic to a regular Dirichlet form, and it is a simple matter to check that our hypotheses and our conclusions are preserved by quasi-homeomorphisms.

The *hitting time* $\inf\{t > 0; X_t \in B\}$ of $B \in \mathcal{B}(E)$ is denoted by T_B ; the *exit time* from B , namely T_{B^c} , is denoted by $\tau(B)$. An increasing sequence $\{B_n\}$ of Borel subsets of E is a *nest* provided $\mathbf{P}^m[\lim_n \tau(B_n) < \zeta] = 0$. The reader is referred to [16, Lemma IV. 4.5] for a characterization of this notion in terms of the Dirichlet form $(\mathcal{E}, \mathcal{D})$.

Each element $u \in \mathcal{D}$ admits a Borel measurable m -modification \tilde{u} such that $t \mapsto \tilde{u}(X_t)$ is right continuous on $[0, \infty[$, \mathbf{P}^m -a.s.; because X is a diffusion, we then have the automatic left continuity of $t \mapsto \tilde{u}(X_t)$ on $]0, \zeta[$, \mathbf{P}^m -a.s. The function \tilde{u} is *quasi-continuous* in the sense that there is a nest (K_n) of compact subsets of E such that $\tilde{u}|_{K_n} \in C(K_n)$ for every n .

A Borel set $N \subset E$ is said to be *exceptional* provided $\mathbf{P}^m[\tau(E \setminus N) < \zeta] = 0$. It can be shown that $N \in \mathcal{B}(E)$ is exceptional if and only if there is a nest of compacts $\{K_n\}$ such that $N \subset \bigcap_n K_n^c$. A statement or property depending on $x \in E$ is said to hold quasi-everywhere (q.e.) provided the set of points where it fails to hold is exceptional.

Given $u \in \mathcal{D}$, we have Fukushima's decomposition [13, Theorem 5.2.2]:

$$(1.11) \quad \tilde{u}(X_t) - \tilde{u}(X_0) = M_t^u + N_t^u, \quad \text{for all } t \geq 0, \mathbf{P}^x\text{-a.s. for q.e. } x \in E,$$

where M^u and N^u are continuous additive functionals (CAFs) of X , M^u is a martingale such that $\sup_{t > 0} t^{-1} \mathbf{P}^m[[M_t^u]^2] < \infty$, and $\lim_{t \rightarrow 0} t^{-1} \mathbf{P}^m[[N_t^u]^2] = 0$. This decomposition is unique.

For $u \in \mathcal{D}$ we have $\mathcal{E}(u) = 0$ if and only if $\tilde{u}(X_t) = \tilde{u}(X_0)$ for all $t > 0$, \mathbf{P}^m -a.s. In particular, if $\mathcal{E}(u) = 0$ then $K \circ u \in \mathcal{D}$ for every Borel measurable function $K : \mathbf{R} \rightarrow \mathbf{R}$ such

that $K \circ u \in L^2(m)$, as follows immediately from (1.9). For this reason we shall impose the following (very mild) non-degeneracy condition, without which no conclusions can be drawn about functions that operate on \mathcal{D} :

(1.12) There exists $u \in \mathcal{D}$ such that $\mathcal{E}(u) > 0$.

If (1.12) were to fail then m -a.e. $x \in E$ would be a *trap*, in the sense that $\mathbf{P}^x[X_t = x \text{ for all } t \geq 0] = 1$.

If $u \in \mathcal{D}$ then the martingale CAF M^u admits a quadratic variation process $\langle M^u \rangle$; i.e., $\langle M^u \rangle$ is a PCAF such that $(M^u)^2 - \langle M^u \rangle$ is a martingale on $[0, \zeta[$. The Revuz measure of $\langle M^u \rangle$ (the so-called energy measure of u) is the smooth measure $\mu_{\langle u \rangle}$ on E determined by

$$(1.13) \quad \mu_{\langle u \rangle}(f) = \uparrow \lim_{t \rightarrow 0} t^{-1} \mathbf{P}^m \left[\int_0^t f(X_s) d\langle M^u \rangle_s \right], \quad f \in p\mathcal{B}(E).$$

The measure $\mu_{\langle u \rangle}$ has finite total mass; indeed,

$$\mathcal{E}(u, u) = \frac{1}{2} \mu_{\langle u \rangle}(E), \quad u \in \mathcal{D}.$$

Given two elements u and v of \mathcal{D} , the quadratic covariation $\langle M^u, M^v \rangle$ is a (signed) CAF of X and, since $\langle M^u, M^v \rangle = (1/4)[\langle M^{u+v} \rangle - \langle M^{u-v} \rangle]$, we have

$$(1.14) \quad \mu_{\langle u, v \rangle} = \frac{1}{4} [\mu_{\langle u+v \rangle} - \mu_{\langle u-v \rangle}]$$

and

$$(1.15) \quad \mathcal{E}(u, v) = \frac{1}{2} \mu_{\langle u, v \rangle}(E), \quad u, v \in \mathcal{D}.$$

We close this section with two results that will be useful in the constructions of the next section.

PROPOSITION 1.16. *Let $F : \mathbf{R} \rightarrow \mathbf{R}$ be an absolutely continuous function with $F(0) = 0$. Given $u \in \mathcal{D}$, if $F \circ u \in L^2(m)$ and $F' \circ \tilde{u} \in L^2(\mu_{\langle u \rangle})$, then $F \circ u \in \mathcal{D}$ and*

$$(1.17) \quad d\mu_{\langle F \circ u \rangle} = [F' \circ \tilde{u}]^2 d\mu_{\langle u \rangle}.$$

PROOF. Let ξ denote the image of $(1/2)\mu_{\langle u \rangle}$ under the mapping $x \mapsto \tilde{u}(x)$. This is a finite measure on the Borel subsets of \mathbf{R} . Moreover, ξ is absolutely continuous with respect to Lebesgue measure on \mathbf{R} by [4, Theorem I.7.1.1]. Thus the derivative $F' \in L^1_{\text{loc}}(\mathbf{R})$ is uniquely determined up to ξ -null sets, and the hypothesis $F' \circ \tilde{u} \in L^2(\mu_{\langle u \rangle})$ (equivalently $F' \in L^2(\xi)$) is meaningful. Let η denote the standard Gaussian measure on \mathbf{R} . Then $\xi + \eta$ is a regular Borel measure on \mathbf{R} , hence there is a sequence $\{\varphi_n\}$ in $C_0(\mathbf{R})$ such that $\varphi_n \rightarrow F'$ in $L^2(\xi + \eta)$; see [12, Proposition 7.9]. Define $F_n(t) := \int_0^t \varphi_n(s) ds$ for $t \in \mathbf{R}$. Evidently, F_n is globally Lipschitz continuous, so that $F_n \circ u \in \mathcal{D}$ by the fundamental contraction property of Dirichlet spaces. Moreover, because $F_n \in C^1(\mathbf{R})$ we can apply [13, Theorem 3.2.2] to deduce that

$$(1.18) \quad d\mu_{\langle F_n \circ u \rangle} = [\varphi_n \circ \tilde{u}]^2 d\mu_{\langle u \rangle}$$

for all $n \in N$. Consequently,

$$(1.19) \quad \mathcal{E}(F_m \circ u - F_n \circ u) = \frac{1}{2} \int_E [\varphi_m \circ \tilde{u} - \varphi_n \circ \tilde{u}]^2 d\mu_{(u)} = \int_{\mathbf{R}} [\varphi_m - \varphi_n]^2 d\xi.$$

It follows from (1.19) and the choice of $\{\varphi_n\}$ that the sequence $\{F_n \circ u\}$ is \mathcal{E} -Cauchy in \mathcal{D} . Also,

$$\begin{aligned} |F_n(t) - F(t)| &\leq \int_{-|t|}^{|t|} |\varphi_n(s) - F'(s)| ds \\ &\leq \sqrt{2\pi} e^{t^2/2} \int_{-|t|}^{|t|} |\varphi_n(s) - F'(s)| (2\pi)^{-1/2} e^{-s^2/2} ds \\ &\leq \sqrt{2\pi} e^{t^2/2} \|\varphi_n - F'\|_{L^2(\eta)}, \end{aligned}$$

from which it follows that $\lim_n F_n(t) = F(t)$ for each $t \in \mathbf{R}$. In particular, $F_n \circ u$ converges pointwise to $F \circ u$ as $n \rightarrow \infty$. Thus, $F \circ u$ is an element of the extended Dirichlet space \mathcal{D}_e , see [13, page 35]. Since $F \circ u \in L^2(m)$ by hypothesis, we even have $F \circ u \in \mathcal{D}$ by [13, Theorem 1.5.2 (iii)].

Because $\mathcal{E}(F_n \circ u - F \circ u) \rightarrow 0$ as $n \rightarrow \infty$, the left side of (1.18) converges in total variation norm to $d\mu_{(F \circ u)}$, see [13, 5.2.12] and [9, Lemma 5.6]. On the other hand, the right side of (1.18) converges (setwise) to $[F' \circ \tilde{u}]^2 d\mu_{(u)}$, because $\varphi_n \circ \tilde{u} \rightarrow F' \circ \tilde{u}$ in $L^2(\mu_{(u)})$. This proves (1.17). \square

LEMMA 1.19. *There is a non-negative $w \in \mathcal{D} \cap C_0(E)$ and constants $0 < \alpha < \beta$ such that*

$$(1.20) \quad \mathbf{P}^m[w(X_0) \leq \alpha, T_{\{w>\beta\}} < \zeta] > 0.$$

PROOF. If not, then by varying α and β in (1.20) we see that

$$(1.21) \quad \mathbf{P}^m[w(X_0) < \sup_{s>0} w(X_s)] = 0,$$

for all non-negative $w \in \mathcal{D} \cap C_0(E)$. An application of the simple Markov property at each positive rational time leads from (1.21) to

$$w(X_t) \geq \sup_{s>t} w(X_s), \quad \text{for all rational } t > 0, \mathbf{P}^m\text{-a.e.},$$

and the continuity of w now yields

$$(1.22) \quad t \mapsto w(X_t) \text{ is non-increasing, } \mathbf{P}^m\text{-a.e.},$$

for every non-negative $w \in \mathcal{D} \cap C_0(E)$. From (1.22) and Fukushima's decomposition we deduce that the martingale M^w has paths of zero quadratic variation. That is, $\langle M^w \rangle \equiv 0$, \mathbf{P}^m -a.e., from which it follows that $\mathcal{E}(w) = 0$. The \mathcal{E}_1 -density of $\mathcal{D} \cap C_0(E)$ in \mathcal{D} now implies that $\mathcal{E}(u) = 0$, first for all non-negative $u \in \mathcal{D}$ and then for all $u \in \mathcal{D}$ as we see by expressing $u \in \mathcal{D}$ as the difference of its positive and negative parts. This contradicts the non-degeneracy hypothesis (1.12), proving the Lemma. \square

2. Theorem and proof. This section is devoted to the proof of the following theorem, our main result. The setting is as detailed in section 1; in particular the hypothesis (1.12) is in force.

By *essentially unbounded* we shall mean that $m(x \in E; |u(x)| > t) > 0$ for each $t > 0$. If \mathcal{D} contains no essentially unbounded elements, then $\mathcal{D} \subset L^\infty(m)$, in which case it is easy to see that every locally Lipschitz function operates on \mathcal{D} . A Borel set $D \subset E$ is said to be an *exit set* provided $\sup_{x \in E} \mathbf{P}^x[\tau(D)] < \infty$.

THEOREM 2.1. *Let $K : \mathbf{R} \rightarrow \mathbf{R}$ be Borel measurable, with $K(0) = 0$.*

(a) *If*

$$(2.2) \quad K \circ u \in \mathcal{D} \quad \text{for all } u \in \mathcal{D} \cap L^\infty(m),$$

then K is locally Lipschitz continuous, in the sense that for each $N \in \mathbf{N}$ there exists $M_N \in]0, \infty[$ such that

$$(2.3) \quad |K(t) - K(s)| \leq M_N |t - s| \quad \text{for all } s, t \in [-N, N].$$

Equivalently, K is absolutely continuous, and the a.e. defined derivative K' is an element of $L^\infty_{\text{loc}}(\mathbf{R})$.

(b) *If*

$$(2.4) \quad K \circ u \in \mathcal{D} \quad \text{for all } u \in \mathcal{D}$$

and if \mathcal{D} contains an essentially unbounded element vanishing outside an exit set, then K is globally Lipschitz continuous in the sense that there exists $M \in]0, \infty[$ such that

$$(2.5) \quad |K(t) - K(s)| \leq M |t - s| \quad \text{for all } s, t \in \mathbf{R}.$$

Equivalently, K is absolutely continuous and the a.e. defined derivative K' is an element of $L^\infty(\mathbf{R})$.

REMARK 2.6. By an argument found on pp. 503–504 of [11], it can be shown that if there is a non-empty exceptional set (for example, if X is Brownian motion in a Euclidean domain of dimension $d \geq 2$), then there is an essentially unbounded element of \mathcal{D} vanishing outside an exit set. If there are no non-empty exceptional sets, then the transition measures of X are absolutely continuous with respect to m and each point of E is regular for itself. The author does not know whether, in this latter case, the existence of an unbounded element of \mathcal{D} guarantees the existence of such an element vanishing outside an exit set. An illustrative example of a process with no non-empty exceptional sets but with unbounded elements in its Dirichlet space is the 3-dimensional Bessel process on $]0, \infty[$.

Condenser potentials play a key role in the proof of Theorem (2.1), and in preparation we discuss some of their basic properties. Recall from [13, page 68] that a Borel set $B \subset E$ is *quasi closed* if (and only if) there is a nest $\{F_n\}$ of closed subsets of E such that $B \cap F_n$ is a closed set for each $n \in \mathbf{N}$. For example, if $u \in \mathcal{D}$ and $\alpha \in \mathbf{R}$, then $\{x \in E; \tilde{u}(x) \leq \alpha\}$ is quasi closed. Let A and B be quasi closed Borel subsets of E such that $A \cap B$ is exceptional.

The *condenser potential*

$$p(x) = p_{A,B}(x) := \mathbf{P}^x[T_A < T_B]$$

is an excessive function of X killed at time T_B ; consequently p is quasi-continuous. Notice that $p(x) = 1$ for q.e. $x \in A$ and $p(x) = 0$ for q.e. $x \in B$. If the convex set

$$\mathcal{U} = \mathcal{U}_{A,B} := \{u \in \mathcal{D}; \tilde{u} = 1 \text{ q.e. on } A, \tilde{u} = 0 \text{ q.e. on } B\}$$

is nonempty, then

$$\mathcal{E}(p) = \min\{\mathcal{E}(u); u \in \mathcal{U}\}.$$

Unless the process X killed at T_{B^c} is transient, there may be other elements of \mathcal{U} at which the above minimum is attained. Evidently, \mathcal{U} is non-empty if and only if $p \in \mathcal{D}$. See [8] and [20, §2] for more details on these matters.

If $p = p_{A,B} \in \mathcal{D}$, then there is a finite signed smooth measure $\nu = \nu_{A,B}$ such that

$$(2.7) \quad \mathcal{E}(u, p) = \int_E \tilde{u} d\nu, \quad u \in \mathcal{D}.$$

Indeed, the Hahn-Jordan decomposition $\nu^+ - \nu^-$ of ν is such that both ν^+ and ν^- are smooth measures, and ν^+ is carried by A while ν^- is carried by B . (See the discussion on pp. 75–76 of [6], and (2.15) below). Notice that

$$(2.8) \quad C(A, B) := \mathcal{E}(p, p) = \int_E p d\nu = \nu(A) = \nu^+(E).$$

The following instance of the Bouleau-Hirsch “energy image density” theorem [4, Theorem I.7.1.1] is our main tool. This formula was suggested by a calculation in [14]. In the context of Brownian motion, a different proof can be fashioned out of the co-area and divergence formulas, as was shown to the author by Bruce Driver.

PROPOSITION 2.9. *Fix A and B as above, and assume that $p = p_{A,B}$ lies in \mathcal{D} . Then for every Borel function $g : [0, 1] \rightarrow [0, \infty]$,*

$$(2.10) \quad \frac{1}{2} \int_E g(p(x)) d\mu_{\langle p \rangle}(dx) = C(A, B) \int_0^1 g(t) dt.$$

PROOF. It suffices to prove (2.10) when g is continuous and real-valued. Define $G(y) := \int_0^y g(t) dt$. Then $G \circ p \in \mathcal{D}$ and, by (2.7) and (2.8),

$$\mathcal{E}(G \circ p, p) = \int_A G \circ p d\nu = G(1)\nu(A) = \int_0^1 g(t) dt \cdot C(A, B).$$

On the other hand, by the polarized form of (1.17),

$$\mathcal{E}(G \circ p, p) = \frac{1}{2} \int_E \mu_{\langle G \circ p, p \rangle}(dx) = \frac{1}{2} \int_E g(p(x)) \mu_{\langle p \rangle}(dx). \quad \square$$

The proof of Theorem (2.1) will be accomplished through a series of lemmas. In the remainder of this section we suppose that $K : \mathbf{R} \rightarrow \mathbf{R}$ is a Borel function with $K(0) = 0$, and that (2.2) holds.

LEMMA 2.11. *K is continuous.*

PROOF. Arguing by contradiction, let us suppose that K is not continuous. We may assume that $\liminf_{s \uparrow 1} K(s) < K(1)$; all other cases can be reduced to this case by suitable linear transformations of K . Thus, $\kappa := \liminf_{t \uparrow 1} K(t) < K(1)$, so that there is a strictly increasing sequence $\{t_n\}$ of real numbers with $t_0 = 0$ and $\lim_n t_n = 1$, such that $\lim_n K(t_n) = \kappa$. We can (and do) assume that $\sum_n \sqrt{t_n - t_{n-1}} < \infty$. (This can always be arranged by passing to a subsequence; let $n(k)$ be the smallest positive integer n such that $t_n \geq 1 - 2^{-k}$, then $t_{n(k)} - t_{n(k-1)} \leq 2^{-(k-1)}$.) Define a second increasing sequence of positive numbers $\{\beta_n\}$ by their differences: $\beta_0 := 0$ and

$$\beta_{2n} - \beta_{2n-1} = \beta_{2n+1} - \beta_{2n} = C\sqrt{t_n - t_{n-1}},$$

where the constant C is chosen so that $\lim_n \beta_n = 1$. By Lemma (1.19) there is a non-negative $w \in \mathcal{D} \cap C_0(E)$ such that $\mathbf{P}^m[w(X_0) \leq \alpha, T_{\{w>1\}} < \zeta] > 0$ for some $\alpha \in]0, 1[$. Since $m\{x \in E; w(x) = s\}$ is non-zero for at most countably many values of $s > 0$, we can adjust α and w slightly (by multiplying both by a suitable constant) to ensure that $m\{x \in E; w(x) = 1\} = 0$ as well. Define relatively compact open sets B_1, B_2, \dots by $B_n := \{w > \beta_n\}$. Let $p_n \in \mathcal{D}$ be the condenser potential defined by $p_n(x) := \mathbf{P}^x[T_{\overline{B_{2n}}} < T_{B_{2n-1}^c}]$, $n = 1, 2, \dots$. Then $p_n = 1$ q.e. on $\overline{B_{2n}}$ and $p_n = 0$ q.e. on B_{2n-1}^c . Define $u(x) := \sum_{n=1}^\infty (t_n - t_{n-1})p_n(x)$. Each p_n is quasi-continuous, and the series defining u converges uniformly, and hence u is quasi-continuous. Now p_n is an element of $\mathcal{U}_{\overline{B_{2n}}, B_{2n-1}^c} := \{v \in \mathcal{D}; \tilde{v} = 1 \text{ q.e. on } \overline{B_{2n}}, \tilde{v} = 0 \text{ q.e. on } B_{2n-1}^c\}$ of least energy; thus, if we set

$$w_n := [(\beta_{2n} - \beta_{2n-1})^{-1}(w - \beta_{2n-1})^+] \wedge 1 \in \mathcal{U}_{\overline{B_{2n}}, B_{2n-1}^c},$$

then

$$\mathcal{E}(p_n) \leq \mathcal{E}(w_n) \leq (\beta_{2n} - \beta_{2n-1})^{-2} \mathcal{E}(w).$$

In view of (2.7) and the subsequent discussion, $\mathcal{E}(p_j, p_k) = 0$ if $j \neq k$. Therefore, writing $u_n = \sum_{k=1}^n (t_k - t_{k-1})p_k$, we have

$$\begin{aligned} \mathcal{E}(u_n) &= \sum_{k=1}^n (t_k - t_{k-1})^2 \mathcal{E}(p_k) \\ &\leq \mathcal{E}(w) \sum_{k=1}^n (t_k - t_{k-1})^2 (\beta_{2k} - \beta_{2k-1})^{-2} \\ &= \mathcal{E}(w) \cdot C^{-2} \sum_{k=1}^n (t_k - t_{k-1}) \leq \mathcal{E}(w) \cdot C^{-2}. \end{aligned}$$

In addition, because u_n vanishes q.e. outside B_1 and $0 \leq u_n \leq 1$, we have $\|u_n\|_2^2 \leq m(B_1) \leq \beta_1^{-1} \int_E w dm < \infty$. By the Banach-Saks theorem, the Cesàro means of some subsequence of $\{u_n\}$ converge in \mathcal{D} ; the limit must coincide with the pointwise limit of the $\{u_n\}$, namely u . It follows that $u \in \mathcal{D}$. Notice that $0 \leq u \leq 1$, $u \equiv 1$ on $\{w > 1\}$, and $u = t_n$ q.e. on $\overline{B_{2n}} \setminus B_{2n+1} \supset \{x \in E; \beta_{2n} < w(x) < \beta_{2n+1}\}$.

If $h : \mathbf{R} \rightarrow \mathbf{R}$ is Borel measurable, then

$$\text{ess lim sup}_{s \downarrow t} h(s) := \lim_{\delta \downarrow 0} \text{ess sup}_{s \in]t, t+\delta[} h(s),$$

where “ess sup” denotes essential supremum with respect to Lebesgue measure. The notions $\text{ess lim inf}_{s \downarrow t}$ and $\text{ess lim}_{s \downarrow t}$ are defined analogously, as are their counterparts from the left. If $v \in \mathcal{D}$ then $t \mapsto \tilde{v}(X_t(\omega))$ is continuous on $[0, \zeta(\omega)[$ for \mathbf{P}^m -a.e. $\omega \in \Omega$. Also, $\mathbf{P}^m[\tilde{v}(X_t) \neq v(X_t)] = mP_t(\tilde{v} \neq v) \leq m(\tilde{v} \neq v) = 0$. Consequently, by Fubini’s theorem, $\tilde{v}(X_t(\omega)) = v(X_t(\omega))$ for a.e. $t \geq 0$, for \mathbf{P}^m -a.e. $\omega \in \Omega$. Therefore,

$$(2.12) \quad \tilde{v}(X_t(\omega)) = \text{ess lim}_{s \uparrow t} v(X_s(\omega)) = \text{ess lim}_{s \downarrow t} v(X_s(\omega)), \quad \text{for all } t \in]0, \zeta(\omega)[,$$

for \mathbf{P}^m -a.e. $\omega \in \Omega$. Define $\mathbf{P}^*[\cdot] := \int_{\{w \leq \alpha\}} \mathbf{P}^x[\cdot] m(dx)$, and let τ denote the hitting time of $\{w > 1\}$. Observe that

$$\mathbf{P}^m \left[\int_0^\infty 1_{\{w=1\}}(X_t) dt \right] = \int_0^\infty mP_t(w = 1) dt \leq \int_0^\infty m(w = 1) dt = 0.$$

This implies that

$$(2.13) \quad \mathbf{P}^*[\text{there exists } \delta > 0 \text{ such that } w(X_t) = 1 \text{ for all } t \in]\tau - \delta, \tau[; \tau < \zeta] = 0.$$

Recall that w and α were chosen so that $\mathbf{P}^*[\tau < \zeta] > 0$, and clearly $\mathbf{P}^*[\tau = 0] = 0$. Let $\omega \in \{X_0 \leq \alpha\}$ be a sample point such that $t \mapsto X_t(\omega)$ is continuous and (2.12) holds with $v = K \circ u$. Because of (2.13) we may suppose that $t \mapsto w(X_t(\omega))$ is *not* identically 1 in any interval of the form $]\tau(\omega) - \delta, \tau(\omega)[$. Thus, given $\delta \in]0, \tau(\omega)[$, there exists $t_\delta \in]\tau(\omega) - \delta, \tau(\omega)[$ such that $w(X_{t_\delta}(\omega)) < 1$, which implies that $t \mapsto w(X_t(\omega))$ takes on every value between $w(X_{t_\delta}(\omega))$ and 1 during the time interval $]\tau(\omega) - \delta, \tau(\omega)[$ because $w(X_{\tau(\omega)}(\omega)) = 1$ if $\tau(\omega) < \zeta(\omega)$. Consequently, there is a positive integer $n_0(\omega, \delta)$ such that for each $n \geq n_0(\omega, \delta)$, $X_t(\omega) \in \{\beta_{2n} < w < \beta_{2n+1}\} \subset \overline{B_{2n}} \setminus B_{2n+1}$ for some $t \in]\tau(\omega) - \delta, \tau(\omega)[$. Because $t \mapsto X_t(\omega)$ is continuous and $\{\beta_{2n} < w < \beta_{2n+1}\}$ is open, we conclude that the Lebesgue measure of $\{t \in]\tau(\omega) - \delta, \tau(\omega)[; X_t(\omega) \in \overline{B_{2n}} \setminus B_{2n+1}\}$ is strictly positive for each $n \geq n_0(\omega, \delta)$. Because $u(x) = t_n$ for q.e. $x \in \overline{B_{2n}} \setminus B_{2n+1}$ we therefore have

$$\text{ess inf}_{\tau(\omega) - \delta < t < \tau(\omega)} K(u(X_t(\omega))) \leq \inf_{n \geq n_0(\omega, \delta)} K(t_n).$$

In view of (2.12) (which, in particular, guarantees the existence of $\text{ess lim}_{t \uparrow \tau} K(u(X_t))$), this last display implies that

$$\widetilde{K \circ u}(X_\tau) = \text{ess lim}_{t \uparrow \tau} K(u(X_t)) \leq \lim_n K(t_n) = \kappa < K(1)$$

\mathbf{P}^* -a.s. on the event $\{\tau < \zeta\}$. On the other hand, because $\{w > 1\}$ is open, if $\tau(\omega) < \zeta(\omega)$ then for every $\delta > 0$ the set $\{t \in]\tau(\omega), \tau(\omega) + \delta[; w(X_t(\omega)) > 1\}$ has positive Lebesgue measure. Because $u \equiv 1$ on $\{w > 1\}$ we therefore have

$$\widetilde{K \circ u}(X_\tau) = \text{ess lim}_{t \downarrow \tau} K(u(X_t)) = K(1)$$

\mathbf{P}^* -a.s. on the event $\{\tau < \zeta\}$. Since $\mathbf{P}^*[\tau < \zeta] > 0$, the last two displays provide the desired contradiction. \square

LEMMA 2.14. *K is absolutely continuous, and $K' \in L^2_{\text{loc}}(\mathbf{R})$.*

PROOF. We begin by showing that K is absolutely continuous on $]0, 1[$ with $K' \in L^2_{\text{loc}}(]0, 1[)$. For this it suffices to prove that there is a constant C such that

$$(2.15) \quad \left| \int_0^1 K(s)\psi'(s)ds \right| \leq C\|\psi\|_2, \quad \text{for all } \psi \in C_0^\infty(]0, 1[).$$

Fix $\psi \in C_0^\infty(]0, 1[)$ and define $\Psi(t) := \int_0^t \psi(s)ds$. Let U and V be non-empty open relatively compact subsets of E such that $\bar{U} \subset V$. Let p be the condenser potential associated with $A = \bar{U}$ and $B = V^c$. From [6, (3.4)] we know that there are PCAFs C^A and C^B with C^A (resp. C^B) increasing only when X is in A (resp. in B) such that the Fukushima decomposition associated with p is

$$(2.16) \quad p(X_t) - p(X_0) = M_t^p + N_t^p = M_t^p - C_t^A + C_t^B, \quad 0 \leq t < \zeta.$$

Indeed, C^A (resp. C^B) is the PCAF with Revuz measure [13, Section 5.1] ν^+ (resp. ν^-), where $\nu^+ - \nu^-$ is the Hahn-Jordan decomposition of the signed smooth measure associated with p . Now $\Psi \circ p \in \mathcal{D}$, and from (2.16) and Itô's formula it follows that

$$(2.17) \quad N_t^{\Psi \circ p} = \frac{1}{2} \int_0^t \psi'(p(X_s))d\langle M^p \rangle_s, \quad t \geq 0.$$

By a result of Nakao [19, §3], (2.17) yields

$$(2.18) \quad \mathcal{E}(v, \Psi \circ p) = -\frac{1}{2} \int_E \tilde{v}(x)\psi'(p(x))\mu_{(p)}(dx),$$

for any bounded $v \in \mathcal{D}$. In particular, by (2.10),

$$\mathcal{E}(K \circ p, \Psi \circ p) = -\frac{1}{2} \int_E K(p(x))\psi'(p(x))\mu_{(p)}(dx) = -C(A, B) \int_0^1 K(s)\psi'(s)ds.$$

On the other hand,

$$\begin{aligned} |\mathcal{E}(K \circ p, \Psi \circ p)| &\leq \mathcal{E}(K \circ p)^{1/2}\mathcal{E}(\Psi \circ p)^{1/2} \\ &= \mathcal{E}(K \circ p)^{1/2}C(A, B)^{1/2}\|\psi\|_2, \end{aligned}$$

where we have used (1.15) and (2.10) for the equality. Thus (2.15) holds with $C = [\mathcal{E}(K \circ p)/C(A, B)]^{1/2}$.

Now notice that $J(x) := K(|4x - 1| - 1)$ satisfies (2.2). By the preceding paragraph, J is absolutely continuous on $]0, 1[$ and $J' \in L^2_{\text{loc}}(]0, 1[)$. But the restriction of J to $[1/4, 3/4]$ is $x \mapsto K(4x - 2)$. It follows that K is absolutely continuous on $[-1, 1]$ and that $K' \in L^2[-1, 1]$. Finally, for $b > 0$, the preceding applies to $K_b(x) := K(bx)$, the upshot being that K is absolutely continuous on $[-b, b]$ with $K' \in L^2[-b, b]$ \square

LEMMA 2.19. $K' \in L^\infty_{\text{loc}}(\mathbf{R})$.

PROOF. Arguing as at the end of the proof of (2.14), it is enough to show that $K' \in L^\infty[0, 1]$. For this we need only to show that if $0 \leq f \in L^1[0, 1]$, then

$$\int_0^1 [K'(s)]^2 f(s) ds < \infty.$$

Fix such an f and define $\varphi : [0, 1] \rightarrow [0, 1]$ by the formula

$$t = \gamma^{-1} \int_0^{\varphi(t)} \frac{1}{1+f(s)} ds, \quad 0 \leq t \leq 1,$$

where $\gamma := \int_0^1 [1+f(s)]^{-1} ds$. Evidently, φ is a strictly increasing map of $[0, 1]$ onto itself, and φ is an absolutely continuous function with $\varphi(0) = 0$ and

$$(2.20) \quad \varphi'(t) = \gamma[1+f(\varphi(t))]$$

for Lebesgue a.e. $t \in [0, 1]$. Let the condenser potential p be as in the proof of Lemma (2.14). Observe that $\varphi \circ p$ is bounded and $m(\varphi \circ p \neq 0) \leq m(p > 0) < \infty$; therefore $\varphi \circ p \in L^2(m)$. By Proposition (1.16), $\varphi \circ p \in \mathcal{D}$ because

$$\frac{1}{2} \int_E [\varphi' \circ p]^2 d\mu_{(p)} = C(A, B) \int_0^1 [\varphi'(s)]^2 ds = \gamma C(A, B) \int_0^1 [1+f(t)] dt < \infty.$$

Therefore $K \circ \varphi \circ p \in \mathcal{D}$, and hence

$$\begin{aligned} \int_0^1 [K'(t)]^2 [1+f(t)] dt &= \frac{1}{\gamma} \int_0^1 [K'(\varphi(s))]^2 [\varphi'(s)]^2 ds \\ &= \frac{1}{2\gamma C(A, B)} \int_E [K'(\varphi(p(x)))]^2 [\varphi'(p(x))]^2 \mu_{(p)}(dx) \\ &= \frac{1}{\gamma C(A, B)} \mathcal{E}(K \circ \varphi \circ p) < \infty. \quad \square \end{aligned}$$

LEMMA 2.21. *If \mathcal{D} contains an essentially unbounded element vanishing outside an exit set, then $K' \in L^\infty(\mathbf{R})$.*

PROOF. It suffices to show that $K' \in L^\infty[0, \infty[$, and for this it is enough to show that

$$\int_0^\infty [K'(s)]^2 f(s) ds < \infty$$

for each non-negative $f \in L^1[0, \infty[$. Fix such an f . Let v be an essentially unbounded element of \mathcal{D} such that $\{\tilde{v} \neq 0\}$ is an exit set; substituting $|v|$ for v if necessary, we can and do assume that $v \geq 0$. For $n \in \mathbf{N}$ define quasi closed sets $A_n := \{\tilde{v} \geq 2n + 1\}$ and $B_n := \{\tilde{v} \leq 2n\}$, and let p_n be the associated condenser potential p_{A_n, B_n} . Because v is essentially unbounded, the condenser capacities

$$C_n := C(A_n, B_n)$$

are strictly positive and finite. Notice that

$$\mathcal{E}(p_n, p_m) = 0 \quad \text{if } m \neq n.$$

Because

$$C_n \leq \mathcal{E}((v - 2n)^+ \wedge 1) = \frac{1}{2} \int_E 1_{[2n, 2n+1]}(\tilde{v}(x)) \mu_{(v)}(dx) \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

we have $\sum_n C_n^{-1} = \infty$. Next observe that

$$x^2 \leq \int_0^x [e^{-s} + f(s)] ds \cdot \int_0^x [e^{-s} + f(s)]^{-1} ds,$$

and hence the integrability of $s \mapsto e^{-s} + f(s)$ implies that

$$\lim_{x \rightarrow \infty} \int_0^x [e^{-s} + f(s)]^{-1} ds = \infty.$$

Therefore a strictly increasing function $\varphi : [0, \infty[\rightarrow [0, \infty[$ is well-defined by

$$t = \int_0^{\varphi(t)} \frac{1}{e^{-s} + f(s)} ds, \quad t \geq 0.$$

Clearly, $\varphi'(s) = e^{-\varphi(s)} + f(\varphi(s))$ for Lebesgue a.e. $s > 0$, $\varphi(0) = 0$, and $\lim_{s \rightarrow \infty} \varphi(s) = \infty$.

Let us now check that the function u defined by

$$u(x) := \varphi \left(\sum_{k=1}^{\infty} C_k^{-1} p_k(x) \right), \quad x \in E,$$

is an element of \mathcal{D} . Proposition (1.16) implies that for each $n \in \mathbb{N}$,

$$u_n(x) := \varphi \left(\sum_{k=1}^n C_k^{-1} p_k(x) \right), \quad x \in E,$$

is an element of \mathcal{D} . Clearly, u is the pointwise limit of the sequence $\{u_n\}$. An application of the Banach-Saks theorem (as in the proof of Lemma (2.11)) shows that the asserted inclusion of $u \in \mathcal{D}$ will follow once we check that both $\mathcal{E}(u_n)$ and $\|u_n\|_2$ are bounded functions of n . Now, because p_k is 1 on the support of p_j for $j > k$, we have $\mu_{(p_j, p_k)} = 0$ for $j > k$. Thus, defining $x_0 := 0$ and $x_k := C_1^{-1} + \dots + C_k^{-1}$ for $k \geq 1$, the key formula (2.10) yields that

$$\begin{aligned}
 \mathcal{E}(u_n) &= \sum_{k=1}^n \int_E [\varphi'(x_{k-1} + C_k^{-1} p_k(x))]^2 C_k^{-2} \mu_{\langle p_k \rangle}(dx) \\
 &= \sum_{k=1}^n C_k \int_0^1 [\varphi'(x_{k-1} + C_k^{-1} s)]^2 C_k^{-2} ds \\
 &= \sum_{k=1}^n \int_0^1 [\varphi'(x_{k-1} + C_k^{-1} s)]^2 C_k^{-1} ds \\
 &= \sum_{k=1}^n \int_0^{C_k^{-1}} [\varphi'(x_{k-1} + t)]^2 dt \\
 &= \int_0^{x_n} [\varphi'(z)]^2 dz \leq \int_0^\infty [\varphi'(z)]^2 dz \\
 &= \int_0^\infty \varphi'(\varphi^{-1}(t)) dt \\
 &= \int_0^\infty [e^{-t} + f(t)] dt < \infty.
 \end{aligned}$$

Because u_n vanishes outside the exit set $\{\tilde{v} > 0\}$, we have (by [22, Theorem 3.1] or [10, (1.19)] applied to X killed at the exit time from $\{\tilde{v} > 0\}$)

$$\|u_n\|_2^2 \leq c_1 \mathcal{E}(u_n) \leq c_1 \int_0^\infty [e^{-t} + f(t)] dt < \infty,$$

where $c_1 := \sup_{x \in E} \mathbf{P}^x [T_{\{\tilde{v}=0\}}] < \infty$. Thus, $u \in \mathcal{D}$, hence $K \circ u \in \mathcal{D}$. Computing as above, we find, because $\lim_{n \rightarrow \infty} x_n = \infty$,

$$\infty > \mathcal{E}(K \circ u) = \int_0^\infty [K'(t)]^2 [e^{-t} + f(t)] dt \geq \int_0^\infty [K'(t)]^2 f(t) dt,$$

as desired. □

Acknowledgment. I thank the referee, whose careful reading of the manuscript led to the elimination of several errors.

REFERENCES

[1] A. ANCONA, Continuité des contractions dans les espaces de Dirichlet, Séminaire de Théorie du Potentiel de Paris, no. 2, pp. 1–26, Lecture Notes in Math. 563, Springer, Berlin, 1976. [MR 58 #28567]
 [2] J. APPELL AND P. P. ZABREJKO, Nonlinear Superposition Operators, Cambridge University Press, Cambridge, 1990. [MR 91k:47168]
 [3] R. M. BLUMENTHAL AND R. K. GETTOOR, Markov Processes and Potential Theory, Academic Press, New York-London, 1968. [MR 41 #9348]
 [4] N. BOULEAU AND F. HIRSCH, Formes de Dirichlet générales et densité des variables aléatoires réelles sur l'espace de Wiener, J. Funct. Anal. 69 (1986), 229–259. [MR 88d:60196]
 [5] G. BOURDAUD, Le calcul fonctionnel dans les espaces de Sobolev, Invent. Math. 104 (1991), 435–446. [MR 93b:46053]

- [6] Z.-Q. CHEN, P. J. FITZSIMMONS AND R. SONG, Crossing estimates for symmetric Markov processes, *Probab. Theory Related Fields* 120 (2001), 68–84.
- [7] Z.-Q. CHEN, Z.-M. MA AND M. RÖCKNER, Quasi-homeomorphisms of Dirichlet forms, *Nagoya Math. J.* 136 (1994), 1–15. [MR 95m:31020]
- [8] K. L. CHUNG AND R. K. GETTOOR, The condenser problem, *Ann. Probab.* 5 (1977), 82–86. [MR 55 #1501]
- [9] P. J. FITZSIMMONS, Absolute continuity of symmetric diffusions, *Ann. Probab.* 25 (1997), 230–258. [MR 98c:60097]
- [10] P. J. FITZSIMMONS, Hardy’s inequality for Dirichlet forms, *J. Math. Anal. Appl.* 250 (2000), 548–560. [MR 2001h:31010]
- [11] P. J. FITZSIMMONS AND R. K. GETTOOR, On the potential theory of symmetric Markov processes, *Math. Ann.* 281 (1988), 495–512. [MR 89k:60110]
- [12] G. B. FOLLAND, *Real Analysis* (Second edition), Wiley, New York, 1999. [MR 2000c:00001]
- [13] M. FUKUSHIMA, Y. ŌSHIMA AND M. TAKEDA, *Dirichlet Forms and Symmetric Markov Processes*, Walter de Gruyter & Co., Berlin, 1994. [MR 96f:60126]
- [14] M. FUKUSHIMA AND J. YING, A note on regular Dirichlet subspaces, *Proc. Amer. Math. Soc.* 131 (2003), 1607–1610.
- [15] D. A. LABUTIN, Superposition operator in Sobolev spaces on domains, *Proc. Amer. Math. Soc.* 128 (2000), 3399–3403. [MR 2001b:47106]
- [16] Z.-M. MA AND M. RÖCKNER, *Introduction to the Theory of (Nonsymmetric) Dirichlet Forms*, Springer-Verlag, Berlin, 1992. [MR 94d:60119]
- [17] M. MARCUS AND V. J. MIZEL, Complete characterization of functions which act, via superposition, on Sobolev spaces, *Trans. Amer. Math. Soc.* 251 (1979), 187–218. [MR 80j:46055]
- [18] M. MARCUS AND V. J. MIZEL, Every superposition operator mapping one Sobolev space into another is continuous, *J. Funct. Anal.* 33 (1979), 217–229. [MR 80h:47039]
- [19] S. NAKAO, Stochastic calculus for continuous additive functionals of zero energy, *Z. Wahrsch. Verw. Gebiete* 68 (1985), 557–578. [MR 86c:60082]
- [20] Y. ŌSHIMA AND T. YAMADA, On some representations of continuous additive functionals locally of zero energy, *J. Math. Soc. Japan* 36 (1984), 315–339. [MR 85j:60136]
- [21] M. SHARPE, *General Theory of Markov Processes*, Academic Press, San Diego, 1988. [MR 89m:60169]
- [22] P. STOLLMANN AND J. VOIGT, Perturbation of Dirichlet forms by measures, *Potential Anal.* 5 (1996), 109–138. [MR 97e:47065]

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF CALIFORNIA, SAN DIEGO
9500 GILMAN DRIVE
LA JOLLA, CA 92093–0112
U.S.A.

E-mail address: pfitz@euclid.ucsd.edu