

## ON RINGS WITH FINITE SELF-INJECTIVE DIMENSION II

(Dedicated to Professor Goro Azumaya on his 60th birthday)

By

Yasuo IWANAGA

For a module  $M$  over a ring  $R$  (with an identity),  $\text{pd}(M)$  and  $\text{id}(M)$  denote the projective and injective dimension of  $M$ , respectively. In the previous paper [5] and [6], we called a (left and right) noether ring  $R$   $n$ -Gorenstein if  $\text{id}({}_R R) \leq n$  and  $\text{id}(R_R) \leq n$  for an  $n \geq 0$ , and Gorenstein if  $R$  is  $n$ -Gorenstein for some  $n$ . This note is concerned with two subjects on Gorenstein rings. In §1, we consider the modules of finite projective or injective dimension over a Gorenstein ring and, first, show that the finiteness of projective dimension coincides with one of injective dimension. Then it follows that the highest finite projective (or injective) dimension is  $n$  for modules over an  $n$ -Gorenstein ring and, next, such modules over an artinian Gorenstein ring are investigated. Finally, we present some example to compare with Auslander's definition of an  $n$ -Gorenstein ring.

In §2, for a Gorenstein ring  $R$ , we consider a quasi-Frobenius extension of  $R$  and show it also is a Gorenstein ring. Further we generalize [3, Corollary 8 and 8'] to the case of a quasi-Frobenius extension. Also an example concerning with a maximal quotient ring of a Gorenstein ring is presented.

### 1. Modules of finite projective or injective dimension

We start with the next proposition which states [4, Korollar 1.12] and [7, Corollary 5] more precisely:

PROPOSITION 1. For a noether ring  $R$ ,

$$\text{id}(R_R) = \sup \{ \text{flat dim}(E); {}_R E \text{ is an injective left } R\text{-module.} \}.$$

PROOF. By [2, Chap. VI, Proposition 5.3],

$$(*) \quad \text{Tor}_i^R(A_R, {}_R E) \cong \text{Hom}_R(\text{Ext}_R^i(A_R, {}_R R), {}_R E)$$

for any finitely generated right  $R$ -module  $A_R$ , injective left  $R$ -module  ${}_R E$  and  $i > 0$ .

First assume  $\text{id}(R_R) = n < \infty$ , then  $\text{Ext}_R^{n+1}(A, R) = 0$  for any finitely generated

$A_R$  and so  $\text{Tor}_{n+1}^R(A, E)=0$  for any injective  ${}_R E$ . Further, for any  $X_R$ , we can represent  $X=\varinjlim A_\alpha$  such that each  $A_\alpha$  is finitely generated and hence

$$\text{Tor}_{n+1}^R(X_R, {}_R E) \cong \varinjlim \text{Tor}_{n+1}^R(A_\alpha, E) = 0.$$

Therefore  $\text{flat dim}(E) \leq n$ .

Conversely, if  $\text{flat dim}(E) \leq n < \infty$  for any injective  ${}_R E$ , (\*) induces

$$\text{Hom}_R(\text{Ext}_R^{n+1}(A, R), E) \cong \text{Tor}_{n+1}^R(A, E) = 0$$

for any finitely generated  $A_R$ . Now then, by taking  ${}_R E$  as an injective cogenerator, it holds that  $\text{Ext}_R^{n+1}(A, R)=0$  for any finitely generated  $A_R$  and hence  $\text{id}(R_R) \leq n$ .

The following was shown in [5] and [6] under certain assumption on the dominant dimension, but now we can release this assumption and include completely the commutative case.

**THEOREM 2.** *For an  $n$ -Gorenstein ring  $R$  and an  $R$ -module  $M$ , the following are equivalent:*

$$(1) \text{pd}(M) < \infty, \quad (2) \text{pd}(M) \leq n, \quad (3) \text{id}(M) < \infty, \quad (4) \text{id}(M) \leq n.$$

**PROOF.** Since the implications (1) $\Rightarrow$ (2) and (2) $\Rightarrow$ (4) are proved in [1] and [5], respectively, we prove only (3) $\Rightarrow$ (2).

Let

$$0 \longrightarrow M \xrightarrow{f_0} E_0 \xrightarrow{f_1} E_1 \longrightarrow \cdots \xrightarrow{f_m} E_m \longrightarrow 0$$

be an injective resolution of  $M$  and  $K_{i-1} = \ker(f_i)$  ( $i=1, \dots, m$ ), then in the exact sequence

$$0 \longrightarrow K_{m-1} \longrightarrow E_{m-1} \longrightarrow E_m \longrightarrow 0,$$

if  $\text{pd}(E_{m-1}), \text{pd}(E_m) \leq n$ , then  $\text{pd}(K_{m-1}) \leq n$  by [5, Lemma 4]. For an arbitrary  $i$ , in the exact sequence

$$0 \longrightarrow K_{i-1} \longrightarrow E_{i-1} \longrightarrow K_i \longrightarrow 0,$$

if  $\text{pd}(K_i), \text{pd}(E_{i-1}) \leq n$ , then  $\text{pd}(K_{i-1}) \leq n$  and therefore  $\text{pd}(M) = \text{pd}(K_0) \leq n$  by the induction. Thus, it is enough to show  $\text{pd}(E) \leq n$  for any injective left module  ${}_R E$ .

Now, since  $\text{flat dim}(E) \leq n$  by Proposition 1, let

$$0 \longrightarrow U_n \xrightarrow{f_n} U_{n-1} \xrightarrow{f_{n-1}} \cdots \xrightarrow{f_1} U_0 \longrightarrow E \longrightarrow 0$$

be a resolution of  ${}_R E$  by flat modules  $U_i$  ( $i=0, 1, \dots, n$ ) and  $C_{i-1} = \text{cok}(f_i)$  ( $i=1, \dots, n$ ), then  $\text{pd}(U_i) < \infty$  for  $i=0, 1, \dots, n$  by [7, Proposition 6]. First,

from the exact sequence

$$0 \longrightarrow U_n \longrightarrow U_{n-1} \longrightarrow C_{n-1} \longrightarrow 0$$

with  $\text{pd}(U_n), \text{pd}(U_{n-1}) < \infty$ , it follows that  $\text{pd}(C_{n-1}) < \infty$ . For an arbitrary  $i$ , in the exact sequence

$$0 \longrightarrow C_{i+1} \longrightarrow U_i \longrightarrow C_i \longrightarrow 0,$$

if  $\text{pd}(C_{i+1}), \text{pd}(U_i) < \infty$ , then it follows that  $\text{pd}(C_i) < \infty$  and hence  $\text{pd}(E) = \text{pd}(C_0) < \infty$  by the induction, which is equivalent to  $\text{pd}(E) \leq n$  by the implication (1)  $\Rightarrow$  (2).

From Theorem 2, we are interested in modules  $M$  satisfying  $\text{pd}(M) = n$  or  $\text{id}(M) = n$  over an  $n$ -Gorenstein ring. Thus we next consider such modules.

For a module  $M$ , we define  $E^i(M)$  for  $i \geq 0$  as the  $(i+1)$ -th term in a minimal injective resolution of  $M$  and  $E(M) = E^0(M)$ , i. e.

$$0 \longrightarrow M \longrightarrow E^0(M) \longrightarrow \dots \longrightarrow E^i(M) \longrightarrow \dots$$

is a minimal injective resolution of  $M$ . Dually, if  $M$  has a minimal projective resolution, we define  $P^i(M)$  for  $i \geq 0$ , similarly.

**THEOREM 3.** *Let  $R$  be an artinian  $n$ -Gorenstein ring,  $0 \rightarrow {}_R R \rightarrow E_0 \rightarrow \dots \rightarrow E_n \rightarrow 0$  a minimal injective resolution for  ${}_R R$  and  ${}_R M$  a left  $R$ -module.*

(1) *If  $\text{id}(M) = n$ , then  $\text{id}(M) = \text{pd}(E^n(M)) = n$  and, for any direct summand  ${}_R E$  of  $E^n(M)$ ,  $\text{pd}(E) = n$ .*

*If  $\text{pd}(M) = n$ , then  $\text{id}(P^n(M)) = \text{pd}(M) = n$  and, for any direct summand  ${}_R P$  of  $P^n(M)$ ,  $\text{id}(P) = n$ .*

*In particular,  $\text{id}(P^n(E_n)) = \text{pd}(E_n) = n$  provided  $\text{id}({}_R R) = n$ .*

(2) *If  $\text{pd}(M) = n$ , then  $E^n P^n(M)$  is isomorphic to a direct summand of a direct sum of copies of  $E_n$ .*

*Especially,  $E^n P^n(E_n)$  is isomorphic to a direct summand of  $E_n$ .*

**PROOF.** (1) Suppose  $\text{id}(M) = n$  and  ${}_R E$  an indecomposable summand of  $E^n(M)$ , then since  $E$  is of the form  $E(S)$  for some simple module  ${}_R S$ , the exact sequence

$$0 \longrightarrow {}_R S \longrightarrow {}_R E(S) \longrightarrow {}_R E(S)/S \longrightarrow 0$$

induces

$$\text{Ext}_R^n(E(S), M) \longrightarrow \text{Ext}_R^n(S, M) \longrightarrow \text{Ext}_R^{n+1}(E(S)/S, M) \quad (\text{exact}).$$

Here,  $\text{Ext}_R^{n+1}(E(S)/S, M) = 0$  but  $\text{Ext}_R^n(S, M) \neq 0$  by [6, Lemma 1] since  ${}_R S$  is monomorphic to  $E^n(M)$ , and hence  $\text{Ext}_R^n(E(S), M) \neq 0$ . So  $\text{pd}(E(S)) \geq n$  implies  $\text{pd}(E(S)) = n$  by Theorem 2.

Next, assume  $\text{pd}(M) = n$  and  ${}_R P$  an indecomposable summand of  $P^n(M)$ , then

for any simple homomorphic image  ${}_R S$  of  $P$ , the exact sequence

$$0 \longrightarrow {}_R K \longrightarrow {}_R P \longrightarrow {}_R S \longrightarrow 0$$

induces

$$\text{Ext}_R^n(M, P) \longrightarrow \text{Ext}_R^n(M, S) \longrightarrow \text{Ext}_R^{n+1}(M, K) \quad (\text{exact}).$$

Now, since  $\text{Ext}_R^{n+1}(M, K)=0$  but  $\text{Ext}_R^n(M, S)\neq 0$  by the dual of [6, Lemma 1],  $\text{Ext}_R^n(M, P)\neq 0$  and hence  $\text{id}(P)=n$  again by Theorem 2.

(2) Decompose  ${}_R R$  into projective indecomposables, then for any projective indecomposable  ${}_R P$  with  $\text{id}(P)=n$ ,  $E^n(P)$  is isomorphic to a direct summand of  $E_n$ . On the other hand, if  $\text{pd}(M)=n$ ,  $\text{id}(P^n(M))=n$  by (1) and hence  $E^n P^n(M)$  is isomorphic to a summand of a direct sum of copies of  $E_n$ .

**COROLLARY 4.** *Let  $R$  be an  $n$ -Gorenstein ring with  $\text{dom}\cdot\text{dim}_R R \geq n$  and assume  ${}_R M$  a left  $R$ -module with  $\text{id}(M)=n$ , then  $E^n(M)$  is isomorphic to a direct summand of a direct sum of copies of  $E_n$ .*

Now we present an example which seems itself interesting.

**EXAMPLE.** Let  $R$  be an artinian Gorenstein ring with  $\text{id}({}_R R)=n$  and  $0 \rightarrow {}_R R \rightarrow E_0 \rightarrow \cdots \rightarrow E_n \rightarrow 0$  a minimal injective resolution of  ${}_R R$ , then we see from Theorem 3 that  $E_n$  has the largest projective dimension  $n$ . Here, we give an example of  $n$ -Gorenstein ring  $R$  with  $\text{pd}(E_0)=\cdots=\text{pd}(E_n)=n$ , which shows that our definition of an  $n$ -Gorenstein ring is different from Auslander's one.

Let  $k$  be a field and  $R$  a subalgebra of  $(k)_8$ , all  $8 \times 8$  matrices over  $k$ , having  $\{c_{11}+c_{88}, c_{22}+c_{55}, c_{33}+c_{44}, c_{66}, c_{77}, c_{21}, c_{31}, c_{32}, c_{54}, c_{86}, c_{87}\}$  as a  $k$ -basis where  $c_{ij}$  is a matrix unit in  $(k)_8$ . Then  $\text{id}({}_R R)=\text{id}(R_R)=2$ , i. e.  $R$  is 2-Gorenstein,  $\text{gl}\cdot\text{dim } R = \infty$  and  $\text{pd}(E_0)=\text{pd}(E_1)=\text{pd}(E_2)=2$ . Further any left  $R$ -module of projective dimension  $=2$  is a summand in a direct sum of copies of  $E_0 \oplus E_1 \oplus E_2$ .

## 2. A quasi-Frobenius extension of a Gorenstein ring

For rings  $R \subseteq T$ ,  $T/R$  is called a *left quasi-Frobenius (=QF) extension* if  ${}_R T$  is finitely generated projective and  ${}_T T_R$  is isomorphic to a direct summand in a direct sum of copies of  ${}_T \text{Hom}_R({}_R T_T, {}_R R_R)_R$ . A *quasi-Frobenius extension* is a left and right quasi-Frobenius extension. See [9] for details.

In this section we show a QF extension of a Gorenstein ring is also a Gorenstein ring. First we observe the following.

Let  $R, T$  be rings and  $F: {}_R \mathbf{M} \rightarrow {}_T \mathbf{M}$  a functor of the category of left  $R$ -modules to one of left  $T$ -modules, which satisfies the condition:

- 1)  $F$  is exact,

2) if  ${}_R E$  is injective, so is  ${}_T F(E)$ ,

then  $\text{id}({}_T F(M)) \leq \text{id}({}_R M)$  for any left  $R$ -module  ${}_R M$ . Further if

3)  $F$  preserves an essential monomorphism

is satisfied,  $\text{id}({}_T F(M)) = \text{id}({}_R M)$  for any  ${}_R M$ .

The next is a generalization of [3, Corollary 8] to a quasi-Frobenius extension and concerns with the case of a Gorenstein order [10, Lemma 5].

PROPOSITION 5. *Let  $T$  be a left quasi-Frobenius extension of a ring  $R$  and  ${}_R M$  a left  $R$ -module, then*

$$\text{id}({}_T T \otimes_R M) \leq \text{id}({}_R M).$$

PROOF. By [2, VI Proposition 5.2],

$$\begin{aligned} {}_T T \otimes_R M &\cong {}_T T \otimes_R \text{Hom}_R({}_R R_R, {}_R M) \\ &\cong {}_T \text{Hom}_R({}_R \text{Hom}_R({}_T T_R, {}_R R_R)_T, {}_R M). \end{aligned}$$

Here,  $T_R$  is projective by [9, Satz 2] and since  $\text{Hom}_R({}_T T_R, {}_R R_R)_T$  is projective ([9, Satz 2]),  ${}_T T \otimes_R E$  is injective for any injective left  $R$ -module  ${}_R E$ . Therefore the functor  ${}_T T \otimes_R - : {}_R \mathbf{M} \rightarrow {}_T \mathbf{M}$  satisfies the conditions 1)–2) and so

$$\text{id}({}_T T \otimes_R M) \leq \text{id}({}_R M).$$

The following should be compared with [9, Satz 3].

COROLLARY 6. *A quasi-Frobenius extension of an  $n$ -Gorenstein ring also is an  $n$ -Gorenstein ring.*

In connection with [1, Example (2)] and [3, Corollary 8'], we state the following.

PROPOSITION 7. (1) *Let  $T$  be a left quasi-Frobenius extension of a ring  $R$  and suppose  $T_R$  a generator, then*

$$\text{id}({}_T T \otimes_R M) = \text{id}({}_R M)$$

for any left  $R$ -module  $M$  and especially  $\text{id}({}_T T) = \text{id}({}_R R)$ .

Moreover, for a finite group  $G$  and a ring  $R$ ,

$$\text{id}({}_{R[G]} R[G]) = \text{id}({}_R R).$$

(2) *Let  $T$  be a quasi-Frobenius extension of a ring  $R$  and suppose  ${}_R T$  (or  $T_R$ ) a generator, then*

$$\text{id}({}_T T) = \text{id}({}_R R) \quad \text{and} \quad \text{id}(T_T) = \text{id}(R_R).$$

PROOF. (1) Let  $F = T \otimes_R - : {}_R \mathbf{M} \rightarrow {}_T \mathbf{M}$ , then  $F$  satisfies the conditions 1)–3) for  $T_R$  is a progenerator by [9, Satz 2].

(2) Let  $F = \text{Hom}_R({}_R T_T, -): {}_R \mathbf{M} \rightarrow {}_T \mathbf{M}$ , then  ${}_R T$  is a progenerator and so  $\text{id}({}_T \text{Hom}_R({}_R T_T, {}_R R)) = \text{id}({}_T F({}_R R)) = \text{id}({}_R R)$ . Now, since  $T/R$  is a left (resp. right) quasi-Frobenius extension,  ${}_T \text{Hom}_R({}_R T_T, {}_R R)$  is a generator (resp. finitely generated projective) and therefore  $\text{id}({}_T \text{Hom}_R({}_R T_T, {}_R R)) = \text{id}({}_T T)$ . Also  $\text{id}({}_T T) = \text{id}({}_R R)$  follows from (1).

REMARK. In Proposition 7, if we replace a ring  $T$  by an  $R$ -module and its endomorphism ring, then we obtain the following.

Let  $R$  be a ring,  ${}_R P$  a projective left  $R$ -module,  $T = \text{End}_R(P)$  and assume  $P_T$  flat, then the functor  $F = \text{Hom}_R({}_R P_T, -): {}_R \mathbf{M} \rightarrow {}_T \mathbf{M}$  satisfies 1)–2) by [2, VI Proposition 5.1] and hence  $\text{id}({}_T F(P)) \leq \text{id}({}_R P)$ . Observing this fact,

(i) Let  $R$  be a left noether ring,  ${}_R P$  a projective generator and  $T = \text{End}_R(P)$ , then  $\text{id}({}_T T) \leq \text{id}({}_R R)$ . Therefore it follows immediately that an endomorphism ring of a faithful finitely generated projective module over a quasi-Frobenius ring also is a quasi-Frobenius ring. (Curtis and Morita)

(ii) If rings  $R$  and  $T$  are Morita equivalent, then  $\text{id}({}_R R) = \text{id}({}_T T)$  and  $\text{id}({}_R R) = \text{id}({}_T T)$ .

Now, if rings  $R$  and  $T$  are Morita equivalent, there exists a finitely generated projective generator (i. e. progenerator)  ${}_R P$  and  $T \cong \text{End}_R(P)$ . However, if we delete that  ${}_R P$  is a generator, it happens that  $R$  is Gorenstein but  $T$  is not and we see also that faithfulness in Curtis-Morita theorem above is necessary. For example, let  $R$  be a self-basic serial ring and  $R = Re_1 \oplus Re_2 \oplus Re_3$  a decomposition into primitive left ideals such that  $|Re_1| = |Re_2| = |Re_3| = 5$  and  $Re_1$  (resp.  $Re_2$ ) is epimorphic to  $Ne_2$  (resp.  $Ne_3$ ) where  $N$  is the radical of  $R$ . Then  $R$  is a quasi-Frobenius ring, but  $\text{id}({}_e R_e e R e)$  is infinite for  $e = e_1 + e_2$ .

Finally we state an example concerning with a maximal quotient ring of a Gorenstein ring.

EXAMPLE. It is easily seen that a classical quotient ring or more generally a flat epimorphic extension of a Gorenstein ring also is a Gorenstein ring, but it is not known yet that a maximal quotient ring of a Gorenstein ring is also so. (See [11] in the special case.) Here we present an example of a Gorenstein ring  $R$  whose left maximal quotient ring  $Q$  has  $\text{id}({}_Q Q) > \text{id}({}_R R)$ .

Let  $k$  be a field,  $R$  a subalgebra of  $(k)_5$  whose  $k$ -basis consists of  $c_{11} + c_{55}$ ,  $c_{22} + c_{44}$ ,  $c_{33}$ ,  $c_{31}$ ,  $c_{32}$ ,  $c_{54}$  and  $Q_l$  (resp.  $Q_r$ ) a left (resp. right) maximal quotient ring of  $R$ . Then  $R$  is 1-Gorenstein,  $\text{id}({}_Q Q) = 2$  and  $Q_r$  is a quasi-Frobenius ring.

ACKNOWLEDGEMENT The author wishes to thank the referee for valuable advices and simplification of the proof of Proposition 5.

**References**

- [1] Bass, H.: Injective dimension in Noetherian rings, *Trans. Amer. Math. Soc.* **102** (1962) 19-29.
- [2] Cartan, H. and Eilenberg, S.: "Homological Algebra", Princeton Univ. Press, Princeton, N. J., 1956.
- [3] Eilenberg, S. and Nakayama, T.: On the dimension of modules and algebras II, *Nagoya Math. J.* **9** (1955) 1-16.
- [4] Ischebeck, F.: Eine Dualität zwischen den Funktoren Ext und Tor, *J. Algebra* **11** (1969) 510-531.
- [5] Iwanaga, Y.: On rings with self-injective dimension  $\leq 1$ , *Osaka J. Math.* **15** (1978) 33-45.
- [6] Iwanaga, Y.: On rings with finite self-injective dimension, *Comm. Algebra* **7** (1979) 393-414.
- [7] Jensen, C. U.: On the vanishing of  $\varprojlim^{(i)}$ , *J. Algebra* **15** (1970) 151-166.
- [8] Levin, G. and Vasconcelos, W.: Homological dimensions and Macaulay rings, *Pac. J. Math.* **25** (1968) 315-323.
- [9] Müller, B. J.: Quasi-Frobenius-Erweiterungen, *Math. Z.* **85** (1964) 345-368.
- [10] Roggenkamp, K. W.: Injective modules for group rings and Gorenstein orders, *J. Algebra* **24** (1973) 465-472.
- [11] Sato, H.: On localizations of a 1-Gorenstein ring, *Sci. Rep. Tokyo Kyoiku Daigaku* **13** (1977) 188-193.

Institute of Mathematics  
University of Tsukuba  
Ibaraki, 305 Japan