

Gibbs States of a One Dimensional Quantum Lattice

HUZIHIRO ARAKI*

I.H.E.S., 91 – Bures-sur-Yvette, France

Received June 20, 1969

Abstract. A one dimensional infinite quantum spin lattice with a finite range interaction is studied. The Gibbs state in the infinite volume limit is shown to exist as a primary state of a UHF algebra. The expectation value of any local observables in the state as well as the mean free energy depend analytically on the potential, showing no phase transition. The Gibbs state is an extremal KMS state.

§ 1. Introduction

A one dimensional infinite classical spin lattice system has been studied in [12] and shown to be without any phase transition for a large class of interactions. We show an analogous result for the quantum case with any finite range interaction.

We first show that the power series for the time displacement automorphism of the algebra of observables has an infinite radius of convergence for local observables in one dimensional lattice. This enables us to use the Tomonaga-Schwinger-Dyson perturbation type formula and pull out each potential from $e^{-\beta H}$ as a factor. The transfer matrix technique for the classical one dimensional Ising model is then applicable in a fashion analogous to [12] and we obtain a formula for the infinite volume Gibbs state in terms of an eigen state of a certain linear bounded operator acting on observables.

A standard perturbation theory of bounded linear operators on a Banach space enables us to find an analytic continuation of the Gibbs state with respect to the interaction potential and to prove the analyticity of the expectation value of local observables in the Gibbs state as well as the analyticity of the mean free energy.

This technique is applicable also to the classical case, provided that the interaction potential decreases exponentially at large separation.

The Gibbs state is shown to be invariant under time and lattice translation, satisfies the KMS boundary condition and has the exponen-

* On leave from Research Institute for Mathematical Sciences, Kyoto University Kyoto, Japan.

tial, uniform clustering property. The last property implies, by a general theorem, that the state is primary (a factor state) and is an extremal KMS state.

§ 2. Notation and Results

We represent a one dimensional lattice by the set of integers $Z = \{0, \pm 1, \dots\}$. For each lattice point j , we have a d -dimensional Hilbert space \mathcal{H}_j where d is finite and independent of j . For each finite subset I of Z , we consider a finite dimensional full matrix algebra $\mathfrak{A}(I) = \mathcal{B}(\mathcal{H}(I))$, $\mathcal{H}(I) = \bigotimes_{j \in I} \mathcal{H}_j$. For $I \subset I'$, $Q \in \mathfrak{A}(I)$ is identified with $Q \otimes 1_{I' \setminus I}$ in $\mathfrak{A}(I')$ where $I' \setminus I$ denotes the complement of I in I' and $1_{I' \setminus I}$ is the identity in $\mathfrak{A}(I' \setminus I)$. The collection of $\mathfrak{A}(I)$ for all finite subsets I of Z together with this identification defines a normed *-algebra \mathfrak{A}_0 (the algebra of local observables). Its completion \mathfrak{A} is taken as the C*-algebra of quasi local observables. The closed *-subalgebra of \mathfrak{A} generated by all $\mathfrak{A}(I)$, $I \subset I'$ will be denoted by $\mathfrak{A}(I')$ for an infinite subset I' of Z (as well as for a finite subset I').

For any two lattice points j and j' , we fix a unitary mapping $w(j, j')$ from $\mathcal{H}_{j'}$ onto \mathcal{H}_j such that $w(j, j') w(j', j'') = w(j, j'')$, $w(j, j')^* = w(j', j)$ and $w(j, j) = 1$. Let $I + a$ denote the set $\{j + a; j \in I\}$. Let $w(I + a, I) = \bigotimes_{j \in I} w(j + a, j)$, which is a unitary map of $\mathcal{H}(I)$ onto $\mathcal{H}(I + a)$. The *-automorphism of \mathfrak{A} , which is induced by a *-isomorphism $Q \in \mathfrak{A}(I) \rightarrow w(I + a, I) Q w(I + a, I)^*$, is denoted by $\tau_s(a)$ and is called a lattice translation. We also need the *-isomorphism of $\mathfrak{A}(Z \setminus [1 - n, n])$ onto \mathfrak{A} , which is induced by the *-isomorphisms $\tau_s(n) \otimes \tau_s(-n)$ of $\mathfrak{A}([n + 1, N]) \otimes \mathfrak{A}([1 - N, -n])$ onto $\mathfrak{A}([1, N - n]) \otimes \mathfrak{A}([1 - N + n, 0])$, $N = n + 1, n + 2, \dots$. It is denoted by $\tau_c(n)$. Here $[a, b]$ denotes the set of integers j satisfying $a \leq j \leq b$.

Let $\Phi(I)$ be the interaction potential among lattice sites in I . $\Phi(I)^* = \Phi(I)$, $\Phi(I) \in \mathfrak{A}(I)$. We require $\Phi(I) = 0$ if I is not within some interval of length r . We also require $\Phi(I + a) = \tau_s(a) \Phi(I)$. The Hamiltonian for a finite interval $[N_1, N_2]$ is

$$U(N_1, N_2) = \sum_{I \subset [N_1, N_2]} \Phi(I). \tag{2.1}$$

The Gibbs state $\varphi_{a,b}^G$ of $\mathfrak{A}([a, b])$ is

$$\varphi_{a,b}^G(Q) = Z(a, b)^{-1} \text{tr}_{ab}[Q \exp - U(a, b)], \tag{2.2}$$

$$Z(a, b) = \text{tr}_{ab}[\exp - U(a, b)], \tag{2.3}$$

where tr_{ab} is the trace of a full matrix algebra $\mathfrak{A}([a, b])$.

Theorem 2.1. (i) *The following limit exists and is finite:*

$$P(\Phi) = \lim_{b \rightarrow a \rightarrow \infty} (b - a)^{-1} \log Z(a, b). \tag{2.4}$$

It is holomorphic in Φ .

(ii) *For every $Q \in A_0$, the following limit exists and defines a state of \mathfrak{A} :*

$$\varphi_\Phi(Q) = \lim_{a \rightarrow -\infty, b \rightarrow +\infty} \varphi_{ab}^\Phi(Q). \tag{2.5}$$

For each $Q \in \mathfrak{A}_0$, it is holomorphic in Φ .

The holomorphy in Φ means the real holomorphy in $\zeta = (\zeta_1, \dots, \zeta_n)$ when each $\Phi(I) = \Phi(I, \zeta)$ is a restriction of an $\mathfrak{A}(I)$ valued holomorphic function $\Phi(I, \zeta)$ to $\zeta = \xi \in$ some real domain.

The analyticity statement can be proved for the following class of quasilocals observables with exponential tail.

Definition 2.2. *Let $Q \in \mathfrak{A}$.*

$$\|Q\|_n = \inf_{Q_n \in \mathfrak{A}([-n, n])} \|Q - Q_n\|. \tag{2.6}$$

$$\|Q\|_{n,x} = \sum_{k=n}^\infty x^k \|Q\|_k. \tag{2.7}$$

$$\| \|Q\|_{n,x} \| = \|Q\| + \|Q\|_{n,x}. \tag{2.8}$$

For $x > 1$, $\mathfrak{A}(x)$ is the set of $Q \in \mathfrak{A}$ such that $\|Q\|_{n,x} < \infty$.

$\varphi_\Phi(Q)$ is holomorphic in Φ if $Q \in \mathfrak{A}(x)$ for some $x > 1$.

The limit

$$\tau_T(t)Q = \lim_{a \rightarrow -\infty, b \rightarrow \infty} e^{itU(a,b)} Q e^{-itU(a,b)} \tag{2.9}$$

exists in \mathfrak{A} for all $Q \in \mathfrak{A}$ and defines a continuous one parameter group of * automorphisms of \mathfrak{A} , which we denote by $\tau_T(t)$ and call a time translation. (The unit of time is $(\beta\hbar)^{-1}$.) A state φ of \mathfrak{A} is time and lattice translation invariant if $\varphi(\tau_T(t) \tau_s(a) Q) = \varphi(Q)$, for any t, a and $Q \in \mathfrak{A}$. It satisfies the KMS boundary condition if $\varphi(Q_1 Q_2(f_0)) = \varphi(Q_2(f_1) Q_1)$ for all $Q_1, Q_2 \in \mathfrak{A}$ and $f \in \mathcal{D}$ where $Q_2(f_\alpha) = \int \tau_T(t) Q_2 f_\alpha(t) dt, f_\alpha(t) = \int_{-\infty}^\infty e^{-ist + \alpha s} f(s) ds$. Let S_1 be a convex subset of set of states of \mathfrak{A} . A state

$\varphi \in S_1$ is extremal in S_1 if $\varphi = \lambda\varphi_1 + (1 - \lambda)\varphi_2, \varphi_1, \varphi_2 \in S_1, 0 < \lambda < 1$ imply $\varphi_1 = \varphi_2 = \varphi$. If S_1 is the set of translation invariant states, φ is an extremal translation invariant state and if S_1 is the set of time translation invariant states satisfying the KMS boundary condition, then φ is an extremal KMS state.

A state φ has an exponential clustering property if there exists $\varrho > 0$ such that

$$\lim_{n \rightarrow \infty} e^{\varrho|n|} [\varphi(Q_1 \tau_s(n) Q_2) - \varphi(Q_1) \varphi(\tau_s(n) Q_2)] = 0 \tag{2.10}$$

for any fixed $Q_1, Q_2 \in \mathfrak{A}_0$. φ has a uniform clustering property if there exists N for any given $\varepsilon > 0$ and $Q_1 \in \mathfrak{A}$ such that

$$|\varphi(Q_1 Q_2) - \varphi(Q_1) \varphi(Q_2)| < \varepsilon \|Q_2\| \tag{2.11}$$

for all $Q_2 \in \mathfrak{A}(Z \setminus [-N, N])$. A state φ is primary (or a factor state) if the cyclic representation π_φ of \mathfrak{A} associated with φ through the GNS construction is such that $\pi_\varphi(\mathfrak{A})''$ is a factor (i.e. the center of $\pi_\varphi(\mathfrak{A})''$ consists of multiples of the identity).

Theorem 2.3. *The Gibbs state φ_Φ for the infinite system is invariant under time and lattice translation, satisfies the KMS boundary condition, has exponential and uniform clustering properties, is primary, is an extremal KMS state, and is an extremal lattice translation invariant state.*

In the following discussion, we use the following combination of $\Phi(I)$:

$$\Phi = \sum_{I \subset [0, r]} n_r(I)^{-1} \Phi(I) \tag{2.12}$$

where $n_r(I)$ is the number of the translates $I + a$ of I which is still in $[0, r]$. We denote

$$H(I) = \sum_{[n, r+n] \subset I} \tau_s(n) \Phi, \tag{2.13}$$

$$H(a, b) = H([a, b]). \tag{2.14}$$

$H(a, b)$ and $U(a, b)$ differs only near the two ends:

$$U(a, b) - H(a, b) = \Delta_a^- + \Delta_b^+, \tag{2.15}$$

$$\Delta_b^+ \in A([b - r, b]), \quad \Delta_a^- \in A([a, a + r]), \tag{2.16}$$

$$\|\Delta_b^+\| \leq \sum_{I \subset [0, r]} \|\Phi(I)\|, \tag{2.17}$$

$$\|\Delta_a^-\| \leq \sum_{I \subset [0, r]} \|\Phi(I)\|. \tag{2.18}$$

§ 3. The Spaces $\mathfrak{A}(M, x)$

Lemma 3.1. $\|Q\|_{n,x}$ is a norm of linear space $\mathfrak{A}(x)$. $\mathfrak{A}(x)$ equipped with the norm $\| \cdot \|_{n,x}$ (denoted as $\mathfrak{A}(n, x)$) is a * Banach algebra.

Proof. There always exists $Q^{(n)} \in \mathfrak{A}([-n, n])$ such that $\|Q - Q^{(n)}\| = \|Q\|_n$ due to the compactness of a bounded closed subset of $A([-n, n])$. Let

$\|Q_1 - Q_1^{(n)}\| = \|Q_1\|_n$, $\|Q_2 - Q_2^{(n)}\| = \|Q_2\|_n$. Then

$$\|Q_1 + Q_2\|_n \leq \|Q_1 + Q_2 - Q_1^{(n)} - Q_2^{(n)}\| \leq \|Q_1\|_n + \|Q_2\|_n. \quad (3.1)$$

Similarly, $\|CQ\|_n \leq |C| \|Q\|_n$. At the same time, $\|Q\|_n \leq |C^{-1}| \|CQ\|_n$. Therefore $\|CQ\|_n = |C| \|Q\|_n$.

Obviously $\|Q^*\|_{n,x} = \|Q\|_{n,x}$. Further, $\|Q_1 Q_2\|_n \leq \|Q_1 Q_2 - Q_1^{(n)} Q_2^{(n)}\| \leq \|Q_1\| \|Q_2\|_n + \|Q_2^{(n)}\| \|Q_1\|_n \leq \|Q_1\|_n \|Q_2\|_n + \|Q_1\| \|Q_2\|_n + \|Q_1\|_n \|Q_2\|_n$. Hence $\|Q_1 Q_2\|_{n,x} \leq \|Q_1\|_{n,x} \|Q_2\|_{n,x}$.

Any Cauchy sequence Q_k with respect to $\| \cdot \|_{n,x}$ is a Cauchy sequence with respect to the norm in \mathfrak{A} and has a limit Q in \mathfrak{A} : $\lim_k \|Q_k - Q\| = 0$.

This implies $\|Q\|_n \leq \liminf_k \|Q_k\|_n \leq \overline{\lim}_k \|Q_k\|_n \leq \|Q\|_n$ and hence $\|Q\|_n = \lim_k \|Q_k\|_n$. Hence

$$\|Q\| + \sum_{l=n}^N x^l \|Q\|_l = \lim_{k \rightarrow \infty} \left(\|Q_k\| + \sum_{l=n}^N x^l \|Q_k\|_l \right) \leq \sup_k \|Q_k\|_{n,x} < \infty. \quad (3.2)$$

Thus $\|Q\|_{n,x} < \infty$ and $Q \in \mathfrak{A}(n, x)$. Given $\varepsilon > 0$. There exists K such that $\|Q_k - Q_k\|_{n,x} < \varepsilon/4$ for $k \geq K$. There exists N such that $\|Q_k\|_{N,x} < \varepsilon/4$ and $\|Q\|_{N,x} < \varepsilon/4$. There exists K' such that $\|Q - Q_k\| + \sum_{l=n}^{N-1} x^l \|Q - Q_k\|_l < \varepsilon/4$ for $k \geq K'$. Then $\|Q - Q_k\|_{n,x} < \varepsilon$ for $k \geq \max(K, K')$. Therefore $\lim_k \|Q_k - Q\|_{n,x} = 0$. Q.E.D.

Lemma 3.2. *The set $\Sigma_N(\gamma)$ of $Q \in \mathfrak{A}$ such that*

$$\|Q\| \leq \gamma_0, \quad \|Q\|_l \leq \gamma_l, \quad l = N, N+1, \dots \quad (3.3)$$

is convex and compact provided that $\gamma_0 < \infty$, $\lim_{l \rightarrow \infty} \gamma_l = 0$.

Proof. The convexity is straight forward from the triangular inequality. By setting $Q_n = 0$ in (2.6), we obtain $\|Q\|_n \leq \|Q\| \leq \gamma_0$. Let $Q^{(n)} \in \mathfrak{A}([-n, n])$ be such that $\|Q - Q^{(n)}\| = \|Q\|_n$. Then $\|Q^{(n)}\| \leq \|Q\| + \|Q\|_n \leq 2\gamma_0$.

Let $Q_k \in \Sigma_N(\gamma)$. Let $k(i, j)$, $j = 1, 2, \dots$ be a subsequence of $k(i-1, j)$, $j = 1, 2, \dots$ such that $Q_k^{(i, j)}$ is convergent in norm as $j \rightarrow \infty$, where $n(N-1, k) = k$ and $i = N, N+1, \dots$. Such choice is inductively possible because $\|Q_k^{(i)}\| \leq 2\gamma_0$ for all i and k .

We can show that $Q_{k(j, j)}$ is a Cauchy sequence in \mathfrak{A} . Let $\varepsilon > 0$ be given. There exist K such that $\gamma_l < \varepsilon/3$ for $l \geq K$ and $K' > K$ such that $\|Q_k^{(K, j)} - Q_k^{(K, j')}\| < \varepsilon/3$ for $j, j' \geq K'$. Since $k(j, j) = k(K, v)$ with $v \geq j$ if $j \geq K$, we have $\|Q_k^{(K, j)} - Q_k^{(K, j')}\| < \varepsilon/3$ if $j, j' \geq K'$. Hence

$$\begin{aligned} & \|Q_{k(j, j)} - Q_{k(j', j')}\| \leq \|Q_{k(j, j)} - Q_{k(j, j)}^{(K)}\| \\ & + \|Q_{k(j, j)}^{(K)} - Q_{k(j', j')}^{(K)}\| + \|Q_{k(j', j')}^{(K)} - Q_{k(j', j')}\| < \varepsilon \end{aligned} \quad (3.4)$$

if $j, j' \geq K'$. Hence Q_k has a convergent subsequence $Q_{k(j, j)}$.

Since \mathfrak{A} is a separable Banach space, its sequentially compact subset is compact. Q.E.D.

Lemma 3.3. *Let $x_2 > x_1 > 1$. The closure of a bounded subset of $\mathfrak{A}(n, x_2)$ with respect to the norm in \mathfrak{A} is in $\mathfrak{A}(n, x_2)$ and is a compact subset of $\mathfrak{A}(n, x_1)$ (with respect to $\| \cdot \|_{n, x_1}$).*

Proof. Let $\|Q_k\|_{n, x_2} \leq a$. Then $\|Q_k\| \leq a$, $\|Q_k\|_l \leq x_2^{-l} a$ for $l = n, n + 1, \dots$ and $\lim_{l \rightarrow \infty} x_2^{-l} a = 0$. Hence Q_k has a subsequence which converges with respect to the norm in \mathfrak{A} . Let now Q_k be a sequence such that $\lim_{k \rightarrow \infty} \|Q - Q_k\| = 0$. By the latter half of the proof of Lemma 3.1, $\|Q\|_n = \lim_k \|Q_k\|_n$ and hence $\|Q\| + \sum_{l=n}^{N-1} \|Q\|_l \leq a$. Therefore $\|Q\|_{n, x_2} \leq a$. Since $\|Q_k\|_l \leq x_2^{-l} a$, $l \geq n$, we have $\|Q_k\|_{N, x_1} \leq a(x_1/x_2)^N (1 - (x_1/x_2))^{-1}$, which tend to 0 as $N \rightarrow \infty$. This is true also when Q_k is replaced by Q . Since $\lim \|Q - Q_k\|_l = 0$ for each l , we have $\lim \|Q - Q_k\|_{n, x_1} = 0$. Q.E.D.

In the above discussion $\mathfrak{A}(n, x)$ for given x and varying n are topologically equivalent. We introduced $\| \cdot \|_{n, x}$ merely for the convenience in later computation.

Definition 3.4. \mathfrak{A}_1 and \mathfrak{A}_2 are sets of $Q \in \mathfrak{A}$ satisfying the following conditions (i) and (ii), respectively.

- (i) $\sum_l x^l \|Q\|_l < \infty$ for all x .
- (ii) $\sup_n n^{-1} [\log \|Q\|_n + (n/r) \log n] < \infty$.

Lemma 3.5. \mathfrak{A}_1 and \mathfrak{A}_2 are *-subalgebra of \mathfrak{A} stable under $\tau_s(a)$. \mathfrak{A}_1 contains \mathfrak{A}_2 .

Proof. (i) Let $Q_1, Q_2 \in \mathfrak{A}$. Then

$$\sum x^l \|c_1 Q_1 + c_2 Q_2\|_l \leq |c_1| \sum x^l \|Q_1\|_l + |c_2| \sum x^l \|Q_2\|_l, \tag{3.5}$$

$$\begin{aligned} & \sup_n n^{-1} [\log \|c_1 Q_1 + c_2 Q_2\|_n + (n/r) \log n] \\ & \leq \sup_n n^{-1} [\log(|c_1| \|Q_1\|_n + |c_2| \|Q_2\|_n) + (n/r) \log n] \tag{3.6} \\ & = \max_{j=1,2} \left\{ \sup_n n^{-1} [\log \|Q_j\|_n + (n/r) \log n] \right\} + \log(|c_1| + |c_2|). \end{aligned}$$

Hence \mathfrak{A}_1 and \mathfrak{A}_2 are linear subset of \mathfrak{A} . Next

$$\begin{aligned} \|Q_1 Q_2\|_n & \leq \|Q_1 Q_2 - Q_1^{(n)} Q_2^{(n)}\| \leq \|Q_1\| \|Q_2\|_n + \|Q_2^{(n)}\| \|Q_1\|_n \\ & \leq \|Q_1\| \|Q_2\|_n + 2 \|Q_2\| \|Q_1\|_n. \tag{3.7} \end{aligned}$$

Hence

$$\Sigma x^l \|Q_1 Q_2\|_l \leq \|Q_1\| \Sigma x^l \|Q_2\|_l + 2 \|Q_2\| \Sigma x^l \|Q_1\|_l, \tag{3.8}$$

$$\sup_n n^{-1} [\log \|Q_1 Q_2\|_n + (n/r) \log n] \leq \max_{j=1,2} \left\{ \sup_n n^{-1} [\log \|Q_j\|_n + (n/r) \log n] \right\} + \log(\|Q_1\| + 2 \|Q_2\|). \tag{3.9}$$

Therefore \mathfrak{A}_1 and \mathfrak{A}_2 are algebras. Since $\|Q^*\|_l = \|Q\|_l$, \mathfrak{A}_1 and \mathfrak{A}_2 are *-algebras. $\|\tau_s(a) Q\|_l \leq \|Q\|_{l+a}$. Therefore \mathfrak{A}_1 and \mathfrak{A}_2 are stable under $\tau_s(a)$. $\mathfrak{A}_2 \subset \mathfrak{A}_1$ is obvious. Q.E.D.

Definition 3.6. Let $Q \in \mathfrak{A}$. Define $\|Q^{-1}\|$ to be ∞ if $Q^{-1} \notin \mathfrak{A}$.

$$\alpha(Q) = \|Q\| \|Q^{-1}\|, \tag{3.10}$$

$$\alpha_n(Q) = \inf_{Q_n \in \mathfrak{A}((-n, n))} \|Q - Q_n\| \|Q_n^{-1}\|. \tag{3.11}$$

Lemma 3.7. Let $Q > 0$. Then

$$\|Q^{-1}\|^{-1} \leq Q \leq \|Q\|, \tag{3.12}$$

$$\alpha(Q) = \sup \varphi'(Q) / \varphi''(Q) \tag{3.13}$$

where sup is taken over all states φ' and φ'' of \mathfrak{A} .

Proof. $\|Q\|$ is the l.u.b. of the spectrum of Q and $\|Q^{-1}\|$ is the l.u.b. of the spectrum of Q^{-1} , which is the inverse of the g.l.b. of the spectrum of Q . Hence (3.12) follows. Since $\sup_{\varphi} \varphi(Q) = \|Q\|$, $\inf_{\varphi} \varphi(Q) = \|Q^{-1}\|^{-1}$ for $Q > 0$, we have (3.13).

Lemma 3.8. If X and Y are elements of a Banach algebra, X^{-1} exists and $\|Y\| \|X^{-1}\| = \delta < 1$, then $X + Y$ has an inverse $\|(X + Y)^{-1}\| \leq \|X^{-1}\| (1 - \delta)^{-1}$, and

$$\|(X + Y)^{-1} - X^{-1}\| \leq \|X^{-1}\| \{(1 - \delta)^{-1} - 1\}. \tag{3.14}$$

Proof. Consider the series

$$f = X^{-1} \sum_{n=0}^{\infty} (-YX^{-1})^n = \left(\sum_{n=0}^{\infty} (-X^{-1}Y)^n \right) X^{-1} \tag{3.15}$$

which is absolutely convergent due to $\|YX^{-1}\| \leq \|Y\| \|X^{-1}\| = \delta < 1$. It satisfies $(X + Y)f = f(X + Y) = 1$ and hence $f = (X + Y)^{-1}$. Further $\|f - X^{-1}\| \leq \|X^{-1}\| \sum_{n=1}^{\infty} \delta^n \leq \|X^{-1}\| \{(1 - \delta)^{-1} - 1\}$.

Lemma 3.9. Let $Q > 0$, $Q \in \mathfrak{A}$.

$$(1) \quad \alpha_l(Q) \leq (\alpha(Q) - 1)(\alpha(Q) + 1)^{-1} < 1. \tag{3.16}$$

$$(2) \quad \alpha_l(Q) \leq \alpha_{l'}(Q) \text{ if } l \leq l'. \tag{3.17}$$

(3) *There exists $Q_{(l)} \in \mathfrak{A}([-l, l])$, such that*

$$\|Q - Q_{(l)}\| \|Q_{(l)}^{-1}\| = \alpha_l(Q) \quad \text{and} \\ \|Q^{-1}\| (1 - \alpha_l(Q)) \leq \|Q_l^{-1}\| \leq \|Q^{-1}\| (1 + \alpha_l(Q)). \quad (3.18)$$

$$(4) \quad \|Q\|_l \leq (\|Q\| - \|Q^{-1}\|^{-1})/2. \quad (3.19)$$

$$(5) \quad \|Q\|_l \leq \|Q\|_{l'} \quad \text{if } l' \leq l. \quad (3.20)$$

$$(6) \quad \alpha_l(Q) \leq \|Q\|_l \|Q^{-1}\| (1 - \|Q\|_l \|Q^{-1}\|)^{-1}. \quad (3.21)$$

$$(7) \quad \|Q\|_l \leq \|Q^{-1}\|^{-1} \alpha_l(Q) (1 - \alpha_l(Q))^{-1}. \quad (3.22)$$

$$(8) \quad \alpha(\lambda Q) = \alpha(Q), \quad \alpha_l(\lambda Q) = \alpha_l(Q) \quad \text{if } \lambda > 0. \quad (3.23)$$

$$(9) \quad \|Q + x\|_l = \|Q\|_l. \quad (3.24)$$

(10) *If $\lambda \|Q^{-1}\| (1 + \alpha_l(Q)) < 1$, then*

$$\alpha_l(Q - \lambda) \leq \alpha_l(Q) (1 - \lambda \|Q^{-1}\| (1 + \alpha_l(Q)))^{-1}, \quad \lambda > 0. \quad (3.25)$$

$$\alpha(Q - \lambda) \leq (\alpha(Q) - \lambda \|Q^{-1}\|) (1 - \lambda \|Q^{-1}\|)^{-1}. \quad (3.26)$$

Proof. (1) If we set $Q^{(0)} = \{\|Q\| + \|Q^{-1}\|^{-1}\}/2$ then

$$\alpha_l(Q) \leq \|Q - Q^{(0)}\| \|Q^{(0)}\|^{-1} \\ = (\|Q\| - \|Q^{-1}\|^{-1}) / (\|Q\| + \|Q^{-1}\|^{-1}) = (\alpha(Q) - 1) (\alpha(Q) + 1)^{-1}. \quad (3.27)$$

(4) follows from $\|Q\|_l \leq \|Q - Q^{(0)}\| = (\|Q\| - \|Q^{-1}\|^{-1})/2$. (2) and (5) are obvious. (3). Since $\|Q_l^{-1}\| = \sup \|\psi\| \|\pi(Q_l)\psi\|^{-1} \geq \|Q_l\|^{-1}$ where the sup is over all non zero vectors in a faithful representation π of \mathfrak{A} . Hence as $\|Q_l\| \rightarrow \infty$, $\|Q - Q_l\| \|Q_l^{-1}\| \geq \|Q_l\| \|Q_l^{-1}\| (1 - \|Q\|/\|Q_l\|)$ becomes ≥ 1 . By (3.16), if $\|Q - Q_l\| \|Q_l^{-1}\| < \alpha_l(Q) + \varepsilon$ with $\varepsilon < (1 - \alpha_l(Q))/2$, then $\|Q_l\|$ is bounded by a constant. Hence there exists $Q_{(l)} \in \mathfrak{A}([-l, l])$ such that $\|Q - Q_{(l)}\| \|Q_{(l)}^{-1}\| = \alpha_l(Q)$ by the compactness of a bounded closed set in $\mathfrak{A}([-l, l])$. We have

$$\|Q Q_{(l)}^{-1} - 1\| \leq \|Q - Q_{(l)}\| \|Q_{(l)}^{-1}\| \leq \alpha_l(Q). \quad (3.28)$$

Hence $1 - \alpha_l(Q) \leq \|Q Q_l^{-1}\| \leq 1 + \alpha_l(Q)$ and

$$\|Q_{(l)}^{-1}\| \leq \|Q^{-1}\| \|Q Q_{(l)}^{-1}\| \leq \|Q^{-1}\| (1 + \alpha_l(Q)). \quad (3.29)$$

From (3.28) and Lemma 3.8, $\|Q_{(l)} Q^{-1}\| = \|(1 + (Q Q_{(l)}^{-1} - 1))^{-1}\| \leq (1 - \alpha_l(Q))^{-1}$. Therefore

$$\|Q_{(l)}^{-1}\| \geq \|Q^{-1}\| \|Q_{(l)} Q^{-1}\|^{-1} \geq \|Q^{-1}\| (1 - \alpha_l(Q)). \quad (3.30)$$

By Lemma 3.8, we have

$$\|(Q^{(0)})^{-1}\| \leq \|Q^{-1}\| (1 - \|Q - Q^{(0)}\| \|Q^{-1}\|)^{-1}. \quad (3.31)$$

Hence we have

$$\alpha_l(Q) \leq \|Q - Q^{(l)}\| \|(Q^{(l)})^{-1}\| \leq \|Q^{-1}\| \|Q\|_l (1 - \|Q\|_l \|Q^{-1}\|)^{-1} \quad (3.32)$$

where $Q^{(l)} \in \mathfrak{A}([-l, l])$, $\|Q - Q^{(l)}\| = \|Q\|_l$. This proves (6). (7) follows from (3).

(8) and (9) are immediate from definitions. To obtain (10), we note that if $Q_{(l)}$ is given by (3), then, by Lemma 3.8,

$$\|(Q_{(l)} - \lambda)^{-1}\| \leq \|Q_{(l)}^{-1}\| (1 - |\lambda| \|Q_{(l)}^{-1}\|)^{-1}$$

and hence

$$\begin{aligned} \alpha_l(Q - \lambda) &\leq \|Q - Q_{(l)}\| \|(Q_{(l)} - \lambda)^{-1}\| \\ &\leq \|Q - Q_{(l)}\| \|Q_{(l)}^{-1}\| (1 - |\lambda| \|Q_{(l)}^{-1}\|)^{-1} \\ &\leq \alpha_l(Q) (1 - |\lambda| \|Q^{-1}\| (1 + \alpha_l(Q)))^{-1} \end{aligned} \quad (3.33)$$

where we have used (3.29).

Since $\alpha(Q) \geq \|Q^{-1}\| Q \geq 1$, $\alpha(Q) - \lambda \|Q^{-1}\| \geq \|Q^{-1}\| (Q - \lambda) \geq 1 - \|Q^{-1}\| \lambda$. Hence we have (3.26).

Lemma 3.10. *Let v be a state of \mathfrak{A} , $a > 0$, $1 < \alpha < \infty$, $\lim \alpha_l = 0$, $\alpha_l > 0$. Let Σ be the set of $Q \in \mathfrak{A}$, $Q \geq 0$ such that $v(Q) = a$, $\alpha(Q) \leq \alpha$, $\|Q\|_l \|Q^{-1}\| \leq \alpha_l$, $l = N, N + 1, \dots$. Then Σ is a convex, compact subset of \mathfrak{A} .*

Proof. First we prove the convexity. Let $Q = \lambda Q_1 + (1 - \lambda) Q_2$, $Q_1 \in \Sigma$, $Q_2 \in \Sigma$, $0 \leq \lambda \leq 1$. $v(Q) = \lambda v(Q_1) + (1 - \lambda) v(Q_2) = a$. Since $Q_1 \geq 0$, $Q_2 \geq 0$, we have $Q \geq 0$.

$$\|Q\| \leq \lambda \|Q_1\| + (1 - \lambda) \|Q_2\| \leq \alpha \{ \lambda \|Q_1^{-1}\|^{-1} + (1 - \lambda) \|Q_2^{-1}\|^{-1} \}. \quad (3.34)$$

$$\begin{aligned} \|Q^{-1}\|^{-1} &= \inf_{\varphi} \varphi(Q) \geq \lambda \inf \varphi(Q_1) \\ &\quad + (1 - \lambda) \inf \varphi(Q_2) \geq \lambda \|Q_1^{-1}\|^{-1} + (1 - \lambda) \|Q_2^{-1}\|^{-1}. \end{aligned} \quad (3.35)$$

Hence $\alpha(Q) \leq \alpha$. Similarly,

$$\|Q\|_n \leq \lambda \|Q_1\|_n + (1 - \lambda) \|Q_2\|_n \leq \alpha_n (\lambda \|Q_1^{-1}\|^{-1} + (1 - \lambda) \|Q_2^{-1}\|^{-1}). \quad (3.36)$$

Hence $\|Q\|_l \|Q^{-1}\| \leq \alpha_l$.

Next we prove the compactness. From $v(Q) = a \geq \|Q^{-1}\|^{-1} \geq \alpha^{-1} \|Q\|$, we have $\|Q\| \leq a\alpha$. Similarly $\|Q\|_l \leq \alpha_l \|Q^{-1}\|^{-1} = \alpha_l a$. Therefore Σ is a subset of a compact set. We now prove that Σ is closed. Let $Q_n \in \Sigma$, $\lim_{n \rightarrow \infty} \|Q - Q_n\| = 0$. We have $v(Q) = \lim v(Q_n) = a$. From $Q_n \geq \|Q_n^{-1}\|^{-1} \geq \alpha^{-1} \|Q_n\| \geq \alpha^{-1} a$, we have $Q \geq \alpha^{-1} a$. Hence by Lemma 3.8, $\lim_{n \rightarrow \infty} \|Q_n^{-1}\| = \|Q^{-1}\|$. Hence $\lim_{n \rightarrow \infty} \alpha(Q_n) = \alpha(Q) \leq \alpha$, $\lim_{n \rightarrow \infty} \|Q_n\|_l \|Q_n^{-1}\| = \|Q\|_l \|Q^{-1}\| \leq \alpha_l$. Q.E.D.

Corollary 3.11. *Let v be a state of \mathfrak{A} , $x > 1$, $1 < \alpha < \infty$, $a > 0$. The set of $Q \in \mathfrak{A}$, $Q \geq 0$ such that $v(Q) = a$, $\|Q\|_{M,x} \|Q^{-1}\| \leq \alpha$ is a convex compact subset of \mathfrak{A} .*

Proof. From $\|Q\|_{M,x} \|Q^{-1}\| \leq \alpha$, we have $\alpha(Q) \leq \alpha$, $\|Q\|_l \|Q^{-1}\| \leq x^{-l} \alpha$, $l = M, M + 1, \dots$. The set is closed and hence compact. From (3.34)~(3.36), $\| \lambda_1 Q_1 + \lambda_2 Q_2 \|_{M,x} \| (\lambda_1 Q_1 + \lambda_2 Q_2)^{-1} \| \leq a$ if Q_1 and Q_2 are in the set, $\lambda_1 + \lambda_2 = 1$, $\lambda_1 \geq 0$, $\lambda_2 \geq 0$. Hence it is convex.

Corollary 3.12. *The set of $Q \in \mathfrak{A}$, $Q \geq 0$ such that $v(Q) = a$, $\alpha(Q) \leq \alpha$, $\alpha_l(Q) \leq \alpha_l$, $l = N, N + 1, \dots$ is a subset of a compact convex set, if $\lim \alpha_l = 0$.*

Proof. This follows from (3.22) and Lemma 3.10.

Remark. In [12], quantities of the form $\beta_l(Q) = \sup_{\varphi, \varphi', \varphi''} \varphi \otimes \varphi'(Q) / \varphi \otimes \varphi''(Q)$

has been used instead of our $\alpha_l(Q)$, where φ runs over states of $\mathfrak{A}([-n, n])$ and φ', φ'' runs over states of $\mathfrak{A}(Z \setminus [-n, n])$. $\beta_l(\lambda Q_1 + (1 - \lambda) Q_2) \leq \max\{\beta_l(Q_1), \beta_l(Q_2)\}$ for $Q_1, Q_2 \geq 0, 0 \leq \lambda \leq 1$. Hence the condition $\beta_l(Q) \leq \beta_l$ is stable under convex combination of Q . \mathfrak{A}_1 is a Fréchet Montel space.

§ 4. Time Translation

For any two elements Q and R in \mathfrak{A} , define

$$\delta(R) Q = [R, Q], \tag{4.1}$$

$$\{\exp \delta(R)\} Q = \sum_{n=0}^{\infty} (n!)^{-1} \delta(R)^n Q. \tag{4.2}$$

We extend this definition to $R = H(I)$

$$\delta(H(I)) Q = \sum_{j: [j, j+r] \subset I} [\tau_s(j) \Phi, Q]. \tag{4.3}$$

If $Q \in \mathfrak{A}_0$, the sum terminates at finite j .

Lemma 4.1. *Let $C_l^m(n)$ be numbers such that*

$$C_l^{m+1}(n) = (l - r + 1) C_l^m(n) + 2 \sum_{k=1}^r C_{l-k}^m(n) \tag{4.4}$$

$$C_l^0(n) = \delta_{ln} \text{ (i.e. } = 1 \text{ if } l = n, = 0 \text{ if } l \neq n) \tag{4.5}$$

where $l, m \in Z, m \geq 0, n \geq r - 1$. Then $C_l^m(n) \geq 0$, and

$$C_l^m(n) = 0 \text{ if } l < n \text{ or } l > n + mr. \tag{4.6}$$

Let

$$f_n(x, y) = \sum_{m,l} C_l^m(n) x^{l-n} y^{mr+n-l} / m!. \tag{4.7}$$

Then, (4.7) is absolutely convergent for all x and y , and is given by

$$f_n(x, y) = \exp[(n - r + 1) y^r + 2 \sum_{k=1}^r k^{-1} (x/y)^k \{\exp k y^r - 1\}]. \quad (4.8)$$

Let

$$F_n(x) = \sum_{m,l} C_l^m(n) x^m/m!, \quad (4.9)$$

$$F_n^L(x) = \sum_m \sum_{l>L} C_l^m(n) x^m/m!. \quad (L \geq n.) \quad (4.10)$$

Then

$$F_n(x) = \exp[(n - r + 1) x + 2 \sum_{k=1}^r k^{-1} \{\exp k x - 1\}], \quad (4.11)$$

$$F_n^{Lr+n}(x) \leq (L + 1)!^{-1} \left[2 \sum_{k=1}^r k^{-1} (e^{kx} - 1) \right]^{L+1} F_n(x) \quad (x > 0), \quad (4.12)$$

where $L \geq 0$.

Proof. $C_l^m(n) \geq 0$ and (4.6) are immediate from (4.4) and (4.5) by induction on m . By (4.4) and (4.6), we have

$$\sup_l |C_l^{m+1}(n)| \leq \{(n + mr) - r + 1 + 2r\} \sup_l |C_l^m(n)|. \quad (4.13)$$

By repetition, we have $\sup_l |C_l^m(n)| \leq \prod_{k=1}^m (n + 1 + kr)$. Therefore (4.7) is absolutely convergent near $x = y = 0$ and defines a holomorphic function.

Due to (4.4), $f_n(x, y)$ satisfies within the polycircle of convergence the following partial differential equation

$$\begin{aligned} (ry^r)^{-1} [x(\partial/\partial x) + y(\partial/\partial y)] f_n \\ = \{x(\partial/\partial x) + n - r + 1\} f_n + 2 \sum_{k=1}^r (x/y)^k f_n. \end{aligned} \quad (4.14)$$

From (4.5), we have the initial condition

$$f(0, 0) = 1. \quad (4.15)$$

After the change of variables

$$s = \log x, \quad t = \log y - y^r, \quad u = (s + t)/2, \quad v = (s - t)/2, \quad (4.16)$$

we have

$$(\partial/\partial u) f_n(x, y) = g(x/y) (\partial y^r/\partial u) f_n(x, y), \quad (4.17)$$

$$g(x/y) = (n - r + 1) + 2 \sum_{k=1}^r (x/y)^k, \quad (4.18)$$

where $(\partial/\partial u)$ is for fixed v . Since $x/y = \exp(2v - y^r)$, we have

$$\begin{aligned} \log f_n(x, y) + k(v) &= \int g(\exp[2v - y^r]) d(y^r) \\ &= (n - r + 1) y^r - 2 \sum_{k=1}^r k^{-1} (x/y)^k, \end{aligned} \tag{4.19}$$

where the unknown function $k(v)$ can be determined from (4.15) by taking the limit $y \rightarrow 0$ with x/y fixed at e^{2e} :

$$k(\varrho) = -2 \sum_{k=1}^r k^{-1} e^{2ke}. \tag{4.20}$$

Therefore we have (4.8). By definition, $F_n(x) = f_n(x^{1/r}, x^{1/r})$ and we have (4.11).

To obtain (4.12), we consider

$$\hat{f}_n(x, y) = \exp[(n - r + 1) y^r + 2(x/y)^r \sum_{k=1}^r k^{-1} \{\exp k y^r - 1\}] \tag{4.21}$$

$$= \sum_{m,l} \hat{C}_l^m(n) x^{l-n} y^{mr+n-1}/m! \tag{4.22}$$

By the power series expansion of exponentials in (4.11) as well as in (4.21), we obtain expressions for $C_l^m(n)$ and $\hat{C}_l^m(n)$ as sums of positive terms. The change of $(x/y)^k$ in (4.11) to $(x/y)^r = (x/y)^k (x/y)^{r-k}$ increases the power of x by $r - k \geq 0$ while keeping the total degree in x and y as well as the numerical coefficient of each term. Therefore all terms in the expression for $C_l^m(n)$ moves into expressions for $\hat{C}_l^m(n)$ with the same m and higher $l \geq l$. Hence

$$\begin{aligned} 0 &\leq \sum_{l>L} C_l^m(n) \leq \sum_{l>L} \hat{C}_l^m(n) \\ 0 &\leq F_n^L(x) \leq \hat{F}_n^L(x) \equiv \sum_n \sum_{l>L} \hat{C}_l^m(n) x^m/m!, \quad (x > 0). \end{aligned} \tag{4.23}$$

By Taylor's expansion theorem, we have

$$\begin{aligned} \hat{F}_n^{L+r+n}(x) &= \left\{ \varphi_n(x, x) - \sum_{k=0}^L (x^k/k!) (\partial/\partial \xi)^k \varphi_n(\xi, x) \right\}_{\xi=0} \\ &= (L + 1)!^{-1} x^{L+1} (\partial/\partial \xi)^{L+1} \varphi_n(\xi, x)|_{\xi=\theta x} \end{aligned} \tag{4.24}$$

where $0 < \theta < 1$ and

$$\begin{aligned} \varphi_n(\xi, \eta) &= \hat{f}_n(\xi^{1/r}, \eta^{1/r}) \\ &= \exp \left[(n - r + 1) \eta + 2 \sum_{k=1}^r k^{-1} (e^{k\eta} - 1) (\xi/\eta) \right]. \end{aligned} \tag{4.25}$$

The main point of introducing \hat{F}_n is that (4.24) is easier to calculate for \hat{F} than for F . From (4.23), (4.24) and (4.25), we obtain (4.12).

Theorem 4.2. (i) $\{\exp \delta(\beta H(I))\} Q$ converges absolutely in the norm for any complex β ; $Q \in \mathfrak{A}_1$, and $I \subset Z$. (ii) $\exp \delta(\beta H(I))$ is a group of automorphisms of $\mathfrak{A}(I') \cap \mathfrak{A}_1$ and $\mathfrak{A}(I') \cap \mathfrak{A}_2$ with one complex parameter β for any $I' \supset I$. $\{\exp \delta(\beta H(I))\} Q$ is analytic in β for each $Q \in \mathfrak{A}_1$. (iii) $\tau_T^I(t) = \exp \delta(it H(I))$ for real t has a unique extension to a continuous one parameter group of *-automorphisms of \mathfrak{A} , commuting with $\tau_s(a)$. $\tau_T^Z = \tau_T$.

(iv) If $Q \in \mathfrak{A}([0, n])$, then for any $N \geq 0$

$$\{\exp \delta(\beta H(I))\} Q = Q_{N,I}(\beta) + \delta Q_{N,I}(\beta), \tag{4.26}$$

$$Q_{N,I}(\beta) \equiv \exp \delta[\beta H(I \cap [-N, n+N])] Q \in \mathfrak{A}([-N, n+N]), \tag{4.27}$$

$$\|\delta Q_{N,I}(\beta)\| \leq (1 + [N/r])!^{-1} \alpha(\beta)^{[N/r]+1} F_n(2|\beta| \|\Phi\|) \|Q\|, \tag{4.28}$$

$$\alpha(\beta) = 2 \sum_{k=1}^r k^{-1} (\exp 2k|\beta| \|\Phi\| - 1), \tag{4.29}$$

where $[N/r]$ denotes the largest integer not exceeding N/r , and $n \geq r - 1$.

Proof. By definition, if $Q \in \mathfrak{A}([0, n])$

$$\delta(\beta H(Z))^m Q = \beta^m \Sigma [\tau_s(j_m) \Phi, [\dots, [\tau_s(j_1) \Phi, Q] \dots]], \tag{4.30}$$

where the sum is over all $j_1 \dots j_m \in Z$ such that $[j_k, j_k + r]$ has a non empty intersection with the interval

$$I(j_1 \dots j_{k-1}) = [0, n] \cup \left(\bigcup_{l < k} [j_l, j_l + r] \right) \tag{4.31}$$

for each $k = 1, 2, \dots m$. Let $C_l^m(n)$ be the number of terms in (4.30) for which the length of the interval $I(j_1 \dots j_m)$ is l . It satisfies (4.4) and (4.5), where the first term of (4.4) represents the case in which $[j_{m+1}, j_{m+1} + r]$ falls in $I(j_1 \dots j_m)$ and the rest represents cases in which $[j_{m+1}, j_{m+1} + r]$ has still non empty intersection with $I(j_1 \dots j_m)$ and sticks out to either side of $I(j_1 \dots j_m)$. We now have two inequalities

$$\|\delta(\beta H(I))^m Q\| \leq (2|\beta| \|\Phi\|)^m \|Q\| \sum_l C_l^m(n), \tag{4.32}$$

$$\begin{aligned} \|\delta(\beta H(I))^m Q - \delta(\beta H(I \cap [-N, n+N]))^m Q\| \\ \leq (2|\beta| \|\Phi\|)^m \|Q\| \sum_{l > N+n} C_l^m(n). \end{aligned} \tag{4.33}$$

Note that the change of Z to $I \subset Z$ only decreases the number of terms in (4.30).

We are now ready to prove (i) and (iv). If we write $Q = \sum_{k=k_0}^{\infty} Q_k$, $Q_k = Q^{(k)} - Q^{(k-1)} \in \mathfrak{A}([-k, k])$, $\|Q - Q^{(k)}\| = \|Q\|_k (k \geq k_0)$, $Q^{(k_0-1)} = 0$, we have $\|Q_k\| \leq \|Q\|_k + \|Q\|_{k-1}$ where $\|Q\|_{k_0-1}$ is to be replaced by $\|Q\|$.

Hence if $Q \in \mathfrak{A}_1$, then $\Sigma x^k \|Q_k\| < \infty$ for any $x > 0$. From our discussion, we have

$$\begin{aligned} & \sum_{m=0}^{\infty} (m!)^{-1} |\beta|^m \sum_{j_1 \dots j_m} \| [\tau_s(j_m) \Phi, [\dots, [\tau_s(j_1) \Phi, Q] \dots]] \| \\ & \leq \sum_{k=k_0}^{\infty} \sum_{m=0}^{\infty} (m!)^{-1} (2|\beta| \|\Phi\|)^m \sum_I C_I^m(2k) (\|Q\|_k + \|Q\|_{k-1}) \quad (4.34) \\ & \leq \sum_{k=k_0}^{\infty} F_{2k}(2|\beta| \|\Phi\|) (\|Q\|_k + \|Q\|_{k-1}). \end{aligned}$$

By Lemma 4.1, $F_{2k}(x) = (e^{2x})^k F_0(x)$. Hence (4.34) is finite and we have (i). Similarly, for $Q \in \mathfrak{A}[0, n]$,

$$\begin{aligned} \|\delta Q_{N,I}(\beta)\| & \leq \sum_{m=0}^{\infty} (m!)^{-1} (2|\beta| \|\Phi\|)^m \sum_{l > N+n} C_I^m(n) \|Q\| \\ & \leq F_n^{N+n}(2|\beta| \|\Phi\|) \|Q\| \quad (4.35) \\ & \leq F_n^{[N/r]r+n}(2|\beta| \|\Phi\|) \|Q\|. \end{aligned}$$

By substituting (4.12) into (4.35), we obtain (4.28).

To obtain (ii), we first show that \mathfrak{A}_1 and \mathfrak{A}_2 are mapped into themselves.

Let $Q = \Sigma Q_k$, $Q_k \in \mathfrak{A}([-k, k])$, $\|Q_k\| \leq \|Q\|_k + \|Q\|_{k-1}$, $k = k_0, k_0 + 1, \dots$ where $\|Q\|_{k_0-1}$ is to be replaced by $\|Q\|$. We have for $n \geq k_0$

$$\begin{aligned} \gamma_n & = \|\exp \delta(\beta H(I)) Q\|_n \\ & \leq \sum_{k=k_0}^n F_{2k}^{(n-k)+2k}(2|\beta| \|\Phi\|) \|Q_k\| + \sum_{k=n+1}^{\infty} F_{2k}(2|\beta| \|\Phi\|) \|Q_k\|. \quad (4.36) \end{aligned}$$

Hence, for $x > 1$,

$$\begin{aligned} & \sum_{n=k_0}^{\infty} x^n \gamma_n \\ & \leq \left(\sum_{N=0}^{\infty} x^N (1 + [N/r])!^{-1} \alpha(\beta)^{[N/r]+1} \right) \left(\sum_{k=k_0}^{\infty} (x e^{4\|\beta\|\Phi\|})^k \|Q_k\| \right) F_0(2\|\beta\|\Phi) \\ & \quad + \sum_{k > k_0} \{x e^{4\|\beta\|\Phi\|}\}^k \|Q_k\| (x-1)^{-1} F_0(2\|\beta\|\Phi), \quad (4.37) \end{aligned}$$

which is finite if $Q \in \mathfrak{A}_1$. Next, if $Q \in \mathfrak{A}_2$, we have

$$\begin{aligned} \log \gamma_n & \leq \log \sum_{k=k_0}^{n+1} \gamma_{nk} \leq \log \{n+2-k_0\} \max \gamma_{nk} \\ & = \log(n+2-k_0) + \max \log \gamma_{nk} \quad (4.38) \end{aligned}$$

where $\gamma_{nk} = \delta_{n-k}^1 \delta_k^2$ for $k = k_0 \dots n$,

$$\delta_N^1 = (1 + [N/r])!^{-1} \alpha(\beta)^{[N/r]+1}, \tag{4.39}$$

$$\delta_k^2 = F_{2k}(2\|\beta\Phi\|) \|Q_k\|, \tag{4.40}$$

$$\gamma_{n,n+1} = \sum_{k=n+1}^{\infty} F_{2k}(2\|\beta\Phi\|) \|Q_k\| \leq (1 - e^{-1})^{-1} F_{2n}(2\|\beta\Phi\|) \gamma'_{n,n+1}, \tag{4.41}$$

$$\gamma'_{n,n+1} = \sup_{k \geq 1} [e^{(1+4\|\beta\Phi\|)}]^k \|Q_{n+k}\|. \tag{4.42}$$

We have, for $k_0 \leq k \leq n$ and $n \geq k_0$,

$$n^{-1} [(n/r) \log n - (k/r) \log k - [(n-k)/r] \log(n-k)] \leq r^{-1} \log 2, \tag{4.43}$$

$$n^{-1} \log F_{2k}(2\|\beta\Phi\|) \leq 4\|\beta\Phi\| + \log F_0(2\|\beta\Phi\|), \tag{4.44}$$

$$\begin{aligned} & n^{-1} \{ \log \delta_{n-k}^1 + ((n-k)/r) \log(n-k) \} \\ & \leq \sup_{N \geq 1} N^{-1} \{ (N/r) \log N - \log \Gamma(N/r + 1) \} + (1 + 1/r) \log \alpha(\beta) < \infty, \end{aligned} \tag{4.45}$$

$$n^{-1} \{ \log \|Q_k\| + (k/r) \log k \} \leq \sup_{l \geq 1} l^{-1} \{ \log \|Q_l\| + (l/r) \log l \} < \infty, \tag{4.46}$$

$$\begin{aligned} & n^{-1} \{ \log \gamma'_{n,n+1} + (n/r) \log n \} \\ & \leq \sup_{k,n \geq 1} \{ (1 + (k/n)) \sup_l \{ l^{-1} (\log \|Q_l\| + (l/r) \log l) \} \\ & \quad - r^{-1} \log(n+k) \} + r^{-1} \log n + (k/n) (1 + 4\|\beta\Phi\|) < \infty. \end{aligned} \tag{4.47}$$

Therefore $\sup_n n^{-1} (\gamma_n + (n/r) \log n) < \infty$.

The isomorphism property of $\exp \delta(\beta H(I))$ follows from the Leibnitz formula:

$$\delta(\beta H(I))^m (Q_1 Q_2) = \sum_{k=0}^m \binom{m}{k} (\delta(\beta H(I))^k Q_1) (\delta(\beta H(I))^{m-k} Q_2). \tag{4.48}$$

To see the group property, we first note that

$$\sum_{m,n} \|\delta(\beta_1 H(I))^m \delta(\beta_2 H(I))^n Q / m! n!\| < \infty.$$

Hence we can change the order of summation to obtain

$$\exp \delta(\beta_1 H(I)) \exp \delta(\beta_2 H(I)) = \exp \delta((\beta_1 + \beta_2) H(I)).$$

Since $\exp \delta(\beta H(I)) Q$ has a power series expansion in β which converges absolutely for all β , it is analytic in β . This completes the proof of (ii).

The *-isomorphism property of $\tau_T^I(t)$ follows from

$$[\delta(it H(I)) Q]^* = \delta(it H(I)) Q. \tag{4.49}$$

Any $*$ -isomorphism of \mathfrak{A}_0 into \mathfrak{A} can be uniquely extended to a $*$ -isomorphism of \mathfrak{A} into \mathfrak{A} . The uniqueness guarantees that the extension to \mathfrak{A} of the restriction to \mathfrak{A}_0 of $\exp \delta(it H(I))$ is the extension of $\exp \delta(it H(I))$. The group property and the continuity in β is preserved in the extension. Hence each $\exp \delta(it H(I))$ has an inverse $\exp \delta(-it H(I))$ and therefore must be a $*$ -automorphism. The commutativity with $\tau_s(a)$ is immediate. This proves (iii). Q.E.D.

Corollary 4.3. *If $|t| \leq (2r \|\Phi\|)^{-1} \log |a|$, then*

$$\lim_{a \rightarrow \infty} e^{|a|^e} \|[Q_1, \tau_s(a) \tau_T(t) Q_2]\| = 0 \tag{4.50}$$

for any $Q_1, Q_2 \in \mathfrak{A}_0$ and $Q > 0$.

The proof is immediate from Theorem 4.2 (iv).

Remark 4.4. The convergence of $\exp \delta(\beta H(Z^v)) Q$ for v dimensional lattice has been proved for $|\beta| < [2r(r-1) \|\Phi\|]^{-1}$. ([11, 13].) In this case, a weaker commutativity can be proven in a region where $|t|$ can grow to infinity as $|a| \rightarrow \infty$.

§ 5. Expansions

Definition 5.1. (cf. [5]). *Let $Q \in \mathfrak{A}_0$, $Q(\beta; I) = \exp \delta(\beta H(I)) Q$. Then*

$$E_r(Q; H(I)) = \sum_{n=0}^{\infty} \int_0^1 d\beta_1 \int_0^{\beta_1} d\beta_2 \dots \int_0^{\beta_{n-1}} d\beta_n \prod_j^{n \rightarrow 1} Q(\beta_j; I), \tag{5.1}$$

$$E_l(Q; H(I)) = \sum_{n=0}^{\infty} \int_0^1 d\beta_1 \int_0^{\beta_1} d\beta_2 \dots \int_0^{\beta_{n-1}} d\beta_n \prod_j^{1 \rightarrow n} Q(-\beta_j; I), \tag{5.2}$$

where

$$\prod_j^{1 \rightarrow n} A_j = A_1 \dots A_n, \quad \prod_j^{n \rightarrow 1} A_j = A_n \dots A_1. \tag{5.3}$$

By a change of integration variables, we also have, for real β ,

$$E_r(\beta Q; \beta H(I)) = \sum_{n=0}^{\infty} \int_0^{\beta} d\beta_1 \int_0^{\beta_1} d\beta_2 \dots \int_0^{\beta_{n-1}} d\beta_n \prod_j^{n \rightarrow 1} Q(\beta_j; I), \tag{5.4}$$

$$E_l(\beta Q; \beta H(I)) = \sum_{n=0}^{\infty} \int_0^{\beta} d\beta_1 \int_0^{\beta_1} d\beta_2 \dots \int_0^{\beta_{n-1}} d\beta_n \prod_j^{1 \rightarrow n} Q(-\beta_j; I). \tag{5.5}$$

For $Q \in \mathfrak{A}([0, n])$, the sums and integrals are absolutely and uniformly (for bounded $\|\beta \Phi\|, \|Q\|$ and n) convergent because $\|Q(\beta_j; I)\|$

$\leq F_n(2 \|\beta \Phi\|) \|Q\|$ for $|\beta_j| \leq |\beta|$ and hence

$$\sum_{n=0}^{\infty} \int_0^{\beta} d\beta_1 \int_0^{\beta_1} d\beta_2 \dots \int_0^{\beta_{n-1}} d\beta_n \prod_{j=1}^n \|Q(\beta_j; I)\| \leq \exp \{|\beta| F_n(2 \|\beta \Phi\|) \|Q\|\} < \infty. \tag{5.6}$$

For bounded operators Q and R ,

$$\{\exp \delta(\beta R)\} Q = e^{\beta R} Q e^{-\beta R}, \tag{5.7}$$

$$E_r(\beta Q; \beta R) \{\exp \beta R\} = \exp \beta(Q + R), \tag{5.8}$$

$$\{\exp \beta R\} E_l(\beta Q; \beta R) = \exp \beta(Q + R). \tag{5.9}$$

These formulas can be easily proved by noting that they are 1 for $\beta = 0$ and each side of 3 equations satisfy differential equations $(d/d\beta) S = [R, S]$; $(d/d\beta) S = S(Q + R)$, and $(d/d\beta) S = (Q + R) S$, respectively. ((5.7) is used in the proof of (5.8) and (5.9).)

From (5.8) and (5.9), the following formulas follow immediately.

$$E_r(Q_1 + Q_2; R) = E_r(Q_1; Q_2 + R) E_r(Q_2; R), \tag{5.10}$$

$$E_r(Q; R) = E_l(Q; -R - Q), \quad E_l(Q; R) = E_r(Q; -Q - R), \tag{5.11}$$

$$E_r(Q; R) E_l(-Q; -R) = E_l(-Q; -R) E_r(Q; R) = 1, \tag{5.12}$$

$$E_r(Q; R) \{\exp \delta(R)\} Q' = [\{\exp \delta(Q + R)\} Q'] E_r(Q; R), \tag{5.13}$$

$$[\{\exp \delta(-R)\} Q'] E_l(Q; R) = E_l(Q; R) \{\exp \delta(-R - Q)\} Q'. \tag{5.14}$$

Lemma 5.2. (i) *If $Q \in \mathfrak{A}([0, n])$, then*

$$E_r(Q; \lambda H(I)) = E_r[Q; \lambda H(I \cap [-N, n + N])] + \delta_r(I, N), \tag{5.15}$$

$$\|\delta_r(I, N)\| \leq C_n \delta_N(\lambda \Phi), \tag{5.16}$$

$$\delta_N(\lambda \Phi) = (1 + [N/r])!^{-1} \alpha(\lambda)^{[N/r]+1}, \tag{5.17}$$

where C_n depends on n , $\|\lambda \Phi\|$ and $\|Q\|$ but is independent of N and I .

(ii) *The same equation holds when the suffix r is replaced by l .*

(iii) *If $Q \in \mathfrak{A}_0$, then $E_r(Q; \lambda H(I)) \in \mathfrak{A}_2$, $E_l(Q; \lambda H(I)) \in \mathfrak{A}_2$.*

(iv) *Formulas (5.10) ~ (5.14) hold when R is replaced by $\lambda H(I)$ and if $Q_1, Q_2, Q, Q' \in \mathfrak{A}_0$.*

Proof. We have $\|Q(\beta'; I)\| \leq F_n(2 \|\lambda \Phi\|) \|Q\|$ and $\|Q_{N,I}(\beta')\| \leq F_n(2 \|\lambda \Phi\|) \|Q\|$ for $|\beta'| \leq 1$. Since

$$\prod_j^{n \rightarrow 1} Q(\beta_j; I) = \sum_{j=1}^n \left(\prod_k^{n \rightarrow j+1} Q(\beta_k; I) \right) \delta_{Q_{N,I}}(\beta_j) \left(\prod_k^{j-1 \rightarrow 1} Q_{N,I}(\beta_k) \right) + \prod_j^{n \rightarrow 1} Q_{N,I}(\beta_j), \tag{5.18}$$

we have (5.16) where, for $n \geq r - 1$,

$$C_n = F_n(2 \|\lambda \Phi\|) \|Q\| \exp \{F_n(2 \|\lambda \Phi\|) \|Q\|\}. \tag{5.19}$$

(ii) follows in exactly the same manner. By the Sterling formula, we obtain (iii). The formulas (5.10) ~ (5.14) hold when R is replaced by $\lambda H(I \cap [-N, n + N])$. By taking the limit $N \rightarrow \infty$, we obtain (5.10) ~ (5.14) for $R = \lambda H(I)$. Q.E.D.

§ 6. The Mapping \mathcal{L}

Definition 6.1. Let I be a finite subset of Z , φ be a state of $\mathfrak{A}(I)$, $Q = \sum u_i Q_i$, $u_i \in \mathfrak{A}(I)$, $Q_i \in \mathfrak{A}(Z \setminus I)$. Then $\varphi(Q) \equiv \sum \varphi(u_i) Q_i \in \mathfrak{A}(Z \setminus I) \subset \mathfrak{A}$.

It is easily proved that $\varphi(Q)$ does not depend on a particular decomposition $Q = \sum u_i Q_i$.

Definition 6.2. Let $Q \in \mathfrak{A}$.

$$\mathcal{L}(Q) = \tau_c(1) d^{-2} \text{tr}_{[0,1]}(K^* Q K), \tag{6.1}$$

$$K = K_+ K_-, \tag{6.2}$$

$$K_+ = E_r[-(1/2) \tau_s(1) \Phi; -(1/2) H(2, \infty)], \tag{6.3}$$

$$K_- = E_r[-(1/2) \tau_s(-r) \Phi; -(1/2) H(-\infty, -1)]. \tag{6.4}$$

Lemma 6.3. (i) If $Q \in \mathfrak{A}_1$, then $\mathcal{L}Q \in \mathfrak{A}_1$. If $Q \in \mathfrak{A}_2$, then $\mathcal{L}Q \in \mathfrak{A}_2$. (ii) If $Q \geq 0$, then $\mathcal{L}Q \geq 0$. If $Q \geq 0$ and $Q \neq 0$, then $\mathcal{L}Q \neq 0$. If $Q \geq 0$ and Q^{-1} exists, then $(\mathcal{L}Q)^{-1} \in \mathfrak{A}$. (iii) If $n > r$, then

$$\mathcal{L}^n Q = p_n \tau_c(n) \varphi_n(K_n^* Q K_n), \tag{6.5}$$

$$K_n = K_{n+} K_{n-}, K_{n+} \in \mathfrak{A}([1, \infty)), K_{n-} \in \mathfrak{A}((-\infty, 0]) \tag{6.6}$$

$$K_{n+} = E_r(-(1/2) \tau_s(n) \Psi; -(1/2) \{H(1, n) + H(n + 1, \infty)\}), \tag{6.7}$$

$$K_{n-} = E_r(-(1/2) \tau_s(-n) \Psi; -(1/2) \{H(-\infty, -n) + H(1 - n, 0)\}), \tag{6.8}$$

$$\Psi = \sum_{j=1-r}^0 \tau_s(j) \Phi \in \mathfrak{A}([1 - r, r]), \tag{6.9}$$

$$d^{2n} \varphi_n(Q') = p_n^{-1} \text{tr}_{[1-n,n]}(Q' \exp - \{H(1, n) + H(1 - n, 0)\}), \tag{6.10}$$

$$d^{2n} p_n = \text{tr}_{[1-n,n]}(\exp - \{H(1, n) + H(1 - n, 0)\}). \tag{6.11}$$

Proof. Since $K_{\pm} \in \mathfrak{A}_2$ by (5.2) (iii), $K^* Q K \in \mathfrak{A}_1$ or \mathfrak{A}_2 according as $Q \in \mathfrak{A}_1$ or \mathfrak{A}_2 . Since $\|\mathcal{L}(Q)\|_n \leq \|K^* Q K\|_{n+1}$, $\mathcal{L}(Q) \in \mathfrak{A}_1$ or \mathfrak{A}_2 . If $Q \geq 0$, then $K^* Q K \geq 0$ and hence $\varphi(\mathcal{L}Q) = (\tau_c(1)^* \varphi) \otimes (d^{-2} \text{tr}_{[0,1]}(K^* Q K)) \geq 0$, for any state φ of \mathfrak{A} , where $\tau_c(1)^* \varphi$ is the state of $\mathfrak{A}(Z \setminus [0, 1])$ such that $\tau_c(1)^* \varphi(Q'') = \varphi(\tau_c(1) Q'')$ for all $Q'' \in \mathfrak{A}(Z \setminus [0, 1])$. Hence $\mathcal{L}Q \geq 0$. If

$\mathcal{L}Q = 0$ in addition, then $\varphi_c(\mathcal{L}Q) = \varphi_c(K^*QK) = 0$ for the central state φ_c of \mathfrak{A} . Since φ_c is faithful on the non negative elements of \mathfrak{A} , we have $K^*QK = 0$. Since K_{\pm} has the inverses

$$K_+^{-1} = E_l(\tau_s(1) \Phi/2; H(2, \infty)/2), \tag{6.12}$$

$$K_-^{-1} = E_l(\tau_s(-r) \Phi/2; H(-\infty, -1)/2), \tag{6.13}$$

we have $Q = 0$. If Q^{-1} exists, then $K^*QK \geq \|Q^{-1}\|^{-1} \|K^{-1}\|^{-2}$ and hence $\mathcal{L}(Q) \geq \|Q^{-1}\|^{-1} \|K^{-1}\|^{-2}$. Therefore $\mathcal{L}(Q)^{-1} \in \mathfrak{A}$.

To obtain (iii), we use (5.10):

$$\begin{aligned} \prod_j^{1 \rightarrow N} \tau_s(j-1) K_+ &= E_r[-(1/2)H(1, N+r); -(1/2)H(N+1, \infty)] \\ &= K_{n+} E_r[-(1/2)H(1, N); -(1/2)H(N+1, \infty)]. \end{aligned} \tag{6.14}$$

Since $H(1, N)$ commutes with every $\tau_s(j) \Phi$ in $H(N+1, \infty)$, the second factor of (6.14) is $\exp -(1/2) H(1, N)$. Similar equation holds for K_- . Therefore

$$\mathcal{L}^n Q = \tau_c(n) d^{-2n} \text{tr}_{[1-n, n]} \{(K'_{n+} K'_{n-})^* Q (K'_{n+} K'_{n-})\}, \tag{6.15}$$

$$K'_{n+} = \prod_j^{1 \rightarrow N} \tau_s(j-1) K_+, \tag{6.16}$$

$$K'_{n-} = \prod_j^{1 \rightarrow N} \tau_s(1-j) K_-. \tag{6.17}$$

This proves (iii).

Lemma 6.4. (i) Let $Q \in \mathfrak{A}$, $Q > 0$, $\alpha(Q) < \infty$. Then

$$\alpha(\mathcal{L}^n Q) \leq \alpha(Q) b(\|\Phi\|), \tag{6.18}$$

$$\alpha_l(\mathcal{L}^n Q) \leq \alpha(Q) b'(\|\Phi\|) \delta_{l-r}(\|\Phi\|/2) + b(\|\Phi\|) \alpha_{l+n}(Q), \tag{6.19}$$

where $\delta_l(\|\Phi\|/2)$ is defined in (5.17) and $b'(\|\Phi\|)$ is another constant.

(ii) Let $\|Q\|_{M,x} \|Q^{-1}\| \leq a$, $0 < a$, $x > 1$. Then there exists $N(a, M, x, \|\Phi\|)$ such that

$$\alpha(\mathcal{L}^n Q) \leq 3b(\|\Phi\|) \tag{6.20}$$

for any $n \geq N(a, M, x, \|\Phi\|)$ uniformly in Q .

Proof. From (5.12) and estimates in Theorem 4.2 (i), we have

$$\|S\| \leq \exp\{F_{2r-1}(\|\Phi\|) \|\Psi\|/2\} \leq \exp\{(r/2) F_{2r-1}(\|\Phi\|) \|\Phi\|\} \tag{6.21}$$

$$\|S^{-1}\| \leq \exp\{(r/2) F_{2r-1}(\|\Phi\|) \|\Phi\|\} \tag{6.22}$$

for $S = K_{n+}, K_{n-}, K_{n+}^{NN'}, K_{n-}^{NN'}$ where $r \leq N < n$, $N' \geq r$, N' may become $+\infty$ and

$$K_{n+}^{NN'} = E_r(-\tau_s(n) \Psi/2; \\ - \{H([n-N, n]) + H([n+1, n+1+N'])\}/2), \quad (6.23)$$

$$K_{n-}^{NN'} = E_r(-\tau_s(-n) \Psi/2; \\ - \{H([-n-N', -n]) + H([1-n, 1-n+N])\}/2). \quad (6.24)$$

Hence

$$\alpha(K_n^* Q K_n) \leq \alpha(Q) \| (K_n^*)^{-1} \| \| K_n^{-1} \| \| K_n \| \| K_n^* \| \leq \alpha(Q) b(\|\Phi\|), \quad (6.25)$$

where

$$b(\|\Phi\|) = \exp \{4r F_{2r-1}(\|\Phi\|) \|\Phi\|\}. \quad (6.26)$$

Since $b_2 \geq Q'' \geq b_1$ implies $b_2 \geq \varphi_n(Q'') \geq b_1$, and since $\alpha(p_n \tau_c(n) Q''') = \alpha(Q''')$, we have (6.18) for $n > r$.

If $n \leq r$, $H(1, n)$ in (6.7) and $H(1-n, 0)$ in (6.8) are absent and Ψ is replaced by $\Psi_n \equiv \sum_{j=1-n}^0 \tau_s(j) \Phi$, which satisfies $\|\Psi_n\| \leq r \|\Phi\|$. Hence we have the same result.

Since $\alpha(R) \geq \alpha_l(R)$ for any R by (3.16), the Eq. (6.19) for $l \leq r$ follows from (6.18). The modification for the case $n \leq r$ is the same as above. Hence we consider the case $n > r$, $l > r$.

Now we prove (6.19). Let $Q_{(n+l)}$ be such that

$$Q_{(n+l)} \in \mathfrak{A}([-n-l, n+l]), \|Q - Q_{(n+l)}\| \|Q_{(n+l)}^{-1}\| = \alpha_{n+l}(Q).$$

Let

$$K_{(n,l)} = K_{n+}^{n-1, l-1} K_{n-}^{n-1, l-1}.$$

Let

$$Q'_i = \tau_c(n) \varphi_n(K_{(n,l)}^* Q_{(n+l)} K_{(n,l)}) \in \mathfrak{A}([-l, l])$$

and compute $\|Q' - Q'_i\| \|(Q'_i)^{-1}\|$ where $Q' = \tau_c(n) \varphi_n(K_n^* Q K_n)$. From Lemma 5.2 (i) and (6.21), we have

$$\|K_n^* Q K_n - K_{(n,l)}^* Q_{(n+l)} K_{(n,l)}\| \\ \leq 4 \|Q\| b(\|\Phi\|)^{1/2} \{(r/2) \|\Phi\| F_{2r-1}(\|\Phi\|)\} \delta_{l-r}(\|\Phi\|/2), \quad (6.27)$$

$$\|K_{(n,l)}^* (Q - Q_{(n+l)}) K_{(n,l)}\| \leq b(\|\Phi\|)^{1/2} \|Q - Q_{(n+l)}\|. \quad (6.28)$$

Hence we have

$$\|Q' - Q'_i\| \leq (2b(\|\Phi\|)^{1/2})^{-1} b'(\|\Phi\|) \delta_{l-r}(\|\Phi\|/2) \\ + b(\|\Phi\|)^{1/2} \|Q - Q_{(n+l)}\|, \quad (6.29)$$

where

$$b'(\|\Phi\|) = b(\|\Phi\|) \{4r\|\Phi\| F_{2r-1}(\|\Phi\|)\}. \quad (6.30)$$

On the other hand,

$$\begin{aligned} K_{(n,l)}^* Q_{(n+l)} K_{(n,l)} &\geq \{K_{(n,l)}^* Q_{(n+l)} K_{(n,l)}\}^{-1} \|\cdot\|^{-1} \\ &\geq b(\|\Phi\|)^{-1/2} \|Q_{(n+l)}^{-1}\|^{-1}. \end{aligned} \quad (6.31)$$

By Lemma 3.9 (3) and (1), we may assume

$$\|Q_{(n+l)}^{-1}\| \leq \|Q^{-1}\| (1 + \alpha_l(Q)) \leq 2\|Q^{-1}\|. \quad (6.32)$$

Hence

$$Q'_l \geq b(\|\Phi\|)^{-1/2} \|Q_{(n+l)}^{-1}\|^{-1} \geq 2^{-1} b(\|\Phi\|)^{-1/2} \|Q^{-1}\|^{-1}. \quad (6.33)$$

Therefore

$$\begin{aligned} \alpha_l(Q') &\leq \|Q' - Q'_l\| \|Q'_l\|^{-1} \\ &\leq b(\|\Phi\|) \alpha_{n+l}(Q) + b'(\|\Phi\|) \alpha(Q) \delta_{l-r}(\|\Phi\|/2). \end{aligned} \quad (6.34)$$

Since $\alpha_l(p_n Q') = \alpha_l(Q')$, we have (6.19).

We now prove (ii). In the previous computation, we consider $Q_{(k)}$ instead of $Q_{(n+l)}$ and $K'_n = K_{n+}^{N,\infty} K_{n-}^{N,\infty}$ instead of $K_{(n,l)}$. We then have bound (6.29) and (6.31) for $K_n^* Q K_n - (K'_n)^* Q_{(k)} K'_n$ and $(K'_n)^* Q_{(k)} K'_n$ where $l-r$ is to be replaced by $N-r+1$ and $(n+l)$ by (k) . Hence for any state φ' and φ'' of \mathfrak{A} , we have

$$\varphi'(\tau_c(n) \varphi_n(K_n^* Q K_n)) \leq \varphi'[\tau_c(n) \varphi_n((K'_n)^* Q_{(k)} K'_n)] (1 + \Delta). \quad (6.35)$$

$$\varphi''(\tau_c(n) \varphi_n(K_n^* Q K_n)) \geq \varphi''[\tau_c(n) \varphi_n((K'_n)^* Q_{(k)} K'_n)] (1 - \Delta), \quad (6.36)$$

$$\Delta \leq b(\|\Phi\|) \alpha_k(Q) + b'(\|\Phi\|) \alpha(Q) \delta_{N-r+1}(\|\Phi\|/2). \quad (6.37)$$

From $\|Q\|_{M,x} \|Q^{-1}\| \leq a$, we have $\|Q^{-1}\| \|Q\|_k \leq x^{-k} a$ for $k \geq M$. By Lemma 3.9 (6), we have

$$\alpha_k(Q) \leq x^{-k} a (1 - x^{-k} a)^{-1}. \quad (6.38)$$

Let L be an integer such that $L \geq M$,

$$x^L \geq (1 + 4b(\|\Phi\|)) a. \quad (6.39)$$

For $k \geq L$, we have $\alpha_k(Q) \leq \{4b(\|\Phi\|)\}^{-1}$. Since $\|Q\| \leq \|Q\|_{M,x}$ we have $\alpha(Q) \leq a$. Let N be an integer such that $N > r$ and

$$4b'(\|\Phi\|) a \delta_{N-r+1}(\|\Phi\|/2) \leq 1. \quad (6.40)$$

We then have $\Delta \leq 1/2$ and hence by (6.5) and (3.13) we have

$$\alpha(\mathcal{L}^n Q) \leq 3 \sup_{\varphi', \varphi''} \varphi' [\tau_c(n) \varphi_n((K'_n)^* Q_{(k)} K'_n)] / \varphi'' [\tau_c(n) \varphi_n((K'_n)^* Q_{(k)} K'_n)]. \quad (6.41)$$

Now we set $N(a, M, x, \|\Phi\|) = L + N + 2$. For $n \geq N(a, M, x, \|\Phi\|)$, we have $Q_k \in \mathfrak{A}([-k, k])$, $K'_n \in \mathfrak{A}(Z \setminus [-k, k])$. Let

$$\bar{\varphi}_1(Q'') \equiv \varphi_1[\tau_c(n) \varphi_n(Q_{(k)} Q'')] = (\varphi_n \otimes \tau_c(n)^* \varphi_1)(Q_{(k)} Q'')$$

be a state on $\mathfrak{A}(Z \setminus [-k, k])$ ($\ni Q''$) induced by $\varphi_1 = \varphi'$ and φ'' . Then we have

$$\alpha(\mathcal{L}^n Q) \leq 3 \sup_{\varphi', \varphi''} \bar{\varphi}'((K'_n)^* K'_n) / \bar{\varphi}''((K'_n)^* K'_n) \leq 3b(\|\Phi\|). \quad (6.42)$$

Lemma 6.5. \mathcal{L} maps $\mathfrak{A}(M, x)$ continuously into itself where $x > 1$.

Proof. By Lemma 6.3 (i), $\mathcal{L}1 \in \mathfrak{A}_1 \subset \mathfrak{A}(M, x)$. Now consider Q such that $\|Q\|_{M, x} \leq 1$, $Q = Q^*$. Let $Q' = 2 + Q$. Then $3 \geq Q' \geq 1$ and hence $\alpha(Q') \leq 3$. By (6.19),

$$\alpha_l(\mathcal{L}Q') \leq b(\|\Phi\|) \alpha_{l+1}(Q') + 3b'(\|\Phi\|) \delta_{l-r}(\|\Phi\|/2). \quad (6.43)$$

By $\|Q'\|_{M, x} \leq 1$, we have $\|Q'\|_n \leq x^{-n}$. By (3.21) we have

$$\alpha_{l+1}(Q') \leq x^{-(l+1)}(1 - x^{-(l+1)}), \quad (6.44)$$

where we have used $\|(Q')^{-1}\| \leq 1$. Let L be such that $y_L < 1$ where

$$y_l = b(\|\Phi\|) x^{-(l+1)}(1 - x^{-(l+1)}) + 3b'(\|\Phi\|) \delta_{l-r}(\|\Phi\|/2). \quad (6.45)$$

From (6.1), we have

$$\|\mathcal{L}Q'\| \leq \|K\|^2 \|Q'\| \leq 3b(\|\Phi\|)^{1/2}. \quad (6.46)$$

By (3.22), $\|R\| \geq \|R^{-1}\|^{-1}$ and (6.46),

$$\|\mathcal{L}Q'\|_l \leq 3b(\|\Phi\|)^{1/2} y_l (1 - y_l)^{-1}. \quad (6.47)$$

Thus

$$\|\mathcal{L}Q'\|_{M, x} \leq 3b(\|\Phi\|)^{1/2} \left\{ 1 + |L - M| + \sum_{l=L}^{\infty} x^l y_L (1 - y_L)^{-1} \right\} < \infty. \quad (6.48)$$

Therefore $\|\mathcal{L}Q\|_{M, x} \leq \|\mathcal{L}Q'\|_{M, x} + 2\|\mathcal{L}1\|_{M, x}$ is uniformly bounded.

Let $Q = Q_1 + iQ_2$, $Q_1^* = Q_1$, $Q_2^* = Q_2$. Then

$$\|Q_1\| = \sup_{\varphi} |\varphi(Q_1)| \leq \sup_{\varphi} |\varphi(Q_1) + i\varphi(Q_2)| \leq \|Q\|. \quad (6.49)$$

Similarly $\|Q_2\| \leq \|Q\|$. Further, let $Q^{(l)}$ be such that $\|Q - Q^{(l)}\| = \|Q\|_l$, $Q^{(l)} = Q_1^{(l)} + iQ_2^{(l)}$, $(Q_1^{(l)})^* = Q_1^{(l)}$, $(Q_2^{(l)})^* = Q_2^{(l)}$, $Q_1^{(l)} \in \mathfrak{A}([-l, l])$, $Q_2^{(l)} \in \mathfrak{A}([-l, l])$.

Then by the same argument as (6.49), we have

$$\|Q_1\|_l \leq \|Q - Q_1^{(l)}\| \leq \|Q - Q^{(l)}\| = \|Q\|_l. \quad (6.50)$$

Similarly $\|Q_2\|_l \leq \|Q\|_l$. Therefore $\|Q_1\|_{M,x} \leq \|Q\|_{M,x}$, $\|Q_2\|_{M,x} \leq \|Q\|_{M,x}$. By using the uniform boundedness for selfadjoint Q , we have the uniform boundedness of $\|\mathcal{L}Q\|_{M,x} \leq \|\mathcal{L}Q_1\|_{M,x} + \|\mathcal{L}Q_2\|_{M,x}$.

§ 7. Convergence Proof

Lemma 7.1. *There exists a state ν of \mathfrak{A} and $\lambda > 0$ such that $\nu(\mathcal{L}Q) = \lambda\nu(Q)$ for all $Q \in \mathfrak{A}$.*

Proof. Let $\hat{\mathcal{L}}$ be the mapping of states of \mathfrak{A} into themselves defined by

$$(\hat{\mathcal{L}}\varphi)(Q) = \varphi(\mathcal{L}1)^{-1}\varphi(\mathcal{L}Q). \quad (7.1)$$

Since

$$\varphi(\mathcal{L}1) \geq \|(\mathcal{L}1)^{-1}\|^{-1} \geq b(\|\Phi\|)^{-1/2} > 0, \quad (7.2)$$

$\hat{\mathcal{L}}$ is weakly continuous. The set of states is convex and weakly compact. Hence $\hat{\mathcal{L}}$ has a fixed point ν due to the Schauder-Tychonov theorem. It satisfies

$$\nu(\mathcal{L}Q) = \lambda\nu(Q), \quad \lambda = \nu(\mathcal{L}1) > 0. \quad (7.3)$$

Definition 7.2. *Let $Q \in \mathfrak{A}$, $LQ = \lambda^{-1}\mathcal{L}Q$. $\Sigma(Q)$ denotes the closure of the convex hull of $\{L^n Q; n = 0, 1, 2, \dots\}$.*

Lemma 7.3. *Let $Q > 0$, $\alpha(Q) < \infty$. $\Sigma(Q)$ is a compact subset of \mathfrak{A} , convex and invariant under L . L is continuous on \mathfrak{A} . If $Q \in \mathfrak{A}_2$, then $\Sigma(Q) \subset \mathfrak{A}_2$.*

Proof. $\Sigma(Q)$ is convex because it is the closure of a convex set. Since $\|L(Q)\| \leq \lambda^{-2}\|K\|^2\|Q\|$, L is continuous. Since the convex hull of $\{L^n Q; n = 0, 1, 2, \dots\}$ is invariant under L , its closure is also invariant due to the boundedness of L . We now show that $\Sigma(Q)$ is compact.

From (6.18) and (6.19), we have

$$\alpha(L^n Q) = \alpha(\mathcal{L}^n Q) \leq b(\|\Phi\|)\alpha(Q), \quad (7.4)$$

$$\alpha_l(L^n Q) = \alpha_l(\mathcal{L}^n Q) \leq b_l,$$

$$b_l = \alpha(Q) b'(\|\Phi\|) \delta_{l-r}(\|\Phi\|/2) + b(\|\Phi\|)\alpha_l(Q), \quad (7.5)$$

where we have used $\alpha_{l+n}(Q) \leq \alpha_l(Q)$ (Eq. (3.17)). Since $\lim_{k \rightarrow \infty} \alpha_k(Q) = 0$ for any $Q \in \mathfrak{A}$, $\lim_{l \rightarrow \infty} b_l = 0$. Further $\nu(L^n Q) = \nu(Q)$. Therefore $\Sigma(Q)$ is compact due to Corollary 3.12.

Now assume $Q \in \mathfrak{A}_2$. Then $\sup l^{-1} \{ \log \alpha_l(Q) + (l/r) \log l \} < +\infty$ and hence

$$\begin{aligned} \overline{\lim}_{l \rightarrow \infty} l^{-1} \{ \log b_l + (l/r) \log l \} &= \overline{\lim}_{l \rightarrow \infty} l^{-1} \{ \log \max(\delta_{l-1}(\|\Phi\|/2), \alpha_l(Q)) \\ &+ (l/r) \log l \} < \infty. \end{aligned} \tag{7.6}$$

From (3.22) and $\|R^{-1}\|^{-1} \leq \|R\|$, we have

$$\|L^n Q\|_l \leq \|L^n Q\| b_l(1 - b_l)^{-1}, \tag{7.7}$$

where $v(L^n Q) = v(Q)$ and (7.4) implies

$$\|L^n Q\| \leq v(Q) \alpha(Q) b(\|\Phi\|). \tag{7.8}$$

(7.7) and (7.8) give a uniform bound for $L^n Q$, which is preserved in taking convex hull and closure. Thus $Q' \in \Sigma(Q)$ satisfies

$$\|Q'\|_l \leq v(Q) \alpha(Q) b(\|\Phi\|) b_l(1 - b_l)^{-1}. \tag{7.9}$$

By (7.6), we see that $Q' \in \mathfrak{A}_2$. Q.E.D.

Remark 7.4. $\Sigma(Q)$ is compact for any Q . This is because

$$Q = Q_1 - Q_2 + i(Q_3 - Q_4) \tag{7.10}$$

where $Q_1 = (Q + Q^*)/2 + 2\|Q\| \geq \|Q\|$, $Q_3 = i(Q^* - Q)/2 + 2\|Q\| \geq \|Q\|$ and $Q_2 = Q_4 = 2\|Q\|$. The estimates like (7.7) and (7.8) hold for each Q_j and hence $\Sigma(Q)$ is compact by Lemma 3.2.

Lemma 7.5. There exists $h \in \mathfrak{A}_2$ such that

$$Lh = h, v(h) = 1, \alpha(h) \leq b(\|\Phi\|). \tag{7.11}$$

Proof. $\Sigma(1)$ has a fixed point h under the mapping L by Lemma 7.3 and the Schauder-Tychonov theorem, and $h \in \mathfrak{A}_2$. Since $\alpha(1) = 1$, (7.4) implies $\alpha(h) \leq b(\|\Phi\|)$. Q.E.D.

Lemma 7.6. Let

$$EQ = v(Q) h. \tag{7.12}$$

Then

$$\lim_{n \rightarrow \infty} (q_x)^n \|L^n(1 - E)\|_{1,x} = 0 \tag{7.13}$$

for any $x > 1$ and some $q_x > 1$.

Proof. Note that $E^2 = E$. Since L and E are linear operators on $\mathfrak{A}_{M,x}$, it is enough to prove the convergence of

$$\lim_{n \rightarrow \infty} q_x^n \|L^n Q\|_{1,x} = 0 \tag{7.14}$$

uniformly in Q such that $\|Q\|_{1,x} \leq 1$ and $(1 - E)Q = Q$. The latter condition is the same as $v(Q) = 0$. Any Q can be decomposed as $Q = Q_1$

$+iQ_2$, $Q_1^* = Q_1$, $Q_2^* = Q_2$ and by the proof of Lemma 6.5, we have $\|Q_1\|_{M,x} \leq \|Q\|_{M,x} = 1$, $\|Q_2\|_{M,x} \leq \|Q\|_{M,x} = 1$. Further $v(Q) = v(Q_1) + iv(Q_2) = 0$ implies $v(Q_1) = v(Q_2) = 0$. Therefore it suffices to prove the uniform convergence of (7.14) for Q such that $Q^* = Q$, $v(Q) = 0$, $\|Q\|_{1,x} \leq 1$. We already know by Lemma 6.5 that L is a continuous map of $\mathfrak{A}_{M,x}$ into $\mathfrak{A}_{M,x}$. Let $Q' = Q + 2$, $A_0(Q') = Q'$ and

$$A_n(Q') = L^N A_{n-1}(Q') - [6b(\|\Phi\|)]^{-1} v(A_{n-1}(Q')), \quad (7.15)$$

$n = 1, 2, \dots$. We fix an N such that

$$N \geq N(a, M, x, \|\Phi\|), \quad N \geq M, \quad (7.16)$$

where $N(a, M, x, \|\Phi\|)$ is given in Lemma 6.4 (ii),

$$a = 2(6b(\|\Phi\|) - 1), \quad (7.17)$$

and M is chosen so as to satisfy

$$x^M \geq 2a, \quad (7.18)$$

$$\sum_{l=M}^{\infty} x^l b^l(\|\Phi\|) \delta_{l-r}(\|\Phi\|/2) \leq 1/10. \quad (7.19)$$

We now prove the following properties of $A_n(Q')$ and $A_n(2)$.

$$A_n(Q') \geq 0, \quad (7.20)$$

$$A_n(2) \geq 0, \quad (7.21)$$

$$A_n(Q') - A_n(2) = L^{nN} Q, \quad (7.22)$$

$$v(A_n(Q')) = 2\bar{q}_x^{-nN} = v(A_n(2)), \quad (7.23)$$

$$\bar{q}_x = [1 - \{6b(\|\Phi\|)\}^{-1}]^{-1/N}. \quad (7.24)$$

$$\|A_n(Q')\|_{M,x} \|A_n(Q')^{-1}\| \leq a, \quad (7.25)$$

$$\|A_n(2)\|_{M,x} \|A_n(2)^{-1}\| \leq a. \quad (7.26)$$

First consider $n = 0$. Since $v(Q') = 2 = v(2)$ due to $v(Q) = 0$, we have (7.23). Since $\|Q\| \leq \|Q\|_{M,x} \leq 1$, we have $1 \leq Q' \leq 3$. Hence (7.20) (7.21) holds. (7.22) is obvious. Since $\|Q'\|_{M,x} \leq 2 + \|Q\|_{M,x} \leq 3 \leq a$, we have (7.25). $\|2\|_{M,x} \|2^{-1}\| = 1 \leq a$.

Next assume (7.20) ~ (7.26) for $n = k - 1$ and consider (7.20) ~ (7.26) for $n = k$. By definition (7.15), (7.22) holds, where we use (7.23) for $n = k - 1$. Further, from (7.15),

$$v(A_k(Q')) = \{1 - [12b(\|\Phi\|)]^{-1}\} v(A_{k-1}(Q')) = 2\bar{q}_x^{-kN} \quad (7.27)$$

and the same holds when Q' is replaced by 2. This proves (7.23) for $n = k$.

From (7.16), (7.25) and Lemma 6.4 (ii), we have

$$\alpha(L^N A_{k-1}(Q')) \leq 3b(\|\Phi\|). \quad (7.28)$$

Therefore

$$\begin{aligned} L^N A_{k-1}(Q') &\geq [3b(\|\Phi\|)]^{-1} v(L^N A_{k-1}(Q')) \\ &= [3b(\|\Phi\|)]^{-1} v(A_{k-1}(Q')). \end{aligned} \quad (7.29)$$

From (7.29) and (7.15), we have

$$A_k(Q') \geq v(A_{k-1}(Q')) [6b(\|\Phi\|)]^{-1}. \quad (7.30)$$

Then same holds when Q' is replaced by 2. Hence we have (7.20) and (7.21) for $n = k$.

From (7.29), we have

$$[6b(\|\Phi\|)]^{-1} v(A_{k-1}(Q')) \|\{L^N A_{k-1}(Q')\}^{-1}\| \leq 1/2. \quad (7.31)$$

Therefore, by (3.26) and (7.28), we have

$$\alpha(A_k(Q')) \leq 2\alpha(L^N A_{k-1}(Q')) - 1 \leq 6b(\|\Phi\|) - 1. \quad (7.32)$$

From Lemma 6.4 (i),

$$\alpha_i(L^N A_{k-1}(Q')) \leq b_i, \quad (7.33)$$

$$b_l \equiv \alpha(A_{k-1}(Q')) b'(\|\Phi\|) \delta_{l-r}(\|\Phi\|/2) + b(\|\Phi\|) \alpha_{l+N}(A_{k-1}(Q')). \quad (7.34)$$

From (7.18) and (7.25) with $n = k - 1$, we have

$$\|A_{k-1}(Q')\|_{l+N} \|A_{k-1}(Q')^{-1}\| \leq x^{-M} a \leq 1/2 \quad (7.35)$$

for $l \geq M$. By (3.21) and (7.35),

$$\alpha_{l+N}(A_{k-1}(Q')) \leq 2 \|A_{k-1}(Q')\|_{l+N} \|A_{k-1}(Q')^{-1}\|. \quad (7.36)$$

Hence

$$\sum_{l=M}^{\infty} x^l \alpha_{l+N}(A_{k-1}(Q')) \leq 2x^{-N} \|A_{k-1}(Q')\|_{M+N,x} \|A_{k-1}(Q')^{-1}\|. \quad (7.37)$$

From (7.18), we have $2b(\|\Phi\|) x^{-M} \leq a^{-1} b(\|\Phi\|) \leq 10^{-1}$. Therefore, by (7.19), (7.34), (7.37) and (7.25), we have

$$\sum_{l=M}^{\infty} x^l b_l \leq a/10. \quad (7.38)$$

From (7.38), it follows for $l \geq M$

$$b_l \leq x^{-M} a/10 \leq 20^{-1}. \quad (7.39)$$

From (3.25), (7.31) and (7.39),

$$\alpha_l(A_k(Q')) \leq b_l \{1 - (1 + b_l)/2\}^{-1} \leq (40/19) b_l. \tag{7.40}$$

From (7.40), (7.39) and (3.22), we have

$$\|A_k(Q')\|_l \|A_k(Q')^{-1}\| \leq (40/19) b_l (1 - (40/19) b_l)^{-1} \leq (40/17) b_l. \tag{7.41}$$

By (7.38), (7.41), and (7.32), we have

$$\|A_k(Q')\|_{M,x} \|A_k(Q')^{-1}\| \leq a. \tag{7.42}$$

This proves (7.25) for $n = k$. The same calculation with Q' replaced by 2 yields (7.26). This completes the inductive proof of (7.20) ~ (7.26).

From (7.25) and (7.28), we have

$$\|A_n(Q')\|_{M,x} \leq a \|A_n(Q')^{-1}\| \leq av(A_n(Q')) \leq 2a\bar{q}_x^{-nN}. \tag{7.43}$$

Similarly,

$$\|A_n(2)\|_{M,x} \leq 2a\bar{q}_x^{-nN}. \tag{7.44}$$

From (7.22), (7.43) and (7.44), we have

$$\|L^{nN}Q\|_{M,x} \leq 4a\bar{q}_x^{-nN}. \tag{7.45}$$

Hence

$$\lim_{n \rightarrow \infty} \|L^{nN}Q\|_{M,x} q_x^{nN} = 0 \tag{7.46}$$

for any $q_x < \bar{q}_x$. This then implies, due to the boundedness of each L (Lemma 6.5),

$$\lim_{n \rightarrow \infty} \|L^{nN+k}Q\|_{M,x} q_x^{nN+k} = 0 \tag{7.47}$$

for $k = 0, 1, \dots, N - 1$. Hence we have

$$\lim_{n \rightarrow \infty} \|L^n Q\|_{M,x} q_x^n = 0. \tag{7.48}$$

Since $\|Q\|_l \leq \|Q\| \leq \|Q\|_{M,x}$ for $l = 1, \dots, M - 1$, we have

$$\|Q\|_{1,x} \leq M \|Q\|_{M,x}. \tag{7.49}$$

Therefore we have (7.13). Q.E.D.

§ 8. Gibbs States

Lemma 8.1. *Let $Q \in \mathfrak{A}_{1,x}$ for some x . Let φ_c be the central state of \mathfrak{A} . Let*

$$\bar{\varphi}_n^{n_1, n_2}(Q) = \varphi_c(Q \exp\{\Delta_1^- + \Delta_0^+ - U(1, n + n_1) - U(1 - n - n_2, 0)\}) \bar{Z}_n^{-1}, \tag{8.1}$$

$$\bar{Z}_n = \varphi_c(\exp\{\Delta_1^- + \Delta_0^+ - U(1, n + n_1) - U(1 - n - n_2, 0)\}). \tag{8.2}$$

Then

$$\lim_{n \rightarrow \infty} \bar{\varphi}_n^{n_1, n_2}(Q) = v(Q), \quad (8.3)$$

$$\lim_{n \rightarrow \infty} \frac{1}{2n} \log \bar{Z}_n = \frac{1}{2} \log \lambda. \quad (8.4)$$

Proof. By (5.8), we have

$$\begin{aligned} \exp(1/2) \{ \Delta_1^- + \Delta_0^+ - U(1, n + n_1) - U(1 - n - n_2, 0) \} \\ = K'_N \exp - (1/2) \{ H(1, N) + H(1 - N, 0) + H' \}, \end{aligned} \quad (8.5)$$

$$H' = H_+ + H_-, \quad (8.6)$$

$$H_+ = H(N + 1, n + n_1) + \Delta_{n+n_1}^+, \quad (8.7)$$

$$H_- = H(1 - n - n_2, -N) + \Delta_{-n-n_2}^-, \quad (8.8)$$

$$K'_N = K'_{N+} + K'_{N-}, \quad (8.9)$$

$$K'_{N+} = E_r(-\tau_s(N) \Psi/2; -\{H(1, N) + H_+\}/2), \quad (8.10)$$

$$K'_{N-} = E_r(-\tau_s(-N) \Psi/2; -\{H(1 - N, 0) + H_-\}/2), \quad (8.11)$$

where $n > N + r$. Let

$$\bar{\varphi}'_{n-N}(Q) = \varphi_c(Q' \exp - \tau_c(N) H') (\bar{Z}'_{n-N})^{-1}, \quad (8.12)$$

$$\bar{Z}'_{n-N} = \varphi_c(\exp - \tau_c(N) H'). \quad (8.13)$$

We then have

$$\bar{\varphi}_n^{n_1, n_2}(Q) = \bar{Z}_n^{-1} \bar{Z}'_{n-N} \bar{\varphi}'_{n-N} [\tau_c(N) p_N \varphi_N ((K'_N)^* Q K'_N)] \quad (8.14)$$

where p_n and φ_n are defined in Lemma 6.3 (iii).

We prove (8.3) for positive Q such that $2 \geq Q \geq 1$. (8.3) for a general Q will immediately follow from this case by linearity.

Let B be a constant such that

$$\|(K'_N)\| \leq B, \quad \|(K_N)\| \leq B, \quad \|(K_N)^{-1}\| \leq B. \quad (8.15)$$

Given ε , there exists $L_1(\varepsilon)$ such that for $n - N > L_1(\varepsilon)$

$$\|K'_N - K_N\| < \varepsilon. \quad (8.16)$$

We then have

$$\|\varphi_N((K'_N)^* Q K'_N) - \varphi_N(K_N^* Q K_N)\| \|\varphi_N(K_N^* Q K_N)^{-1}\| \leq 4B^3 \varepsilon. \quad (8.17)$$

There also exists $L_2(\varepsilon)$ such that for $N > L_2(\varepsilon)$

$$\|\lambda^{-N} \tau_c(N) p_N \varphi_N(K_N^* Q K_N) - v(Q) h\| < \varepsilon. \quad (8.18)$$

Since $\|\{\lambda^{-N}\tau_c(N) p_N \varphi_N(K'_N{}^* Q K_N)\}^{-1}\|^{-1} \leq \|h^{-1}\|^{-1} + \varepsilon$ by (8.18), we have

$$\begin{aligned} \|\lambda^{-N}\tau_c(N) p_N \varphi_N((K'_N{}^* Q K'_N) - v(Q) h)\| \|h^{-1}\| \\ \leq \{4B^3(1 + \varepsilon\|h^{-1}\|) + \|h^{-1}\|\} \varepsilon = \varepsilon_1. \end{aligned} \tag{8.19}$$

Similarly

$$\begin{aligned} \|\lambda^{-N}\tau_c(N) p_N \varphi_N((K'_N{}^* K'_N) - h)\| \|h^{-1}\| \\ \leq \{2B^3(1 + \varepsilon\|h^{-1}\|) + \|h^{-1}\|\} \varepsilon = \varepsilon_2. \end{aligned} \tag{8.20}$$

For any state φ_1 , we have

$$|\varphi_1(\lambda^{-N} p_N \tau_c(N) \varphi_N(K'_N{}^* Q K'_N) / [v(Q) \varphi_1(h)] - 1| \leq \varepsilon_1, \tag{8.21}$$

$$|\varphi_1(\lambda^{-N} p_N \tau_c(N) \varphi_N(K'_N{}^* K'_N) / [\varphi_1(h)] - 1| \leq \varepsilon_2. \tag{8.22}$$

Therefore for $n \geq L_1(\varepsilon) + L_2(\varepsilon)$,

$$|\overline{\varphi}_n^{n_1 n_2}(Q) [\overline{\varphi}_n^{n_1 n_2}(1) v(Q)]^{-1} - 1| < (\varepsilon_1 + \varepsilon_2)(1 - \varepsilon_2)^{-1}. \tag{8.23}$$

Since $\overline{\varphi}_n^{n_1 n_2}(1) = 1$ for all n , we have

$$\lim_{n \rightarrow \infty} \overline{\varphi}_n^{n_1 n_2}(Q) / v(Q) = 1. \tag{8.24}$$

We note that the convergence is uniform in n_1 and n_2 .

Next we prove (8.4). In (8.14), we set $Q = 1$, $n - N = L$. Given ε , we choose $L > L_1(\varepsilon)$, and for this L we choose $L_3(\varepsilon)$ such that $L_3(\varepsilon)^{-1} L < \varepsilon$, $L_3(\varepsilon)^{-1} |\log \overline{Z}'_L| < \varepsilon$. We then have, for $n - N = L$ and $N > \max(L_3(\varepsilon), L_2(\varepsilon))$,

$$\begin{aligned} |n^{-1} \log \overline{Z}_n - \log \lambda| \\ \leq n^{-1} |\log \overline{Z}'_L| + n^{-1} L |\log \lambda| \\ + n^{-1} |\log \overline{\varphi}'_L[\lambda^{-N} \tau_c(N) p_N \varphi_N(K'_N{}^* K'_N)]| \\ < (1 + \log \lambda) \varepsilon + L_3(\varepsilon)^{-1} |\log \overline{\varphi}'_L(h)| + L_3(\varepsilon)^{-1} \max\{\log(1 + \varepsilon_2), -\log(1 - \varepsilon_2)\}. \end{aligned} \tag{8.25}$$

Since $|\log \overline{\varphi}'_L(h)| \leq \max\{|\log \|h\|, |\log \|h^{-1}\|\},$ we have (8.4). Q.E.D.

Lemma 8.2. Let $Q \in \mathfrak{A}_0$ and

$$F = E_r(-\Psi/2; -\{H(-\infty, 0) + H(1, \infty)\}/2). \tag{8.26}$$

Then

$$\lim_{a \rightarrow -\infty, b \rightarrow \infty} \varphi_{ab}^G(Q) = v(F^* Q F) / v(F^* F), \tag{8.27}$$

$$\lim_{b \rightarrow a \rightarrow \infty} (b - a)^{-1} \log Z(a, b) = \frac{1}{2} \log \lambda + \log d \tag{8.28}$$

where φ_{ab}^G and $Z(a, b)$ are given by (2.2) and (2.3).

Proof. Let $a < -r, b > r$ and

$$F_{ab} = E_r(-\Psi/2; \{\Delta_0^+ + \Delta_1^- - U(a, 0) - U(1, b)\}/2) \quad (8.29)$$

Then

$$\varphi_{ab}(Q) = \bar{\varphi}_n^{n_1 n_2}(F_{ab}^* Q F_{ab}) / \bar{\varphi}_n^{n_1 n_2}(F_{ab}^* F_{ab}) \quad (8.30)$$

where $a = 1 - (n + n_2), b = n + n_1$. By a variance of Lemma 5.2,

$$\lim_{n \rightarrow \infty} \|F_{ab} - F\| = 0 \quad (8.31)$$

uniformly in n_1 and n_2 . Furthermore $\|F^{-1}\| < \infty$ and hence $v(F^* F)^{-1} \leq \|F^{-1}\|^2 < \infty$. By Lemma 8.1, we obtain (8.27).

To prove (8.28), we note the formula

$$Z(a, b) = \bar{Z}_n \bar{\varphi}_n^{n_1 n_2}(F_{ab}^* F_{ab}) d^{(2n + n_1 + n_2)}, \quad (8.32)$$

if $a = 1 - n - n_2, b = n + n_1$. Since $\bar{\varphi}_n^{n_1 n_2}(F_n^* F_n)$ converges to a non zero constant $v(F^* F)$, we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{1}{2n} \log Z(-(n + n_1), n + n_2) \\ = \lim_{n \rightarrow \infty} \frac{1}{2n} \log \bar{Z}_n + \log d = \frac{1}{2} \log \lambda + \log d. \end{aligned} \quad (8.33)$$

Since $Z(a, b)$ depends only on $b - a$, we have (8.28). Q.E.D.

This lemma proves Theorem 2.1 except for the analyticity.

Lemma 8.3. $\varphi_\Phi(Q)$ is lattice translation invariant.

Proof. The following two quantities coincide for $n > 0$.

$$v(F^* F) \varphi_\Phi(Q) = \lim_{N \rightarrow \infty} \bar{\varphi}_N^{0, n}(Q), \quad (8.34)$$

$$v(F^* F) \varphi_\Phi(\tau_s(n) Q) = \lim_{N \rightarrow \infty} \bar{\varphi}_N^{n, 0}(\tau_s(n) Q). \quad (8.35)$$

Hence $\varphi_\Phi(Q) = \varphi_\Phi(\tau_s(n) Q)$. Q.E.D.

Lemma 8.4. $\varphi_\Phi(Q)$ is time translation invariant.

Proof. We have

$$\varphi_{a,b}^G([U(a, b), Q]) = 0. \quad (8.36)$$

Hence

$$\varphi_\Phi([\delta(U(Z)) Q]) = 0. \quad (8.37)$$

Hence $\varphi_\Phi(\tau_T(t) Q) = 0$. Q.E.D.

Lemma 8.5. $\varphi_\Phi(Q)$ satisfies the KMS condition.

Proof. Let $Q_1 \in \mathfrak{A}_0, Q_2 \in \mathfrak{A}_0$. We have

$$\varphi_{a,b}^G(\{\exp \delta(U(a, b)) Q_2\} Q_1) = \varphi_{a,b}^G(Q_1 Q_2). \tag{8.38}$$

Hence we have

$$\varphi_\Phi(\{\exp \delta(U(Z)) Q_2\} Q_1) = \varphi_\Phi(Q_1 Q_2). \tag{8.39}$$

By continuity, it holds for any $Q_2 \in \mathfrak{A}_1$.

By Lemma 4.2, $\exp \delta(-sU(Z)) \tau_T(t) Q_2$ is holomorphic in $t + is$. Hence

$$\int \exp \delta(U(Z)) (\tau_T(t) Q_2) f_0(t) dt = \int \tau_T(t) Q_2 f_1(t) dt \tag{8.40}$$

where $f_\alpha(t) = \int_{-\infty}^{\infty} e^{-ist + as} f(s) ds$ is holomorphic in $t + i\alpha$. Note that $\tau_T(t) Q_2 \in \mathfrak{A}_1$. Hence we have $\varphi_\Phi(Q_1 Q_2 (f_0)) = \varphi_\Phi(Q_2 (f_1) Q_1)$ for $Q_1, Q_2 \in \mathfrak{A}_0$. By continuity, this equation holds for any $Q_1, Q_2 \in \mathfrak{A}$. Hence φ_Φ satisfies the KMS condition.

Lemma 8.6. $\varphi_\Phi(Q)$ has a uniform exponential clustering property.

Proof. Let $Q_1 \in \mathfrak{A}_1$. Given $\varepsilon > 0$, we prove the existence of N_ε such that for $N > N_\varepsilon$ and $Q_2 \in \mathfrak{A}(Z[-N, N])$, we have

$$|\varphi_\Phi(Q_1 Q_2) - \varphi_\Phi(Q_1) \varphi_\Phi(Q_2)| e^{qN} < \varepsilon \|Q_2\| \tag{8.41}$$

where q is some positive constant.

We first prove the corresponding property for v . Let $Q \in \mathfrak{A}_1, v(Q) = 0$. Let $x > 1$ and $q < (\log q_x)/2, q < \log x, q > 0$.

Since $v(Q') = v(L^n Q')$ and $\alpha(L^n Q') = \alpha(\mathcal{L}^n Q') \leq \alpha(Q') b(\|\Phi\|)$ by (6.18), we have $\|L^n Q'\| \leq \alpha(Q') b(\|\Phi\|) v(Q')$ if $Q' > 0$. If Q' is selfadjoint and $\|Q'\| \leq 1$, then $\|2 + Q'\| \leq 3, \alpha(2 + Q') \leq 3$. Hence $\|L^n(2 + Q')\| \leq 9b(\|\Phi\|)$. Similarly $\|L^n 1\| \leq b(\|\Phi\|)$. Hence $\|L^n Q'\| \leq 11b(\|\Phi\|)$. For general $Q', Q' = Q_1 + iQ_2, Q_1^* = Q_1, Q_2^* = Q_2, \|Q_1\| \leq \|Q'\|, \|Q_2\| \leq \|Q'\|$. Hence

$$\|L^n(Q')\| \leq 22b(\|\Phi\|) \|Q'\| \tag{8.42}$$

for any $Q' \in \mathfrak{A}$.

Let K_n be given by (6.6), $K_{n\pm}^{NN'}$ by (6.23), (6.24), $K_{(n,l)} = K_{n+}^{n-1, l-1} K_{n-}^{n-1, l-1}$. There exists $L_1(\varepsilon)$ such that for $l \geq L_1(\varepsilon)$

$$e^{2ql} \|K_{(n,l)} K_n^{-1} - 1\| \leq \varepsilon, \tag{8.43}$$

due to $\|K_{(n,l)} K_n^{-1} - 1\| \leq \|K_{(n,l)} - K_n\| \|K_n^{-1}\|$ and Lemma 5.2. Since $Q \in \mathfrak{A}_1$, there exists $L_2(\varepsilon)$ such that for $l \geq L_2(\varepsilon)$

$$e^{ql} \|Q - Q^{(l)}\| < \varepsilon, \quad Q^{(l)} \in \mathfrak{A}([-l, l]). \tag{8.44}$$

By Lemma 7.6, there exists $L_3(\varepsilon)$ such that for $l \geq L_3(\varepsilon)$

$$e^{2\varepsilon l} \|L^l Q\| \leq e^{2\varepsilon l} \|L^l Q\|_{1,x} < \varepsilon \quad (8.45)$$

due to $v(Q) = 0$.

We now have the following series of estimates. Let $N \geq L(\varepsilon) = \max(L_1(\varepsilon), L_2(\varepsilon), L_3(\varepsilon))$ and $Q_2 \in \mathfrak{A}(Z \setminus [-2N, 2N])$. By (8.43) and (8.44),

$$e^{2\varepsilon N} \|Q - (K_N^*)^{-1} K_{(N,N)}^* Q^{(2N)} K_{(N,N)} K_N^{-1}\| < \varepsilon', \quad (8.46)$$

$$e^{2\varepsilon N} \|Q Q_2 - (K_N^*)^{-1} K_{(N,N)}^* Q^{(2N)} Q_2 K_{(N,N)} K_N^{-1}\| < \varepsilon' \|Q\|, \quad (8.47)$$

$$\varepsilon' = \varepsilon \{ \|Q\| (2 + \varepsilon) + (1 + \varepsilon)^2 \}. \quad (8.48)$$

By (8.45), (8.46) and (8.42), we have

$$e^{-2\varepsilon N} (\varepsilon + 22b(\|\Phi\|)\varepsilon') > \|L^N \{(K_N^*)^{-1} K_{(N,N)}^* Q^{(2N)} K_{(N,N)} K_N^{-1}\}\|. \quad (8.49)$$

By (6.5) and $K_{(N,N)} \in \mathfrak{A}([-2N, 2N])$, we have

$$\begin{aligned} L^N \{(K_N^*)^{-1} K_{(N,N)}^* Q^{(2N)} K_{(N,N)} K_N^{-1}\} \tau_c(N) Q_2 \\ = L^N \{(K_N^*)^{-1} K_{(N,N)}^* Q^{(2N)} Q_2 K_{(N,N)} K_N^{-1}\}. \end{aligned} \quad (8.50)$$

By (8.49), (8.50), (8.47) and (8.42), we have

$$\|L^N(Q Q_2)\| e^{2\varepsilon N} < \|Q_2\| \{ \varepsilon + 44b(\|\Phi\|)\varepsilon' \}. \quad (8.51)$$

Hence, for $N > L([1 + 44b(\|\Phi\|)\{4 + 3\|Q\|\}]^{-1}\varepsilon)$, $\varepsilon < 1$, we have

$$|v(Q Q_2)| = |v(L^N(Q Q_2))| < \varepsilon e^{-2\varepsilon N} \|Q_2\| \quad (8.52)$$

for any $Q_2 \in \mathfrak{A}(Z \setminus [-2N, 2N])$.

For general Q , apply (8.52) for $Q - v(Q)$ and we obtain

$$|v(Q Q_2) - v(Q) v(Q_2)| < \varepsilon e^{-2\varepsilon N} \|Q_2\|. \quad (8.53)$$

We now apply (8.53) for $F^* Q Q_2 F$, and $F^* Q F$. Although one F is on the right of Q_2 , the same formula (8.53) holds for sufficiently large N because $\|[Q_2, F]\|$ has a similar bound due to Lemma 5.2. Therefore

$$|v(F^* Q Q_2 F) - v(F^* Q F) v(Q_2)| < \varepsilon e^{-2\varepsilon N} \|Q_2\|, \quad (8.54)$$

$$|v(F^* Q_2 F) - v(F^* F) v(Q_2)| < \varepsilon e^{-2\varepsilon N} \|Q\|. \quad (8.55)$$

By (8.27), we have

$$e^{2\varepsilon N} |\varphi_\Phi(Q Q_2) - \varphi_\Phi(Q) \varphi_\Phi(Q_2)| < \varepsilon v(F^* F)^{-1} (1 + \|Q\|) \|Q_2\|. \quad (8.56)$$

This proves the uniform exponential clustering property of φ_Φ . Q.E.D.

§ 9. Analyticity

Lemma 9.1. *If $\Phi \in \mathfrak{A}([0, r])$ and $Q \in \mathfrak{A}([0, N])$ are holomorphic function of $\zeta = (\zeta_1 \dots \zeta_n)$ in a domain D where r and N are fixed, then $E_r(Q; H(I))$ is holomorphic in ζ in D with respect to $\| \cdot \|_{1,x}, x > 1$ and $\| \cdot \|$.*

Proof. Let $\zeta^{(0)} \in D$ and $\Phi = \sum \Phi_m (\zeta - \zeta^{(0)})^m, Q = \sum Q_m (\zeta - \zeta^{(0)})^m, m = (m_1 \dots m_n), (\zeta - \zeta^{(0)})^m = \prod_j (\zeta_j - \zeta_j^{(0)})^{m_j}$. Then $\sum \| \Phi_m \| |\zeta - \zeta^{(0)}|^m \leq a_\Phi, \sum \| Q_m \| |\zeta - \zeta^{(0)}|^m \leq a_Q$ uniformly in ζ in a neighbourhood of $\zeta^{(0)}$.

We substitute these expansions into estimates in Theorem 4.2 and Lemma 5.2. The estimate there only uses the property $Q \in \mathfrak{A}([0, r])$ and their norms. Hence all estimates holds when $\| \Phi \|$ and $\| Q \|$ are replaced by a_Φ and a_Q . In particular, $E_r(Q; H(I)) = \sum E_m (\zeta - \zeta^{(0)})^m$ and

$$\sum_m \| E_m \| |\zeta - \zeta^{(0)}|^m \leq \exp \{ F_N(2a_\Phi) a_Q \}, \tag{9.1}$$

$$\sum_m \| E_m \|_{N+L} |\zeta - \zeta^{(0)}|^m \leq C'_N \delta_L(a_\Phi), \tag{9.2}$$

where C'_N now depends on a_Φ, a_Q and N . Therefore $E_r(Q; H(I))$ has a convergent power series expansion at $\zeta^{(0)}$ and is holomorphic in ζ .

Lemma 9.2. *Let \mathfrak{B} be a Banach space with a norm $\| Q \|$ for $Q \in \mathfrak{B}$. Let $\mathcal{L}(\zeta)$ be a bounded linear operator on \mathfrak{B} , holomorphic in $\zeta = (\zeta_1 \dots \zeta_n)$ in a neighbourhood D of a real point $\xi_0, \mathcal{L}(\xi) h(\xi) = \lambda(\xi) h(\xi)$ for real ξ in $D, h(\xi) \in \mathfrak{B}, \lambda(\xi) > 0$. Let v_ξ be in the dual \mathfrak{B}^* of $\mathfrak{B}, E_\xi Q = v_\xi(Q) h(\xi), v_\xi(h(\xi)) = 1, v_\xi(\mathcal{L}(\xi) Q) = \lambda(\xi) v_\xi(Q)$ for real ξ in D and $Q \in \mathfrak{B}$. Assume that there exists $\mu_\xi < \lambda(\xi)$ satisfying*

$$\lim_{N \rightarrow \infty} \mu_\xi^{-N} \| \mathcal{L}(\xi)^N (1 - E_\xi) \| = 0. \tag{9.3}$$

Let $Q_0 \in \mathfrak{B}$ be fixed and $v_{\xi_0}(Q_0) = 1$. Then there exist extensions $h(\zeta), \lambda(\zeta)$ and v_ζ for ζ in some neighbourhood D' of ξ_0 , such that $\lambda(\zeta)$ is a holomorphic function of $\zeta, h(\zeta)$ is a \mathfrak{B} valued holomorphic function of ζ and v_ζ is a \mathfrak{B}^ valued holomorphic function of ζ .*

Proof. The series

$$(Z - \mathcal{L}(\xi))^{-1} = (Z - \lambda(\xi))^{-1} E_\xi + \sum_{n=1}^{\infty} Z^{-n} \mathcal{L}(\xi)^{n-1} (1 - E_\xi) \tag{9.4}$$

is convergent for $|Z| > \mu_\xi, Z \neq \lambda(\xi)$, by (9.3) and is the inverse of $Z - \mathcal{L}(\xi)$. Let $\mu_1 > \mu_{\xi_0}, \lambda(\xi_0) - \mu_1 > 0$. If $|Z| \geq \mu_1$, we have

$$\| (Z - \mathcal{L}(\xi_0))^{-1} \| \leq |Z - \lambda(\xi_0)|^{-1} \| E_{\xi_0} \| + \sum_{n=1}^{\infty} \mu_1^{-n} \| \mathcal{L}(\xi_0)^{n-1} (1 - E_{\xi_0}) \|. \tag{9.5}$$

Let $\delta = \sum_{n=1}^{\infty} \mu_1^{-n} \|\mathcal{L}(\xi_0)^{n-1} (1 - E_{\xi_0})\|$. By Lemma 3.7, we have

$$(Z - \mathcal{L}(\xi_0) - \Delta)^{-1} = \sum_{n=1}^{\infty} \{(Z - \mathcal{L}(\xi_0))^{-1} \Delta\}^{n-1} (Z - \mathcal{L}(\xi_0))^{-1} \tag{9.6}$$

provided that

$$\|\Delta\| < (\delta + \delta')^{-1}, \|Z - \lambda(\xi_0)\| > \|E_{\xi_0}\| (\delta')^{-1}. \tag{9.7}$$

If Δ is a holomorphic function of ζ , (9.6) is holomorphic in ζ if $\|\Delta\| < (\delta + \delta')^{-1}$. (The uniform limit of a holomorphic function is holomorphic.) Let $S(\delta')$ be the circle of radius $2\|E_{\xi_0}\| (\delta')^{-1}$ with the center $\lambda(\xi_0)$. Let $\Delta = \mathcal{L}(\zeta) - \mathcal{L}(\xi_0)$. Define

$$E'_\zeta = (2\pi i)^{-1} \oint_{S(\delta')} (Z - \mathcal{L}(\xi_0) - \Delta)^{-1} dZ. \tag{9.8}$$

Provided that

$$2\|E_{\xi_0}\| (\delta')^{-1} \leq \lambda(\xi_0) - \mu_1, \|\Delta\| \leq (\delta + \delta')^{-1}, \tag{9.9}$$

we have from (9.4) and (9.8)

$$E'_\zeta = E_{\xi_0} \quad \text{if} \quad \zeta = \xi_0. \tag{9.10}$$

Since $(Z - \mathcal{L}(\xi_0) - \Delta)^{-1}$ is holomorphic in ζ as long as $Z \in S(\delta')$ and (9.9) holds, E'_ζ is holomorphic in ζ .

As is easily seen the dimension of $E'_\zeta \mathfrak{B}$ is continuous in ζ : If $\dim E'_a \mathfrak{B} < \dim E'_b \mathfrak{B}$, $\dim E'_a \mathfrak{B} < \infty$ then by an orthogonalization procedure there exists $\psi \in E'_b \mathfrak{B}$, $\psi \neq 0$ such that $E'_a \psi = 0$, which contradict the continuity. Therefore $\dim E'_\zeta \mathfrak{B} = \dim E_{\xi_0} \mathfrak{B} = 1$ as long as E'_ζ is holomorphic.

This then implies $\mathcal{L}(\zeta) E'_\zeta = \lambda'(\zeta) E'_\zeta$ because $\mathcal{L}(\zeta)$ commutes with E'_ζ . Since $\lambda'(\zeta) = v_{\xi_0}(\mathcal{L}(\zeta) E'_\zeta h(\xi_0)) v_{\xi_0}(E'_\zeta h(\xi_0))^{-1}$, $\lambda'(\zeta)$ is holomorphic in ζ as long as E'_ζ is holomorphic and $v_{\xi_0}(E'_\zeta h(\xi_0)) \neq 0$. The latter is guaranteed in a neighbourhood of $\zeta = \xi_0$ because $v_{\xi_0}(E_{\xi_0} h(\xi_0)) = 1$. Let $h'(\zeta) = E'_\zeta Q_0$. It is holomorphic and $h'(\zeta) = h(\xi_0)$ when $\zeta = \xi_0$. Finally, let $v'_\zeta(Q) = v_{\xi_0}(E'_\zeta Q) v_{\xi_0}(h'(\zeta))^{-1}$. It is holomorphic as long as $v_{\xi_0}(h'(\zeta)) \neq 0$, and $v'_\zeta(Q) = v_{\xi_0}(Q)$ when $\zeta = \xi_0$. Since $v_{\xi_0}(h(\xi_0)) = 1$, $v_{\xi_0}(h'(\zeta)) \neq 0$ in some neighbourhood of $\zeta = \xi_0$.

By (9.4), $\lambda(\xi)$ is the only singularity of $(Z - \mathcal{L}(\xi))^{-1}$ outside of a circle of radius μ_ξ . If ζ is in sufficiently small neighbourhood of ξ_0 so that $\|\Delta\|$ is small, then $\lambda'(\zeta)$ is the only singularity of $(Z - \mathcal{L}(\zeta))^{-1}$ outside of a circle of radius μ_1 . Hence $\lambda'(\xi) = \lambda(\xi)$ for real ξ in a neighbourhood of ξ_0 . The expansion (9.4) then proves $E'_\xi = E_\xi$, $h'(\xi) = h(\xi)$ and $v'_\xi = v_\xi$. Q.E.D.

$\mathcal{L}(\zeta) E'_\zeta = \lambda'(\zeta) E'_\zeta$ implies $\mathcal{L}(\zeta) h'(\zeta) = \lambda'(\zeta) h'(\zeta)$ and $\mathcal{L}(\zeta)^* v'_\zeta = \lambda'(\zeta) v'_\zeta$.

Lemma 9.3. $\varphi_\Phi(Q), Q \in \mathfrak{A}_{M,x}, x > 1$ and $P(\Phi)$ is holomorphic in Φ .

Proof. Let Φ be holomorphic in ζ and hermitian when ζ is real. Let

$$\mathcal{L}(\zeta) Q = \tau_c(1) d^{-2} \text{tr}_{[0,1]}(K(\zeta)^* Q K(\zeta)) \tag{9.11}$$

where $K(\zeta)$ is defined by (6.2)~(6.4) where Φ is now a holomorphic function of ζ . $K(\zeta) \in \mathfrak{A}_{M,x}$ for any $x > 1$ and is holomorphic with respect to $\| \cdot \|_{M,x}$ by Lemma 9.1. Since $\mathfrak{A}_{M,x}$ is a $*$ -Banach algebra, this implies that $K(\zeta)^* Q K(\zeta)$ is holomorphic in ζ with respect to $\| \cdot \|_{M,x}$. Since $\| \tau_c(1) d^{-2} \text{tr}_{[0,1]} Q \|_{1,x} \leq \| Q \|_{1,x}$, $\mathcal{L}(\zeta)$ is also holomorphic in ζ .

Now Lemma 9.2 is applicable for $\mathfrak{B} = \mathfrak{A}_{M,x}$ and $\mathcal{L}(\zeta)$. We see that λ, ν and h are holomorphic in ζ . $F \in \mathfrak{A}_{M,x}$ is also holomorphic in ζ . Hence $P(\Phi)$ and φ_Φ are also holomorphic in ζ . Q.E.D.

Remark 9.4. The present proof of analyticity is applicable to the one dimensional classical spin lattice with an exponentially decreasing potentials. For higher dimensional quantum lattice, the analyticity for low activity is proved in [6, 7, 8].

§ 10. Factor States and Extremal KMS States

Lemma 10.1. Let \mathfrak{A} be a C^* -algebra, $\mathfrak{A}_n \subset \mathfrak{A}, n \in Z, \mathfrak{A}(I)$ be the C^* -algebra generated by $\mathfrak{A}_n, n \in I, \mathfrak{A}(Z) = \mathfrak{A}, Q \in \mathfrak{A}_n$ commutes with $Q' \in \mathfrak{A}_{n'}$ for $n \neq n'$ and π be a representation of \mathfrak{A} such that $\pi(\mathfrak{A}(I))$ is a factor of type I for any finite I . Then $\pi(\mathfrak{A}(I))' \cap \pi(\mathfrak{A})'' = \pi(\mathfrak{A}(Z \setminus I))'$ for any finite I .

Proof. Let $\mathfrak{A}_0(I)$ be the $*$ -algebra generated by $\mathfrak{A}_n, n \in I$. Let u_{ij} be the matrix unit of $\pi[\mathfrak{A}(I)]$. Then $Q \in \pi(\mathfrak{A})''$ is written as $Q = \sum u_{ij} Q_{ij}, Q_{ij} = \sum_k u_{ki} Q u_{jk} \in \pi(\mathfrak{A}(I))'$. Since $\pi(\mathfrak{A})''$ is the σ -weak closure of $\pi(\mathfrak{A}_0(Z))$ and $Q_i^j \in \pi(\mathfrak{A}_0(Z \setminus I))$ for $Q' \in \pi(\mathfrak{A}_0(Z))$, we have $Q_{ij} \in \mathfrak{A}(Z \setminus I)$. If $Q \in \pi(\mathfrak{A}(I))'$, then $Q_{ij} = \delta_{ij} Q$. Q.E.D.

This is essentially Lemma 2.3 of [10]. This lemma for somewhat more general case follows from Lemma 3.2 of [4].

A state φ of A in Lemma 10.1 is said to be uniformly clustering if there exists finite I for given $\varepsilon > 0$ and $Q \in \mathfrak{A}$ such that

$$|\varphi(Q Q_1) - \varphi(Q) \varphi(Q_1)| < \varepsilon \|Q_1\| \tag{2.1}$$

for every $Q_1 \in \mathfrak{A}(Z \setminus I)$. This condition may be replaced by a number of equivalent conditions. We may require (2.1) for any given $Q \in \mathfrak{A}_0(Z)$ and for any $Q_1 \in \mathfrak{A}_0(Z \setminus I)$. If we denote the representation of \mathfrak{A} associated with the state φ by π_φ , then another equivalent condition is

$$|(\Psi_1, \pi_\varphi(Q_1) \Psi_2) - (\Psi_1, \Psi_2) \varphi(Q_1)| < \varepsilon \|Q_1\| \tag{2.2}$$

for any given Ψ_1, Ψ_2 . In fact (2.1) is a special case of (2.2) with $\Psi_1 = \pi_\varphi(Q^*) \Omega_\varphi, \Psi_2 = \Omega_\varphi$. On the other hand, (2.2) for a dense set of vectors $\Psi_1 = \pi_\varphi(Q_a^*) \Omega_\varphi, \Psi_2 = \pi_\varphi(Q_b) \Omega_\varphi, Q_a, Q_b \in \mathfrak{A}_0(Z)$ follows from (2.1) for sufficiently big I such that $Q_a, Q_b \in \mathfrak{A}(I)$. Hence (2.2) holds for every given Ψ_1 and Ψ_2 for sufficiently large N . The condition (2.2) is equivalent to (2.2) with the specialization $\Psi_1 = \Psi_2$.

Lemma 10.2. *Let \mathfrak{A} be as in Lemma 10.1 where $\pi = \pi_\varphi$ is canonically associated with a given state φ . Then $\pi_\varphi(\mathfrak{A})''$ is a factor if and only if φ is uniformly clustering.*

Proof. The only if part is in Lemma 4.12 of [1]. For the if part, any central element S of $\pi_\varphi(\mathfrak{A})''$ is in $\pi_\varphi(\mathfrak{A}(Z \setminus I))''$ for any I . Given ε , we choose I satisfying (2.2) and then $Q_1 \in \mathfrak{A}(Z \setminus I)$ such that

$$|(\Psi_1, (\pi_\varphi(Q_1) - S) \Psi_2)| < \varepsilon, \quad |(\Psi_1, \Psi_2) \{ \varphi(Q_1) - (\Omega_\varphi, S \Omega_\varphi) \}| < \varepsilon.$$

Then we have $|(\Psi_1, S \Psi_2) - (\Psi_1, \Psi_2) (\Omega_\varphi, S \Omega_\varphi)| < 3\varepsilon$. Since ε is arbitrary, $S = (\Omega_\varphi, S \Omega_\varphi) 1$. Q.E.D.

This is essentially Theorem 2.5 of [10]. It is used under slightly more general circumstances around Eq. (3.6) of [4]. Lemma 10.2 and Lemma 10.1 are also derived in [9] in connection with a characterization of pure phase in both classical and quantum statistical mechanics.

The central decomposition of states into factor states always exists and is unique [14]. If the state is a KMS state, then the factor states are KMS states at least if A is separable (Corollary 3.7[2]). Further Theorem 4.1 in [2] essentially implies, though not explicitly stated, the following theorem.

Theorem 10.3. *A KMS state is a factor state if and only if it is an extremal KMS state.*

Proof. Let the representation π_φ of a C^* -algebra \mathfrak{A} and a cyclic vector Ω_φ in the representation space \mathfrak{H}_φ of π_φ be canonically associated with a KMS state φ (i.e. $\varphi(Q) = (\Omega_\varphi, \pi_\varphi(Q) \Omega_\varphi)$). It is shown in [3] that the center of $\pi_\varphi(\mathfrak{A})''$ is elementwise invariant under time translation. For any central projection $F \neq 0, \varphi_F(Q) = (F \Omega_\varphi, \pi_\varphi(Q) F \Omega_\varphi) \|F \Omega_\varphi\|^{-2}$ is a KMS state ($\|F \Omega_\varphi\| \neq 0$ always) and is different from $\varphi(Q)$ unless $F = 1$. Hence if φ is not a factor state, $\varphi = \lambda \varphi_F + (1 - \lambda) \varphi_{1-F}, 0 < \lambda = \|F \Omega_\varphi\|^2 < 1$ and φ is not an extremal KMS state.

Conversely, let $\varphi = \lambda \varphi_1 + (1 - \lambda) \varphi_2, 0 < \lambda < 1$ where φ_1 and φ_2 are KMS states. There exists an operator $F \geq 0$ in $\pi_\varphi(\mathfrak{A})'$ such that $\lambda \varphi_1(Q) = (\Omega_\varphi, \pi_\varphi(Q) F \Omega_\varphi)$. Since $\varphi_1(\tau_T(t) Q) = \varphi_1(Q)$ by assumption, $(\pi_\varphi(Q_1) \Omega_\varphi, F \pi_\varphi(Q_2) \Omega_\varphi) = (\pi_\varphi(\tau_T(t) Q_1) \Omega_\varphi, F \pi_\varphi(\tau_T(t) Q_2) \Omega_\varphi)$, from which we have $F = U_\varphi(t)^* F U_\varphi(t)$. Namely $F \in R_1'$ in the notation of [1]. By the KMS

condition on φ_1 , we have $(\Omega_\varphi, \hat{Q}_1 \hat{Q}_2(f_0) F \Omega_\varphi) = (\Omega_\varphi, \hat{Q}_2(f_1) \hat{Q}_1 F \Omega_\varphi)$. This is assumed for $\hat{Q}_1 \in \pi_\varphi(\mathfrak{A})$ and hence it holds for $\hat{Q}_1 \in \pi_\varphi(\mathfrak{A})''$ by the weak closure. Rewriting the equation as $(\Omega_\varphi, \hat{Q}_1(f'_1) \hat{Q}_2 F \Omega_\varphi) = (\Omega_\varphi, \hat{Q}_2 \hat{Q}_1(f'_0) \Omega_\varphi)$ with $f'_0(t) = f_1(-t)$, we see that it also holds for $\hat{Q}_2 \in \pi_\varphi(\mathfrak{A})''$, if it is assumed for $\hat{Q}_2 \in \pi_\varphi(\mathfrak{A})$. Now we restrict our attention to $E_0 \mathfrak{H}_\varphi$ of t invariant vectors in \mathfrak{H}_φ . $\hat{\varphi}$ is a cyclic and separating trace of $E_0 \pi_\varphi(\mathfrak{A})'' E_0$, where $\hat{\varphi}(Q) = (\Omega_\varphi, Q \Omega_\varphi)$ for $Q \in \mathcal{B}(\mathfrak{H}_\varphi)$. This property should also hold for $\hat{\varphi}$ similarly defined from φ_1 . Namely $\hat{\varphi}_1$ is a cyclic and separating trace of $F E_0 \pi_\varphi(\mathfrak{A})'' F E_0$ in $F E_0 \mathfrak{H}_\varphi$. This implies that $F E_0$ must be in the center of $E_0 \pi_\varphi(\mathfrak{A})'' E_0$ by an easy calculation. The argument in the proof of Theorem 4.1 of [2] then shows that there must be a central element F_1 of $\pi_\varphi(\mathfrak{A})''$ such that $F E_0 = F_1 E_0$. Since $R'_1 \rightarrow R'_1 E_0$ is an isomorphism, $F = F_1$ and F is a non trivial central element of $\pi_\varphi(\mathfrak{A})''$. Namely $\pi_\varphi(\mathfrak{A})''$ is not a factor if φ is not an extremal KMS state. Q.E.D.

The decomposition of a KMS states into extremal KMS states coincides with the decomposition into extremal time translation invariant states if and only if π_φ is η abelian where η is taken as the mean $(2T)^{-1} \int_{-T}^T dT$ as $T \rightarrow \infty$. ([2].)

For the one dimensional quantum spin lattice, Lemma 10.2 and Theorem 10.3 are applicable and φ_ϕ is a factor state and is an extremal KMS state because it is uniformly clustering. The asymptotic abelian property relative to the lattice translation also implies that the factor state φ_ϕ is an extremal lattice translation invariant state.

Acknowledgement. The author thanks Director L. Motchane and the members of the Institut des Hautes Etudes Scientifiques for their kind hospitality and Dr. Ruelle for discussions.

References

1. Araki, H., and E. J. Woods: A complete Boolean lattice of type I factors. Publ. Res. Inst. Math. Sci. Kyoto Univ. Ser. A, **2**, 157—242 (1966).
2. —, and H. Miyata: On KMS boundary condition. Publ. Res. Inst. Math. Sci. Kyoto Univ. Ser. A, **4**, 373—385 (1968).
3. — Multiple time analyticity of a quantum statistical states satisfying the KMS boundary condition. Publ. Res. Inst. Math. Sci. Kyoto Univ. Ser. A, **4**, 361—371 (1968).
4. — A classification of factors II. Publ. Res. Math. Sci. Kyoto Univ. Ser. A, **4**, 585—593 (1968).
5. Fujiwara, I.: Operator calculus of quantized operator. Progr. Theoret. Phys. **7**, 433—448 (1952).
6. Galavotti, G., S. Miracle-Sole, and D. W. Robinson: Analyticity properties of the anisotropic Heisenberg model. Commun. Math. Phys. **10**, 311—324 (1968).
7. Ginibre, J.: Reduced density matrices of the anisotropic Heisenberg model. Commun. Math. Phys. **10**, 140—154 (1968).

8. Greenberg, W.: Correlation functionals of infinite volume quantum spin systems. *Commun. Math. Phys.* **11**, 314—320 (1969).
9. Landford III, O. E., and D. Ruelle: Observables at infinity and states with short range correlation in statistical mechanics. (preprint).
10. Powers, R. T.: Representations of uniformly hyperfinite algebras and their associated von Neumann rings. *Ann. of Math.* **86**, 138—171 (1967).
11. Robinson, D. W.: Statistical mechanics of quantum spin system II. *Commun. Math. Phys.* **7**, 337—348 (1968).
12. Ruelle, D.: Statistical mechanics of a one dimensional lattice gas. *Commun. Math. Phys.* **9**, 267—278 (1968).
13. Streater, R. F.: The Heisenberg ferromagnet as a quantum field theory. *Commun. Math. Phys.* **6**, 233—247 (1967).
14. Wils, W.: Désintégration centrale des formes positives des C^* -algebres. (preprint).

H. Araki
Research Institute for Mathematical Sciences
Kyoto University
Kyoto, Japan